

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

6-Amino-4-(3-iodoanilino)-2-methylpyrimidin-1-ium chloride

 Wafaa A. Zaghary,^a Detlef Geffken^b and Seik Weng Ng^{c,d*}

^aDepartment of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia, ^bInstitut für Pharmazie, Bundesstrasse 45 20146, Hamburg, Germany, ^cDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^dChemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203 Jeddah, Saudi Arabia
Correspondence e-mail: seikweng@um.edu.my


Received 19 June 2012; accepted 22 June 2012

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.006$ Å; R factor = 0.032; wR factor = 0.078; data-to-parameter ratio = 17.3.

In the cation of the title salt, $\text{C}_{11}\text{H}_{12}\text{IN}_4^+\cdot\text{Cl}^-$, the two aromatic rings are oriented to each other at 9.3 (2)°. In the crystal, the two independent Cl^- anions lie on twofold rotation axes. $\text{N}-\text{H}\cdots\text{Cl}$ hydrogen bonds between the cations and anions generate a supramolecular layer parallel to (010).

Related literature

For the synthesis of 6-amino-4-[(4-chlorophenyl)amino]-2-methylpyrimidine hydrochloride, see: Craveri & Zoni (1958). For the synthesis of the reactants, see: Dox (1941); Foldi *et al.* (1942).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{12}\text{IN}_4^+\cdot\text{Cl}^-$	$V = 1286.23$ (8) Å ³
$M_r = 362.60$	$Z = 4$
Orthorhombic, $Pnn2$	Mo $K\alpha$ radiation
$a = 12.6323$ (5) Å	$\mu = 2.68$ mm ⁻¹
$b = 19.8608$ (7) Å	$T = 100$ K
$c = 5.1267$ (2) Å	$0.25 \times 0.05 \times 0.03$ mm

Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2012)
 $T_{\min} = 0.554$, $T_{\max} = 0.924$

10009 measured reflections
2981 independent reflections
2833 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.032$
 $wR(F^2) = 0.078$
 $S = 1.07$
2981 reflections
172 parameters
5 restraints

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.51$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.70$ e Å⁻³
Absolute structure: Flack (1983), 1321 Friedel pairs
Flack parameter: -0.04 (3)

Table 1

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{N1}-\text{H1}\cdots\text{Cl1}$	0.88 (4)	2.48 (4)	3.325 (4)	162 (4)
$\text{N3}-\text{H3}\cdots\text{Cl2}$	0.88 (5)	2.23 (4)	3.096 (4)	168 (4)
$\text{N4}-\text{H4A}\cdots\text{Cl2}$	0.89 (4)	2.66 (4)	3.462 (4)	150 (3)
$\text{N4}-\text{H4B}\cdots\text{Cl1}^{\dagger}$	0.88 (2)	2.64 (3)	3.404 (4)	146 (4)

 Symmetry code: (i) $x, y, z + 1$.

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank the Deanship of Scientific Research and the Research Center of the College of Pharmacy, King Saud University, and the Ministry of Higher Education of Malaysia (grant No. UM.C/HIR/MOHE/SC/12) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5569).

References

- Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.
Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
Craveri, F. & Zoni, G. (1958). *Bol. Sci. Fac. Chim. Ind. Bologna*, **16**, 126–131.
Dox, A. W. (1941). *Organic Syntheses*, Collected Vol. 1, p. 5. New York: Wiley.
Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
Foldi, Z., Fodor, G. V., Demjen, I., Szekere, H. & Halmos, I. (1942). *Chem. Ber.* **75**, 755–763.
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2012). E68, o2235 [doi:10.1107/S1600536812028401]

6-Amino-4-(3-iodoanilino)-2-methylpyrimidin-1-ium chloride

Wafaa A. Zaghary, Detlef Geffken and Seik Weng Ng

Comment

The synthesis of 6-amino-4-[(3-iodophenyl)amino]-2-methylpyrimidine, which was synthesized with the intention of labeling it with ^{99m}Tc for a study of its biotransformation, requires a small amount of hydrochloric acid as catalyst. A mole of hydrochloric acid is incorporated into the final product, so that the compound is formally 6-amino-4-[(3-iodophenyl)amino]-2-methylpyrimidin-1-ium chloride (Scheme I).

Protonation occurs on the aromatic nitrogen atom that is *para* to the secondary amino substituent. The non-hydrogen atoms of the cation lie on an approximate plane (r.m.s. deviation 0.132 Å); the two aromatic rings were twisted by 9.3 (2)°. The secondary amino and the tertiary pyrimidinium N atoms each forms a hydrogen bonds to a chloride ion to generate a layer parallel to (0 1 0) (Table 1).

Experimental

A mixture of 4-amino-2-methyl-6-chloropyrimidine (1.55 g, 0.01 mol) and 3-iodoaniline (2.19, 0.01 mol) in absolute ethanol (10 ml) and drop of hydrochloric acid was refluxed for 12 h; 4-amino-2-methyl-6-chloropyrimidine was synthesized from two other reactants (Dox, 1941; Foldi *et al.*, 1942). The reaction mixture was cooled and poured onto ice water. The formed precipitate was filtered, washed with water and recrystallized from ethanol to give the title compound in 60% yield; m.p. 535–537 K. The synthesis duplicates that used for of 6-amino-4-[(4-chlorophenyl)amino]-2-methylpyrimidine, which also exists as a hydrochloride (Craveri & Zoni, 1958).


Refinement

Carbon-bound H-atoms were placed in calculated positions [C-H 0.95 to 0.98 Å, $U_{\text{iso}}(\text{H})$ 1.2 to 1.5 $U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation.

The amino H-atoms were located in a difference Fourier map, and were refined with a distance restraint of N–H 0.88±0.01 Å; their temperature factors were refined.

Computing details

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO* (Agilent, 2012); data reduction: *CrysAlis PRO* (Agilent, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).


Figure 1

Thermal ellipsoid plot (Barbour, 2001) of $C_{11}H_{12}N_4I^+ Cl^-$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

6-Amino-4-(3-iodoanilino)-2-methylpyrimidin-1-ium chloride

Crystal data

$C_{11}H_{12}IN_4^+ Cl^-$

$M_r = 362.60$

Orthorhombic, $Pnn2$

Hall symbol: $P\ 2\ -2n$

$a = 12.6323\ (5)\ \text{\AA}$

$b = 19.8608\ (7)\ \text{\AA}$

$c = 5.1267\ (2)\ \text{\AA}$

$V = 1286.23\ (8)\ \text{\AA}^3$

$Z = 4$

$F(000) = 704$

$D_x = 1.872\ \text{Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 5603 reflections

$\theta = 2.6\text{--}27.5^\circ$

$\mu = 2.68\ \text{mm}^{-1}$

$T = 100\ \text{K}$

Prism, colorless

$0.25 \times 0.05 \times 0.03\ \text{mm}$

Data collection

Agilent SuperNova Dual

diffractometer with an Atlas detector

Radiation source: SuperNova (Mo) X-ray

Source

Mirror monochromator

Detector resolution: $10.4041\ \text{pixels mm}^{-1}$

ω scan

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2012)

$T_{\min} = 0.554$, $T_{\max} = 0.924$

10009 measured reflections

2981 independent reflections

2833 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.036$

$\theta_{\max} = 27.6^\circ$, $\theta_{\min} = 2.6^\circ$

$h = -16 \rightarrow 11$

$k = -25 \rightarrow 25$

$l = -6 \rightarrow 6$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.032$

$wR(F^2) = 0.078$

$S = 1.07$

2981 reflections

172 parameters

5 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H atoms treated by a mixture of independent
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0455P)^2 + 0.098P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.51 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.70 \text{ e } \text{\AA}^{-3}$

Absolute structure: Flack (1983), 1321 Friedel
pairs

Flack parameter: $-0.04 (3)$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
II	0.859538 (17)	0.859281 (10)	0.49964 (10)	0.01803 (9)
Cl1	0.5000	0.5000	0.7100 (3)	0.0165 (3)
Cl2	1.0000	0.5000	1.8561 (3)	0.0189 (3)
N1	0.6726 (3)	0.62226 (17)	0.8346 (7)	0.0133 (7)
H1	0.616 (3)	0.597 (2)	0.815 (12)	0.030 (15)*
N2	0.8301 (3)	0.63625 (14)	1.0653 (6)	0.0115 (8)
N3	0.8681 (3)	0.56332 (17)	1.4102 (7)	0.0127 (7)
H3	0.911 (4)	0.551 (2)	1.537 (8)	0.032 (14)*
N4	0.7588 (3)	0.47680 (17)	1.5483 (7)	0.0158 (8)
H4A	0.809 (3)	0.471 (2)	1.667 (7)	0.019 (12)*
H4B	0.6921 (14)	0.465 (2)	1.559 (12)	0.037 (16)*
C1	0.7485 (3)	0.78043 (16)	0.4921 (11)	0.0132 (7)
C2	0.6725 (3)	0.7825 (2)	0.2979 (8)	0.0153 (8)
H2	0.6724	0.8175	0.1716	0.018*
C3	0.5960 (3)	0.7318 (2)	0.2927 (8)	0.0174 (9)
H3A	0.5418	0.7328	0.1644	0.021*
C4	0.5988 (3)	0.68030 (16)	0.4730 (9)	0.0138 (8)
H4	0.5464	0.6460	0.4668	0.017*
C5	0.6770 (3)	0.67774 (17)	0.6634 (8)	0.0116 (7)
C6	0.7534 (3)	0.72929 (18)	0.6783 (9)	0.0141 (8)
H6	0.8062	0.7292	0.8101	0.017*
C7	0.7378 (3)	0.60225 (16)	1.0287 (9)	0.0116 (8)
C8	0.7081 (3)	0.54705 (17)	1.1824 (8)	0.0126 (7)
H8	0.6438	0.5235	1.1515	0.015*
C9	0.7757 (3)	0.52821 (18)	1.3798 (8)	0.0121 (8)
C10	0.8926 (3)	0.61518 (19)	1.2527 (8)	0.0110 (7)
C11	0.9960 (4)	0.64830 (19)	1.3029 (9)	0.0190 (9)
H11A	0.9953	0.6938	1.2292	0.029*
H11B	1.0528	0.6221	1.2212	0.029*
H11C	1.0083	0.6509	1.4914	0.029*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
I1	0.01513 (14)	0.01341 (13)	0.02556 (15)	-0.00259 (7)	-0.00545 (16)	0.00610 (14)
C11	0.0112 (6)	0.0149 (6)	0.0234 (8)	-0.0017 (5)	0.000	0.000
C12	0.0187 (7)	0.0262 (7)	0.0117 (7)	0.0092 (5)	0.000	0.000
N1	0.0139 (17)	0.0138 (14)	0.0122 (16)	-0.0050 (14)	-0.0026 (14)	0.0015 (14)
N2	0.0101 (16)	0.0122 (14)	0.012 (2)	-0.0018 (11)	-0.0021 (13)	0.0017 (11)
N3	0.0130 (18)	0.0130 (15)	0.0121 (16)	0.0006 (12)	-0.0030 (13)	0.0017 (13)
N4	0.0135 (16)	0.0214 (15)	0.012 (2)	-0.0011 (12)	-0.0009 (14)	0.0067 (14)
C1	0.0133 (16)	0.0117 (14)	0.0146 (16)	0.0003 (12)	0.003 (2)	0.002 (2)
C2	0.014 (2)	0.0166 (18)	0.015 (2)	0.0006 (16)	-0.0014 (17)	0.0031 (16)
C3	0.016 (2)	0.0180 (18)	0.018 (2)	0.0008 (16)	-0.0055 (17)	0.0016 (17)
C4	0.0132 (16)	0.0123 (14)	0.016 (2)	0.0005 (12)	0.000 (2)	-0.0008 (18)
C5	0.0115 (18)	0.0116 (16)	0.0118 (18)	0.0008 (14)	0.0002 (16)	0.0019 (16)
C6	0.0126 (19)	0.0157 (17)	0.0139 (17)	0.0002 (14)	-0.0026 (16)	-0.0006 (16)
C7	0.0101 (16)	0.0120 (14)	0.013 (2)	0.0001 (11)	-0.0003 (17)	-0.0014 (17)
C8	0.0117 (19)	0.0122 (16)	0.0139 (18)	-0.0014 (14)	-0.0001 (16)	0.0026 (16)
C9	0.016 (2)	0.0103 (15)	0.0099 (18)	-0.0011 (14)	0.0015 (15)	-0.0004 (14)
C10	0.0123 (19)	0.0121 (18)	0.0085 (17)	0.0006 (15)	0.0004 (15)	0.0006 (15)
C11	0.016 (2)	0.022 (2)	0.020 (2)	-0.0062 (16)	-0.0060 (18)	0.0083 (17)

Geometric parameters (\AA , $^\circ$)

I1—C1	2.103 (3)	C2—H2	0.9500
N1—C7	1.351 (5)	C3—C4	1.380 (6)
N1—C5	1.410 (5)	C3—H3A	0.9500
N1—H1	0.88 (1)	C4—C5	1.390 (6)
N2—C10	1.313 (5)	C4—H4	0.9500
N2—C7	1.360 (5)	C5—C6	1.409 (5)
N3—C10	1.345 (5)	C6—H6	0.9500
N3—C9	1.369 (5)	C7—C8	1.401 (5)
N3—H3	0.88 (1)	C8—C9	1.375 (5)
N4—C9	1.354 (5)	C8—H8	0.9500
N4—H4A	0.88 (1)	C10—C11	1.485 (6)
N4—H4B	0.88 (1)	C11—H11A	0.9800
C1—C2	1.384 (6)	C11—H11B	0.9800
C1—C6	1.395 (6)	C11—H11C	0.9800
C2—C3	1.396 (6)		
C7—N1—C5	131.6 (3)	C4—C5—N1	116.0 (3)
C7—N1—H1	114 (4)	C6—C5—N1	124.1 (4)
C5—N1—H1	114 (4)	C1—C6—C5	117.5 (4)
C10—N2—C7	117.3 (3)	C1—C6—H6	121.2
C10—N3—C9	121.2 (3)	C5—C6—H6	121.2
C10—N3—H3	122 (3)	N1—C7—N2	118.5 (3)
C9—N3—H3	117 (3)	N1—C7—C8	118.8 (3)
C9—N4—H4A	115 (3)	N2—C7—C8	122.7 (4)
C9—N4—H4B	113 (4)	C9—C8—C7	117.4 (4)
H4A—N4—H4B	128 (5)	C9—C8—H8	121.3

C2—C1—C6	123.0 (3)	C7—C8—H8	121.3
C2—C1—I1	117.0 (3)	N4—C9—N3	116.5 (3)
C6—C1—I1	120.0 (3)	N4—C9—C8	125.2 (4)
C1—C2—C3	118.2 (4)	N3—C9—C8	118.3 (4)
C1—C2—H2	120.9	N2—C10—N3	123.0 (4)
C3—C2—H2	120.9	N2—C10—C11	121.0 (4)
C4—C3—C2	120.3 (4)	N3—C10—C11	116.0 (4)
C4—C3—H3A	119.8	C10—C11—H11A	109.5
C2—C3—H3A	119.8	C10—C11—H11B	109.5
C3—C4—C5	121.1 (3)	H11A—C11—H11B	109.5
C3—C4—H4	119.5	C10—C11—H11C	109.5
C5—C4—H4	119.5	H11A—C11—H11C	109.5
C4—C5—C6	119.9 (4)	H11B—C11—H11C	109.5
C6—C1—C2—C3	1.5 (7)	C5—N1—C7—C8	174.7 (4)
I1—C1—C2—C3	-178.2 (3)	C10—N2—C7—N1	-179.0 (4)
C1—C2—C3—C4	-1.8 (6)	C10—N2—C7—C8	0.4 (6)
C2—C3—C4—C5	0.3 (7)	N1—C7—C8—C9	-179.5 (4)
C3—C4—C5—C6	1.6 (6)	N2—C7—C8—C9	1.1 (6)
C3—C4—C5—N1	-179.2 (4)	C10—N3—C9—N4	179.8 (4)
C7—N1—C5—C4	175.8 (4)	C10—N3—C9—C8	0.3 (6)
C7—N1—C5—C6	-5.0 (7)	C7—C8—C9—N4	179.1 (4)
C2—C1—C6—C5	0.4 (7)	C7—C8—C9—N3	-1.4 (6)
I1—C1—C6—C5	-179.9 (3)	C7—N2—C10—N3	-1.6 (6)
C4—C5—C6—C1	-1.9 (6)	C7—N2—C10—C11	178.6 (4)
N1—C5—C6—C1	179.0 (4)	C9—N3—C10—N2	1.3 (6)
C5—N1—C7—N2	-5.8 (6)	C9—N3—C10—C11	-178.9 (4)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots C11	0.88 (4)	2.48 (4)	3.325 (4)	162 (4)
N3—H3 \cdots C12	0.88 (5)	2.23 (4)	3.096 (4)	168 (4)
N4—H4A \cdots C12	0.89 (4)	2.66 (4)	3.462 (4)	150 (3)
N4—H4B \cdots C11 ⁱ	0.88 (2)	2.64 (3)	3.404 (4)	146 (4)

Symmetry code: (i) *x*, *y*, *z*+1.