

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

4-Nitro-*N*-[(*E*)-thiophen-2-ylmethylidene]anilineAbdullah M. Asiri,<sup>a,‡</sup> Hassan M. Faidallah,<sup>a</sup> Seik Weng Ng<sup>b</sup> and Edward R. T. Tiekink<sup>b,\*</sup><sup>a</sup>Chemistry Department, Faculty of Science, King Abdulaziz University, PO Box 80203, Jeddah, Saudi Arabia, and <sup>b</sup>Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia

Correspondence e-mail: edward.tiekink@gmail.com


Received 22 June 2012; accepted 22 June 2012

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.040;  $wR$  factor = 0.111; data-to-parameter ratio = 16.4.

In the title compound,  $\text{C}_{11}\text{H}_8\text{N}_2\text{O}_2\text{S}$ , there is a twist in the molecule, with the dihedral angle between the five- and six-membered rings being  $31.77(9)^\circ$ . The nitro group is slightly twisted out of the plane of the benzene ring to which it is attached [ $\text{O}-\text{N}-\text{C}-\text{C}$  torsion angle =  $9.0(3)^\circ$ ]. The S and N atoms are *syn*. In the crystal, supramolecular layers parallel to  $(\bar{2}04)$  are formed by  $\text{C}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{N}$  interactions. These layers are connected into a three-dimensional architecture by  $\pi-\pi$  interactions occurring between centrosymmetrically related benzene rings [centroid-centroid distance =  $3.6020(11)$  Å].

## Related literature

For background to 2-substituted thiophenes, see: Kleemann *et al.* (2006). For a related structure, see: Asiri *et al.* (2012).



## Experimental

## Crystal data

 $\text{C}_{11}\text{H}_8\text{N}_2\text{O}_2\text{S}$  $M_r = 232.25$ Monoclinic,  $C2/c$  $a = 9.2754(5)$  Å $b = 11.9983(9)$  Å $c = 18.4996(13)$  Å $\beta = 92.772(6)^\circ$   
 $V = 2056.4(2)$  Å<sup>3</sup>  
 $Z = 8$   
Mo  $K\alpha$  radiation $\mu = 0.30$  mm<sup>-1</sup>  
 $T = 100$  K  
 $0.25 \times 0.15 \times 0.05$  mm

## Data collection

Agilent SuperNova Dual diffractometer with an Atlas detector  
Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2012)  
 $T_{\min} = 0.692$ ,  $T_{\max} = 1.000$ 8697 measured reflections  
2377 independent reflections  
1869 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.050$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.111$   
 $S = 1.04$   
2377 reflections145 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\max} = 0.29$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.31$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C1}-\text{H1}\cdots\text{O2}^{\text{i}}$	0.95	2.50	3.412 (2)	160
$\text{C2}-\text{H2}\cdots\text{N1}^{\text{ii}}$	0.95	2.62	3.556 (2)	169

Symmetry codes: (i)  $x - 1, -y + 1, z - \frac{1}{2}$ ; (ii)  $-x + \frac{1}{2}, y - \frac{1}{2}, -z + \frac{1}{2}$ .

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).

The authors are grateful to King Abdulaziz University for providing research facilities. The authors also thank the Ministry of Higher Education (Malaysia) for funding structural studies through the High-Impact Research scheme (UM.C/HIR/MOHE/SC/12).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5953).

## References

- Agilent (2012). *CrysAlis PRO*. Agilent Technologies, Yarnton, Oxfordshire, England.  
Asiri, A. M., Faidallah, H. M., Khan, K. A., Ng, S. W. & Tiekink, E. R. T. (2012). *Acta Cryst. E* **68**, o1026.  
Brandenburg, K. (2006). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.  
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.  
Kleemann, A., Engel, J. B., Kutscher, B. & Reichert, D. (2006). In *Pharmaceutical Substances*. New York, Stuttgart: Georg Thieme Verlag.  
Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.  
Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

‡ Additional correspondence author, e-mail: aasiri2@kau.edu.sa.

## supplementary materials

*Acta Cryst.* (2012). E68, o2288 [doi:10.1107/S1600536812028346]

**4-Nitro-*N*-[(*E*)-thiophen-2-ylmethylidene]aniline**

Abdullah M. Asiri, Hassan M. Faidallah, Seik Weng Ng and Edward R. T. Tiekink

**Comment**

Among the various useful properties exhibited by thiophenes are their biological activities (Kleemann *et al.*, 2006). In continuation of structural studies of thienyl derivatives (Asiri *et al.*, 2012), herein the crystal structure determination of the title compound, (4-nitrophenyl)thiophene-2-ylmethylene-amine (I), is described.

In (I), Fig. 1, the conformation about the N1=C5 bond [1.283 (2) Å] is *E*. A twist in the molecule is evident as seen in the value of the dihedral angle of 31.77 (9)° between the five- and six-membered rings. The major deviation from a planar torsion angle is -37.0 (3)° for C5—N1—C6—C11 indicating that the most significant twist occurs around the N1—C6 bond. The nitro group is slightly inclined with respect to the plane of the benzene ring to which it is attached as seen in the O2—N2—C9—C8 torsion angle of 9.0 (3)°. The S and N atoms are *syn*.

In the crystal packing, supramolecular layers parallel to (-2 0 4) are formed by C—H...O and C—H...N interactions, Table 1, which lead to 22-membered {...HC<sub>2</sub>H...ONC<sub>4</sub>N}<sub>2</sub> synthons (Fig 2). Layers aggregate to form a three-dimensional architecture by  $\pi$ — $\pi$  interactions occurring between centrosymmetrically related benzene rings [inter-centroid distance = 3.6020 (11) Å for symmetry operation 1 - *x*, 1 - *y*, 1 - *z*], Fig. 3.

**Experimental**


A mixture of thiophen-2-carboxaldehyde (1.1 g, 0.01 *M*) and *p*-nitroaniline (1.4 g, 0.0 1*M*) in ethanol (10 ml) was heated on a water bath for 30 min, the solid which separated out was filtered, dried and recrystallized from ethanol as yellow prisms; *M. pt*: 375–376 K. Yield: 96%.

**Refinement**


Carbon-bound H-atoms were placed in calculated positions [C—H = 0.95 Å,  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ ] and were included in the refinement in the riding model approximation.

**Computing details**

Data collection: *CrysAlis PRO* (Agilent, 2012); cell refinement: *CrysAlis PRO* (Agilent, 2012); data reduction: *CrysAlis PRO* (Agilent, 2012); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 2006); software used to prepare material for publication: *publCIF* (Westrip, 2010).


**Figure 1**

The molecular structure of (I) showing the atom-labelling scheme and displacement ellipsoids at the 50% probability level.



**Figure 2**

A view of the supramolecular layer in the  $(-2\ 0\ 4)$  plane in (I) mediated by  $C-H\cdots O$  and  $C-H\cdots N$  interactions shown as orange and blue dashed lines, respectively.

**Figure 3**

A view in projection down the  $b$  axis of the unit-cell contents of (I), showing the stacking of supramolecular layers. The C—H...O, C—H...N and C—H— $\pi$  interactions are shown as orange, blue and purple dashed lines, respectively.

#### 4-Nitro-*N*-[(*E*)-thiophen-2-ylmethylidene]aniline

##### Crystal data

$C_{11}H_8N_2O_2S$

$M_r = 232.25$

Monoclinic,  $C2/c$

Hall symbol:  $-C 2yc$

$a = 9.2754 (5) \text{ \AA}$

$b = 11.9983 (9) \text{ \AA}$

$c = 18.4996 (13) \text{ \AA}$

$\beta = 92.772 (6)^\circ$

$V = 2056.4 (2) \text{ \AA}^3$

$Z = 8$

$F(000) = 960$

$D_x = 1.500 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 2057 reflections

$\theta = 2.8\text{--}27.5^\circ$

$\mu = 0.30 \text{ mm}^{-1}$

$T = 100$  K  
Prism, yellow

$0.25 \times 0.15 \times 0.05$  mm

*Data collection*

Agilent SuperNova Dual  
diffractometer with an Atlas detector  
Radiation source: SuperNova (Mo) X-ray  
Source  
Mirror monochromator  
Detector resolution:  $10.4041$  pixels  $\text{mm}^{-1}$   
 $\omega$  scan  
Absorption correction: multi-scan  
(*CrysAlis PRO*; Agilent, 2012)

$T_{\min} = 0.692$ ,  $T_{\max} = 1.000$   
8697 measured reflections  
2377 independent reflections  
1869 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.050$   
 $\theta_{\max} = 27.6^\circ$ ,  $\theta_{\min} = 2.8^\circ$   
 $h = -11 \rightarrow 12$   
 $k = -13 \rightarrow 15$   
 $l = -22 \rightarrow 24$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.111$   
 $S = 1.04$   
2377 reflections  
145 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.049P)^2 + 1.4742P]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} < 0.001$   
 $\Delta\rho_{\max} = 0.29 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.31 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.22121 (5)	0.35279 (4)	0.24739 (3)	0.01985 (16)
O1	0.94206 (15)	0.49221 (12)	0.60866 (8)	0.0287 (4)
O2	0.92261 (17)	0.65565 (13)	0.55892 (8)	0.0330 (4)
N1	0.45609 (16)	0.40163 (13)	0.36388 (8)	0.0172 (3)
N2	0.88841 (17)	0.55649 (14)	0.56311 (9)	0.0213 (4)
C1	0.1448 (2)	0.24635 (16)	0.19692 (10)	0.0216 (4)
H1	0.0693	0.2567	0.1610	0.026*
C2	0.2041 (2)	0.14558 (16)	0.21402 (10)	0.0191 (4)
H2	0.1742	0.0776	0.1916	0.023*
C3	0.3151 (2)	0.15307 (15)	0.26874 (10)	0.0178 (4)
H3	0.3687	0.0909	0.2871	0.021*
C4	0.33686 (19)	0.26104 (15)	0.29257 (10)	0.0165 (4)
C5	0.44640 (19)	0.29889 (16)	0.34505 (10)	0.0168 (4)
H5	0.5128	0.2465	0.3661	0.020*

C6	0.56731 (19)	0.43424 (15)	0.41452 (10)	0.0156 (4)
C7	0.6258 (2)	0.54040 (16)	0.40547 (10)	0.0186 (4)
H7	0.5925	0.5852	0.3658	0.022*
C8	0.7316 (2)	0.58087 (16)	0.45373 (10)	0.0195 (4)
H8	0.7716	0.6530	0.4477	0.023*
C9	0.77767 (19)	0.51350 (16)	0.51104 (9)	0.0172 (4)
C10	0.7220 (2)	0.40827 (16)	0.52155 (10)	0.0186 (4)
H10	0.7565	0.3637	0.5612	0.022*
C11	0.6151 (2)	0.36890 (16)	0.47345 (10)	0.0190 (4)
H11	0.5742	0.2974	0.4805	0.023*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0209 (3)	0.0158 (3)	0.0223 (3)	0.00164 (19)	-0.00515 (19)	-0.00144 (18)
O1	0.0258 (8)	0.0328 (9)	0.0264 (8)	0.0027 (7)	-0.0107 (6)	0.0029 (6)
O2	0.0361 (9)	0.0281 (9)	0.0334 (9)	-0.0135 (7)	-0.0118 (7)	0.0019 (7)
N1	0.0164 (8)	0.0180 (8)	0.0169 (8)	-0.0001 (6)	-0.0022 (6)	-0.0005 (6)
N2	0.0182 (8)	0.0258 (10)	0.0196 (9)	-0.0014 (7)	-0.0017 (7)	-0.0019 (7)
C1	0.0191 (10)	0.0246 (11)	0.0203 (10)	-0.0014 (8)	-0.0060 (8)	-0.0028 (8)
C2	0.0213 (10)	0.0182 (10)	0.0175 (10)	-0.0024 (8)	-0.0015 (8)	-0.0021 (7)
C3	0.0190 (9)	0.0158 (10)	0.0183 (9)	0.0004 (7)	-0.0019 (7)	0.0011 (7)
C4	0.0160 (9)	0.0173 (9)	0.0160 (9)	0.0010 (7)	-0.0002 (7)	0.0003 (7)
C5	0.0161 (9)	0.0183 (10)	0.0161 (9)	0.0014 (8)	-0.0002 (7)	0.0023 (7)
C6	0.0133 (8)	0.0174 (9)	0.0160 (9)	0.0012 (7)	-0.0003 (7)	-0.0033 (7)
C7	0.0176 (9)	0.0203 (10)	0.0176 (9)	0.0011 (8)	-0.0008 (7)	0.0040 (7)
C8	0.0189 (9)	0.0181 (10)	0.0214 (10)	-0.0040 (8)	-0.0003 (8)	0.0006 (7)
C9	0.0124 (9)	0.0243 (10)	0.0148 (9)	-0.0006 (7)	-0.0009 (7)	-0.0035 (7)
C10	0.0202 (9)	0.0190 (10)	0.0165 (9)	0.0016 (8)	-0.0014 (7)	0.0008 (7)
C11	0.0220 (10)	0.0165 (9)	0.0184 (10)	-0.0003 (8)	-0.0001 (8)	-0.0005 (7)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

S1—C1	1.7150 (19)	C4—C5	1.444 (2)
S1—C4	1.7246 (18)	C5—H5	0.9500
O1—N2	1.230 (2)	C6—C11	1.398 (3)
O2—N2	1.235 (2)	C6—C7	1.397 (3)
N1—C5	1.283 (2)	C7—C8	1.382 (3)
N1—C6	1.415 (2)	C7—H7	0.9500
N2—C9	1.467 (2)	C8—C9	1.384 (3)
C1—C2	1.359 (3)	C8—H8	0.9500
C1—H1	0.9500	C9—C10	1.382 (3)
C2—C3	1.411 (3)	C10—C11	1.383 (3)
C2—H2	0.9500	C10—H10	0.9500
C3—C4	1.380 (3)	C11—H11	0.9500
C3—H3	0.9500		
C1—S1—C4	91.13 (9)	C4—C5—H5	119.3
C5—N1—C6	119.00 (16)	C11—C6—C7	119.63 (17)
O1—N2—O2	123.41 (16)	C11—C6—N1	123.71 (17)

O1—N2—C9	118.47 (16)	C7—C6—N1	116.58 (16)
O2—N2—C9	118.12 (16)	C8—C7—C6	120.68 (17)
C2—C1—S1	112.55 (15)	C8—C7—H7	119.7
C2—C1—H1	123.7	C6—C7—H7	119.7
S1—C1—H1	123.7	C7—C8—C9	118.21 (17)
C1—C2—C3	112.54 (17)	C7—C8—H8	120.9
C1—C2—H2	123.7	C9—C8—H8	120.9
C3—C2—H2	123.7	C10—C9—C8	122.53 (17)
C4—C3—C2	112.28 (17)	C10—C9—N2	118.91 (16)
C4—C3—H3	123.9	C8—C9—N2	118.56 (17)
C2—C3—H3	123.9	C9—C10—C11	118.88 (18)
C3—C4—C5	126.67 (18)	C9—C10—H10	120.6
C3—C4—S1	111.50 (14)	C11—C10—H10	120.6
C5—C4—S1	121.73 (14)	C10—C11—C6	120.05 (18)
N1—C5—C4	121.49 (17)	C10—C11—H11	120.0
N1—C5—H5	119.3	C6—C11—H11	120.0
C4—S1—C1—C2	0.33 (16)	N1—C6—C7—C8	177.85 (16)
S1—C1—C2—C3	-0.5 (2)	C6—C7—C8—C9	0.0 (3)
C1—C2—C3—C4	0.5 (2)	C7—C8—C9—C10	-0.1 (3)
C2—C3—C4—C5	-176.49 (17)	C7—C8—C9—N2	-179.10 (16)
C2—C3—C4—S1	-0.2 (2)	O1—N2—C9—C10	9.7 (3)
C1—S1—C4—C3	-0.06 (15)	O2—N2—C9—C10	-170.11 (18)
C1—S1—C4—C5	176.43 (16)	O1—N2—C9—C8	-171.17 (17)
C6—N1—C5—C4	-178.81 (16)	O2—N2—C9—C8	9.0 (3)
C3—C4—C5—N1	-179.32 (19)	C8—C9—C10—C11	-0.6 (3)
S1—C4—C5—N1	4.7 (3)	N2—C9—C10—C11	178.44 (16)
C5—N1—C6—C11	-37.0 (3)	C9—C10—C11—C6	1.3 (3)
C5—N1—C6—C7	146.04 (18)	C7—C6—C11—C10	-1.4 (3)
C11—C6—C7—C8	0.8 (3)	N1—C6—C11—C10	-178.30 (17)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C1—H1...O2 <sup>i</sup>	0.95	2.50	3.412 (2)	160
C2—H2...N1 <sup>ii</sup>	0.95	2.62	3.556 (2)	169

Symmetry codes: (i)  $x-1, -y+1, z-1/2$ ; (ii)  $-x+1/2, y-1/2, -z+1/2$ .