

**Benildo S. Cavada,<sup>a\*</sup>**  
**Rolando E. R. Castellón,<sup>a</sup>**  
**Georg G. Vasconcelos,<sup>a</sup>**  
**Bruno A. M. Rocha,<sup>a</sup> Gustavo A.**  
**Bezerra,<sup>a</sup> Henri Debray,<sup>b</sup> Plínio**  
**Delatorre,<sup>a,c</sup> Celso S. Nagano,<sup>a</sup>**  
**Marcos Toyama,<sup>d</sup> Vicente P. T.**  
**Pinto,<sup>a</sup> Frederico B. M. B.**  
**Moreno,<sup>e</sup> Fernanda Canduri<sup>e</sup> and**  
**Walter F. de Azevedo Jr<sup>e\*</sup>**

<sup>a</sup>Departamento de Bioquímica e Biologia Molecular, Universidade Federal do Ceará, Biomol-Lab, CEP 60451-970, Caixa Postal 6043, Fortaleza-Ceará, Brazil,

<sup>b</sup>Laboratoire de Chimie Biologique et Unité Mixte de Recherche du CNRS No. 8576, Université des Sciences et Technologies de Lille, Villeneuve d'Ascq, France, <sup>c</sup>Departamento de Ciências Biológicas, Universidade Regional do Cariri, Crato, CE 63105-000, Brazil,

<sup>d</sup>Departamento de Bioquímica, Instituto de Biologia, Universidade Estadual de Campinas (UNICAMP), Campinas, SP, Brazil, and

<sup>e</sup>Departamento de Física, UNESP, São José do Rio Preto, SP 15054-000, Brazil

Correspondence e-mail: bscavada@ufc.br, walterfa@ibilce.unesp.br

Received 29 March 2005

Accepted 29 July 2005

Online 31 August 2005

## Crystallization and preliminary X-ray diffraction analysis of a new chitin-binding protein from *Parkia platycephala* seeds

A chitin-binding protein named PPL-2 was purified from *Parkia platycephala* seeds and crystallized. Crystals belong to the orthorhombic space group  $P2_12_12_1$ , with unit-cell parameters  $a = 55.19$ ,  $b = 59.95$ ,  $c = 76.60$  Å, and grew over several days at 293 K using the hanging-drop method. Using synchrotron radiation, a complete structural data set was collected to 1.73 Å resolution. The preliminary crystal structure of PPL-2, determined by molecular replacement, presents a correlation coefficient of 0.558 and an  $R$  factor of 0.439. Crystallographic refinement is in progress.

### 1. Introduction

Chitin, a natural homopolymer composed of  $\beta(1-4)$ -linked  $N$ -acetylglucosamine (GlcNAc)<sub>*n*</sub>, is a major component of the exoskeleton of fungi (comprising up to 30% of fungal cell walls) and invertebrates. It is easily obtained from marine invertebrates, insects and algae (Patil *et al.*, 2000).

The complete enzymatic hydrolysis of chitin to free  $N$ -acetylglucosamine residues is performed by a chitinolytic system and is known to be a continuous reaction. Different organisms produce a wide variety of hydrolytic enzymes that exhibit different substrate specificities. Some of them are called chitinases, which are enzymes that catalyze the hydrolysis of chitin. These proteins are a large and diverse group of enzymes that differ in their molecular structure, substrate specificity and catalytic mechanism (Kasprzewska, 2003). Specificity for chitin oligosaccharide is not a feature that is exclusive to the chitinases. Proteins named 'chitin-binding lectins' or 'hevein-like lectins' also possess affinity for  $N$ -acetylglucosamine residues, but cannot catalyze the hydrolysis of chitin (Van Damme *et al.*, 1998).

Several chitinases have been found in plants (angiosperms and gymnosperms) and are present in diverse tissues. Most are expressed by stress factors such as infection. Plants use chitinases as a defence against pathogenic fungi, but the enzymes may also perform other functions (Peumans *et al.*, 2002). Some chitinases have industrial and agricultural applications, such as in the biocontrol of pathogenic fungi and insects, as a target for biopesticides and in the production of chitoooligosaccharides (Kasprzewska, 2003; Patil *et al.*, 2000).

Plant lectins with chitinase activity are poorly described in the literature. The acidic chitinase from *Brassica juncea* shows a structure that is distinct from those observed for chitinases studied previously. This difference is characterized by the presence of two chitin-binding sites (Zhao & Chye, 1999), which permit this protein to agglutinate cells and may provide an advantage over other chitinases in antimicrobial and antifungal activity (Chye *et al.*, 2005).

Many carbohydrate-binding proteins have been reported, in particular those purified from plants (Moreno *et al.*, 2004; Gadelha *et al.*, 2005). The majority are from the Leguminosae family and comprise lectins and chitinases from diverse sources. Legume lectins have been well studied as a model of carbohydrate recognition. In the subfamily Mimosoideae, however, apart from *Parkia platycephala* 2 (PPL-2), only the seed lectins from *P. speciosa* (Suvachittanont & Peutpaiboon, 1992), *P. javanica* (Utarabhand & Akkayanont, 1995), *P. platycephala* (Cavada *et al.*, 1997) and *P. discolor* (Cavada *et al.*, 2000) have been isolated and characterized in detail. Moreover, crystal structures are only available for *P. platycephala* lectin (PPL-1)



© 2005 International Union of Crystallography  
All rights reserved

in native form (PDB code 1zgr) and in complex with 5-bromo-4-chloro-3-indolyl- $\alpha$ -D-mannose (PDB code 1zgs).

Mass-spectrometric analysis indicates that the PPL-2 monomer is not glycosylated and contains six cysteine residues that are involved in three disulfide bonds; PPL-2 gives a main mass peak at 29 407. Functional analysis reveals that PPL-2 recognizes carbohydrates on red blood cells and agglutinates trypsin-treated rabbit erythrocytes (128 haemagglutinating units per millilitre). In addition, PPL-2 can hydrolyze  $\beta$ (1-4)-glycosidic linkages between 2-acetoamido-2-deoxy- $\beta$ -D-glucopyranoses present in chitin. The exact mechanism of glycoside hydrolysis has been described by Cavada *et al.* (2005) and this mechanism reveals an endochitinase activity to be associated with PPL-2 from the elution times found for the GlcNAc, (GlcNAc)<sub>2</sub> and (GlcNAc)<sub>3</sub> standards. Hence, PPL-2 is the first and is a remarkable chimerolectin from the Mimosoideae, with the dual property of hydrolyzing chitin and binding sugar moieties on red blood cells (Cavada *et al.*, 2005).

In order to establish the crystal structure of this new member of the chitin-binding proteins, this work reports the crystallization and preliminary X-ray diffraction analysis of a hevamine-like protein from *P. platycephala* seeds, named PPL-2, that has the ability to agglutinate cells and shows inhibitory effects in the growth of bacterial colonies and nematode-egg eclosion (Castellón, 2004; Cavada *et al.*, 2005).

## 2. Material and methods

### 2.1. Purification and crystallization

Soluble proteins were extracted from the seeds of *P. platycephala* Benth in an extraction solution (0.1 M HCl with 0.1 M NaCl). After centrifugation, the supernatant was neutralized with sodium hydroxide (NaOH) and the neutralized solution was submitted to precipitation with ammonium sulfate. The fraction 0/60 was resuspended in 0.05 M Tris-HCl buffer with 0.1 M NaCl pH 7.0 and exhaustively dialyzed against this buffer. The protein was purified by affinity chromatography on a Red-Sepharose CL-6B (23.0  $\times$  2.5 cm) column equilibrated with the same buffer; elution of the non-interacting material took place using the equilibration buffer and the protein was eluted with 0.05 M Tris-HCl with 3.0 M NaCl pH 7.0 and finally dialyzed against Milli-Q water and lyophilized (Castellón, 2004; Cavada *et al.*, 2005).

For crystallization trials, the purified lectin was dissolved at a concentration of 7.5 mg ml<sup>-1</sup> in Milli-Q water. Microcrystals of PPL-2 were grown in Linbro plates at 293 K by the vapour-diffusion/sparse-matrix method (Jancarik & Kim, 1991) in hanging drops using Crystal Screen from Hampton Research. The drops were composed

of equal volumes (3  $\mu$ l) of protein solution and reservoir solution [0.2 M ammonium acetate, 0.1 M trisodium citrate dehydrate pH 5.6 and 30% (w/v) polyethylene glycol 4000] and were equilibrated against 500  $\mu$ l reservoir solution. Microcrystals were seeded into a new drop containing the same crystallization solution and an equal protein volume.

### 2.2. X-ray data collection

X-ray diffraction data were collected at a wavelength of 1.4727 Å using a synchrotron-radiation source (CPr station, Laboratório Nacional de Luz Síncrotron, Campinas, Brazil) and a CCD detector (MAR Research) with a crystal-to-detector distance of 70.00 mm at a temperature of 100 K. To avoid freezing, crystals were soaked in a cryoprotectant solution containing 75% mother liquor and 25% glycerol. Using an oscillation range of 1.0° and an exposure time of 30 s per frame, 90 images were collected to a maximum resolution of 1.73 Å. Data were indexed, integrated and scaled using *MOSFLM* and *SCALA* (Collaborative Computational Project, Number 4, 1994).


### 2.3. Molecular replacement

Sequence-alignment analysis was performed using programs that compared the N-terminal sequence of PPL-2 with those of all the non-redundant bank of proteins deposited in the National Center of Biotechnology Information (NCBI). Local and multiple alignments were carried out using *BLAST* (Altschul *et al.*, 1990) and *CLUS-TALW* (Thompson *et al.*, 1994), respectively. To perform multiple alignments, plant chitinases from *Nicotiana tabacum*, *Phytolacca americana*, *Glycine max*, *Zea mays*, *Vitis vinifera*, *Arabidopsis thaliana*, *Vigna unguiculata* and *Hevea brasiliensis* were used.

The molecular-replacement method was used to determine the crystal structure of PPL-2 using the *AMoRe* software (Navaza, 1994). Rotation and translation functions were performed using data in the resolution range 15–3.0 Å. The best solution for each model was selected based on the magnitude of the correlation coefficient and the *R* factor. Four space groups were tested (*P*222, *P*2<sub>1</sub>2<sub>1</sub>2, *P*222<sub>1</sub> and *P*2<sub>1</sub>2<sub>1</sub>2<sub>1</sub>) using the hevamine protein coordinates (PDB code 1hvq, chain A; Terwisscha van Scheltinga *et al.*, 1996).

## 3. Results and discussion

Microcrystals grew in a month using condition No. 9 of Crystal Screen from Hampton Research; when submitted to the seeding experiment, a crystal cluster was observed (Fig. 1*a*) after another month. The drop was perturbed using a fine hair and after about 20 weeks suitable



**Figure 1**  
PPL-2 crystals. (a) Crystal cluster from PPL-2 seeding. (b) Crystals of PPL-2 diffracted to 1.73 Å.

**Table 1**

X-ray diffraction data collection.

Values in parentheses are for the highest resolution shell.

Wavelength (Å)	1.431
Space group	$P2_12_12_1$
Unit-cell parameters (Å)	$a = 55.19, b = 59.95, c = 76.70$
Resolution range (Å)	32.27–1.73
Unique reflections	25945
Completeness (%)	95.5 (95.5)
$I/\sigma(I)$	13.1 (2.4)
$R_{\text{sym}}$ (%)	4.0 (22.8)
$R_{\text{full}}$ (%)	3.6 (16.3)

PPL2	GGIVV-WGQNGEGTLTSTCESGLYQIVNIAFLSQFGGRRV	41
Hevamina	GGIAIYWGQNGEGTLTQTCTSTRKYSYVNI AFLNKFGNGQ	40
NtCHI	GDIVVVWGQDVGEGKLDTCNSGLYNI VNI AFLSSFG	37
PaCHI	GGIAIYWGQNGEGTLRDTCSGLYSYVNI GFLSTFGNG	39
GmCHI	G-IAVYWGQNGEGTLAEACNTGNYQYVNI AFLSTFGNG	38
ZmCHI	GNIAVYWGQNGEGSLADACNSGLYAYVNI AFLTTFGNG	39
VvCHI	GGIAIYWGQNGEGTLTQTCTNTGKYSYVNI AFLNKFGNG	39
AtCHI	GGIAIYWGQNGEGNLSATCATGRYAYVNI AFLVVKFGNGR	40
VuCHI	GGIAIYWGQNGEGTLSEACDTRGYTHVNI AFLNKFGNG	39
	* * : * * * : * * * : * : * * * : * * * *	

**Figure 2**

Multiple alignment of the N-terminal sequence of PPL-2 with those of plant chitinases. GmCHI, acidic chitinase from *G. max*; ZmCHI, acidic chitinase from *Z. mays*; Hevamina, chitinase/lysozyme from *H. brasiliensis*; VvCHI, precursor of acidic chitinase from *Vitis vinifera*; AtCHI, acidic endochitinase from *A. thaliana*; VuCHI, basic endochitinase type III from *Vigna unguiculata*; PaCHI, chitinase-B from *Phytolacca americana*; NtCHI, basic endochitinase type III from *N. tabacum*.

crystals were obtained (Fig. 1*b*). The best crystals grew to approximate dimensions of  $0.3 \times 0.2 \times 0.3$  mm.

Crystals of PPL-2 were grown by the hanging-drop vapour-diffusion method. PPL-2 crystals belong to the orthorhombic space group  $P2_12_12_1$ , with unit-cell parameters  $a = 55.19, b = 59.95, c = 76.70$  Å. The volume of the unit cell is  $253\,757.48$  Å<sup>3</sup>, which is compatible with one monomer in the asymmetric unit, with a  $V_M$  of  $2.3$  Å<sup>3</sup> Da<sup>-1</sup> (Matthews, 1968). A summary of the data-collection statistics is given in Table 1.

Sequence-alignment analysis permitted the retrieval of a viable search model to submit these data to molecular replacement. The N-terminal amino-acid sequence from PPL-2 (GGIVVWGQNGEGTLTSTCESGLYQIVNIAFLSQFGGRRV) is completely different from that found in PPL-1, a lectin isolated from seeds of *P. platycephala* (SLKGMISVGPWGSGGNYWSFKANHAITEI-VIHVKDNIKS; Cavada *et al.*, 1997). Based on local alignments of PPL-2, similarity has been found with chitinases, proteins that are reported to be related to defence mechanisms in plants. These alignments show that PPL-2 exhibits a high sequence similarity to type III chitinases. One of these proteins is hevamine, a chitinase and lysozyme protein found in latex from *H. brasiliensis*, which has the N-terminal sequence GGIAIYWGQNGEGTLTQTCTSTRKYSY-VNIAFLNKFGNGQ. Acidic chitinases extracted from the leaves of *G. max* (Watanabe *et al.*, 1999), *Z. mays* (Didierjean *et al.*, 1996) and *A. thaliana* (Kawabe *et al.*, 1997) also show similarity with PPL-2, as can be observed in Fig. 2.

The N-terminal alignment of PPL-2 and several chitinases demonstrated a degree of similarity estimated at 72% with basic endochitinase type III from *N. tabacum*, 71% with chitinase-B from *Phytolacca americana*, 71% with acidic chitinases from *G. max* and *Z. mays*, 66% with the precursor of the acidic chitinase (Precursor) from *Vitis vinifera*, 60% with acidic endochitinase from *A. thaliana*, 61% with basic endochitinase type III from *Vigna unguiculata* and 64% with hevamine, a chitinase/lysozyme from *H. brasiliensis*. Despite the high similarity observed among all the alignments, hevamine is the only one in which the polypeptidic fragment corre-

sponds to the N-terminal region as in PPL-2 and is thus a good search model for molecular replacement. The best results were obtained for space group  $P2_12_12_1$ , resulting in a correlation coefficient and *R* factor of 55.8 and 43.9%, respectively.

Initial crystallographic refinement was performed using rigid-body refinement followed by the maximum-likelihood method with the *REFMAC5* software (Collaborative Computational Project, Number 4, 1994), resulting in a model with an *R* factor of 29.7% and an  $R_{\text{free}}$  of 33.1%. Complete refinement of the structure of PPL-2 is in progress.

This work was partly financed by Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico (FUNCAP), Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Universidade Regional do Cariri (URCA), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Laboratório Nacional de Luz Síncrotron (LNLS), Campinas, Brazil and FAPESP (SMOLBNet, 01/07532-0). BSC and WFA are senior investigators of CNPq.

**References**

Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990). *J. Mol. Biol.* **215**, 403–410.

Castellón, R. E. R. (2004). Tese de Doutorado apresentada ao Departamento de Fitotecnia, Universidade Federal do Ceará, Fortaleza-Ceará, Brazil.

Cavada, B. S., Castellón, R. E. R., Vasconcelos, G. G., Delatorre, P., Nagano, C. S., Nascimento, K. S., Souza, E. P., Radis-Baptista, G., Leroy, Y., Toyama, M. H., Sampaio, A. H., Pinto, V. P. T. & Debray, H. (2005). Submitted.

Cavada, B. S., Madeira, S. V. F., Calvete, J. J., Sousa, L. A. G., Bomfim, L. R., Dantas, A. R., Lopes, M. C., Grangeiro, T. B., Freitas, B. T., Pinto, V. P. T., Leite, K. B. & Ramos, M. V. (2000). *Prep. Biochem. Biotech.* **30**, 271–280.

Cavada, B. S., Santos, C. F., Grangeiro, T. B., Silva, L. I. M. M., Campos, M. J. O., Sousa, F. A. M. & Calvete, J. J. (1997). *Physiol. Mol. Biol. Plants*, **3**, 109–115.

Chye, M. L., Zhao, K. J., He, Z. M., Ramalingam, S. & Fung, K. L. (2005). *Planta*, **220**, 717–730.

Collaborative Computational Project, Number 4 (1994). *Acta Cryst.* **D50**, 760–763.

Didierjean, L., Frendo, P., Nasser, W., Genot, G., Marivet, J. & Burkard, G. (1996). *Planta*, **199**, 1–8.

Gadelha, C. A. A., Moreno, F. B. M. B., Santi-Gadelha, T., Cajazeiras, J. B., Rustiguel, J. K. R., Freitas, B. T., Canduri, F., Delatorre, P. & de Azevedo, W. F. Jr (2005). *Acta Cryst.* **F61**, 87–89.

Jancarik, J. & Kim, S.-H. (1991). *J. Appl. Cryst.* **24**, 409–411.

Kasprzewska, A. (2003). *Cell Mol. Biol. Lett.* **8**, 809–824.

Kawabe, A., Innan, H., Terauchi, R. & Miyashita, N. T. (1997). *Mol. Biol. Evol.* **14**, 1303–1315.

Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.

Moreno, F. B. M. B., Delatorre, P., Freitas, B. T., Rocha, B. A. M., Souza, E. P., Facó, F., Canduri, F., Cardoso, A. L. H., Freire, V. N., Lima-Filho, J. L., Sampaio, A. H., Calvete, J. J., de Azevedo, W. F. Jr & Cavada, B. S. (2004). *Acta Cryst.* **D60**, 1493–1495.

Navaza, J. (1994). *Acta Cryst.* **A50**, 157–163.

Patil, R. S., Ghormade, V. V. & Deshpande, M. V. (2000). *Enzyme Microb. Technol.* **26**, 473–483.

Peumans, W. J., Proost, P., Swennen, R. L. & Van Damme, E. J. (2002). *Plant Physiol.* **130**, 1063–1072.

Suvachittanont, W. & Peutpaiboon, A. (1992). *Phytochemistry*, **31**, 4065–4070.

Terwisscha van Scheltinga, A. C., Hennig, M. & Dijkstra, B. W. (1996). *J. Mol. Biol.* **262**, 243–257.

Thompson, J. D., Higgins, D. G. & Gibson, T. J. (1994). *Nucleic Acids Res.* **22**, 4673–4680.

Utarabhand, P. & Akkayanont, P. (1995). *Phytochemistry*, **38**, 281–285.

Van Damme, J. M., Peumans, W. J., Barre, A. J. & Rougé, P. J. (1998). *Crit. Rev. Plant Sci.* **17**, 575–692.

Watanabe, A., Nong, V. H., Zhang, D., Arahira, M., Yeboah, N. A., Udaka, K. & Fukazawa, C. (1999). *Biosci. Biotechnol. Biochem.* **63**, 251–256.

Zhao, K. J. & Chye, M. L. (1999). *Plant Mol. Biol.* **40**, 1009–1018.