

Robert Serek,<sup>a</sup> Jade K. Forwood,<sup>a</sup>  
David A. Hume,<sup>a,b,c,d</sup> Jennifer L.  
Martin<sup>a,b,d</sup> and Bostjan Kobe<sup>a,b,d\*</sup>

<sup>a</sup>School of Molecular and Microbial Sciences, University of Queensland, Brisbane, Queensland 4072, Australia, <sup>b</sup>Institute for Molecular Bioscience, University of Queensland, Brisbane, Queensland 4072, Australia, <sup>c</sup>Cooperative Research Centre for Chronic Inflammatory Diseases, University of Queensland, Brisbane, Queensland 4072, Australia, and <sup>d</sup>Special Research Centre for Functional and Applied Genomics, University of Queensland, Brisbane, Queensland 4072, Australia

Correspondence e-mail: b.kobe@uq.edu.au

Received 25 November 2005

Accepted 3 January 2006

## Crystallization of the C-terminal domain of the mouse brain cytosolic long-chain acyl-CoA thioesterase

The mammalian long-chain acyl-CoA thioesterase, the enzyme that catalyses the hydrolysis of acyl-CoAs to free fatty acids, contains two fused 4HBT (4-hydroxybenzoyl-CoA thioesterase) motifs. The C-terminal domain of the mouse long-chain acyl-CoA thioesterase (Acot7) has been expressed in bacteria and crystallized. The crystals were obtained by vapour diffusion using PEG 2000 MME as precipitant at pH 7.0 and 290 K. The crystals have the symmetry of space group *R*32 (unit-cell parameters  $a = b = 136.83$ ,  $c = 99.82$  Å,  $\gamma = 120^\circ$ ). Two molecules are expected in the asymmetric unit. The crystals diffract to 2.4 Å resolution using the laboratory X-ray source and are suitable for crystal structure determination.

### 1. Introduction

Long-chain acyl-CoAs are intermediates in lipid metabolism and regulators of cellular processes including ion transport, vesicle trafficking, protein phosphorylation and gene expression (Faergeman & Knudsen, 1997; Hunt & Alexson, 2002; Yamada, 2005). Intracellular levels of acyl-CoAs are controlled by the ratio of synthesizing and hydrolysing activities. Acyl-CoA thioesterases (also known as acyl-CoA hydrolases; EC 3.1.2.1 and 3.1.2.2) catalyse the hydrolysis of acyl-CoAs to free fatty acids and CoA-SH. These enzymes are widely distributed among different tissues in mammals (Hunt & Alexson, 2002).

High levels of long-chain acyl-CoA hydrolysing activity (EC 3.1.2.2) have been detected in mouse and human brain (Anderson & Erwin, 1971; Yamada, 2005). Full-length cDNAs corresponding to six different isoforms were cloned from the human brain and shown to be derived from a single gene (Hunt *et al.*, 2005; Yamada *et al.*, 2002). When expressed in bacteria, four of the six isoforms showed acyl-CoA hydrolase activity, while two C-terminally truncated isoforms did not (Yamada *et al.*, 2002). Because several different groups have independently cloned the enzyme, several different names exist, including BACH (Kuramochi *et al.*, 2002; Yamada *et al.*, 1997, 1999), CTE-II (Engberg *et al.*, 1997) and ACT (Broustas *et al.*, 1996). The recently suggested revised nomenclature designates this protein as Acot7 (Hunt *et al.*, 2005). The mouse brain contains a 43 kDa Acot7 as the major isoform and other lesser isoforms including a 50 kDa isoform (Takagi *et al.*, 2004). The enzyme is specific for acyl-CoA hydrolysis, but has a broad chain-length specificity, hydrolysing acyl-CoAs with carbon numbers C6–C20 (Broustas & Hajra, 1995; Yamada *et al.*, 1994, 1996, 1999, 2002).

Acot7 contains two copies of the 4HBT domain (named after 4-hydroxybenzoyl-CoA thioesterase). The structure of 4HBT from *Pseudomonas* revealed a 'Hotdog' fold comprising a  $\beta$ -sheet 'bun' wrapped around an  $\alpha$ -helical 'sausage' (Benning *et al.*, 1998). This fold was first observed in the structure of *Escherichia coli*  $\beta$ -hydroxydecanoyl thiol ester dehydratase (FabA; Leesong *et al.*, 1996) and has since been found in a number of apparently unrelated proteins (Dillon & Bateman, 2004). The acyl-CoA thioesterases form a distinct class of Hotdog-fold proteins with little available structural information.

To shed light on the molecular function of long-chain acyl-CoA thioesterases, including the enzymatic mechanism and the specificity, we set out to determine the three-dimensional structure of mouse

**Table 1**

Diffraction data-collection statistics.

Values in parentheses are for the highest resolution shell.

Reservoir solution	0.1 M Tris, 20–30% PEG 2000 MME pH 7.0
Space group	R32
Unit-cell parameters	
$a = b$ (Å)	136.83
$c$ (Å)	99.82
$\gamma$ (°)	120
Resolution range (Å)	30.0–2.4 (2.48–2.40)
Observations	60405
Unique reflections	14175
Average redundancy	4.26 (4.19)
Completeness (%)	99.9 (99.6)
$R_{\text{merge}}$	0.05 (0.56)
Average $I/\sigma(I)$	12.7 (2.2)

†  $R_{\text{merge}} = \sum_{hkl} (\sum_i (I_{hkl,i} - \langle I_{hkl} \rangle)) / \sum_{hkl,i} I_{hkl,i}$ , where  $I_{hkl,i}$  is the intensity of an individual measurement of the reflection with Miller indices  $h, k$  and  $l$  and  $\langle I_{hkl} \rangle$  is the mean intensity of that reflection for  $I > -3\sigma(I)$ .

Acot7. As the first step towards this goal, we report here the crystallization and preliminary X-ray diffraction analysis of the C-terminal 4HBT domain of this protein.

## 2. Experimental methods

### 2.1. Expression and purification

The DNA encoding amino acids 160–338 of variant 1 of mouse (*Mus musculus*) Acot7 (GeneInfo Identification No. 19923052; Kuramochi *et al.*, 2002) was amplified by PCR using cDNA prepared from LPS-stimulated mouse macrophage cells as a template (Wells *et al.*, 2003). The reactions were catalysed by Triplemaster proofreading, blunt-ended polymerase mix (Eppendorf), using gene-specific primers at a concentration of 10 ng  $\mu\text{l}^{-1}$ . The PCR conditions were initial denaturation at 369 K for 4 min, followed by 25 cycles of 369 K for 30 s, 328 K for 30 s and 345 K for 1 min, and finally one cycle of 345 K for 5 min. The PCR product was cloned into the Gateway entry vector pENTR-D-TOPO (Invitrogen) following the manufacturer's instructions and transformed into chemically competent Top10 cells (Invitrogen) by heat-shock. Colonies were inoculated in 4 ml LB medium in 15 ml Falcon tubes and plasmid DNA was purified from these cultures using a DNA-extraction kit (Roche). Genes cloned into pENTR-D-TOPO were recombined into the expression vector pDEST-17 (Invitrogen) by the Gateway LR reaction following the manufacturer's instructions; this vector allows expression of a fusion protein containing an N-terminal hexahistidine tag (His tag). The resulting entry and expression vectors were initially assessed by digestion with the restriction enzymes *NotI* (entry vector) and *HindIII* and *BamHI* (expression vector) and confirmed by DNA sequencing.

The expression vector was used to transform the chemically competent *E. coli* strain BL21(DE3)pLysS by heat-shock. A single colony was inoculated into 20 ml Luria–Bertani (LB) medium and grown overnight at 310 K in the presence of ampicillin and then used to inoculate 2 l LB containing ampicillin. The culture was grown aerobically at 310 K until the  $\text{OD}_{600\text{nm}}$  reached  $\sim 1$  and induced with either 1 mM IPTG or 10–30 mM lactose. The temperature was reduced to room temperature upon induction and the culture was grown for a further 12 h. At harvest, the  $\text{OD}_{600\text{nm}}$  was typically  $\sim 5$ –6 and the yield of recombinant Acot7 was 7–8 mg per litre of culture. The His-tagged protein was purified by immobilized metal-affinity chromatography (IMAC; nickel-affinity gel, Sigma). After elution, the buffer was exchanged using a PD-10 desalting column (BioRad)

with 200 mM Tris pH 8.5, 150 mM NaCl, 1 mM DTT and the protein was diluted to  $\sim 3.5$  mg  $\text{ml}^{-1}$  for crystallization. The protein was  $>95\%$  pure as assessed by SDS–PAGE.

### 2.2. Crystallization


Prior to crystallization, the protein sample was supplemented with 0.25 mg  $\text{ml}^{-1}$  coenzyme A (CoA). Crystallization conditions were screened by the sparse-matrix approach using the hanging-drop vapour-diffusion technique (Jancarik & Kim, 1991; McPherson, 1982). 1  $\mu\text{l}$  protein solution was combined with 1  $\mu\text{l}$  reservoir solution and suspended over 0.1 ml reservoir solution. Small crystals were initially observed in 25% polyethylene glycol (PEG) 3350, 0.1 M Tris pH 5.5, 0.2 M ammonium acetate. Crystals could be grown in the pH range 5.5–8, with the best crystals obtained in 20–30% of either PEG 3350, 0.1 M Tris pH 7–8 or 20% PEG 2000 monomethylether (MME), 0.1 M Tris pH 7.0. SDS–PAGE of dissolved crystals confirmed that they were formed by the Acot7 fragment.

### 2.3. Diffraction data collection

For X-ray diffraction experiments, crystals were transiently soaked in a solution corresponding to the reservoir solution but supplemented with 15% glycerol and were cooled at 100 K in a nitrogen stream (Cryocool, Cryo Industries, New Hampshire, USA). Data were collected from single crystals using an R-AXIS IV<sup>++</sup> image-plate detector and Cu  $K\alpha$  radiation from a Rigaku FR-E rotating-anode generator (Rigaku/MSC, Texas, USA). Data were autoindexed and processed with the program *CrystalClear* (Rigaku/MSC; Table 1). A representative oscillation image is shown in Fig. 1.

## 3. Results and discussion

We obtained crystals of the protein corresponding to the C-terminal 4HBT domain of the mouse long-chain acyl-CoA thioesterase



**Figure 1**

Oscillation image (1°) of the crystal described in Table 1. The crystal-to-detector distance is 120 mm; the resolution at the edge of the image is 2.0 Å.

(Acot7/mBACH) using either PEG 3350 or PEG 2000 MME as the precipitant. The crystals are rod-shaped (dimensions  $0.2 \times 0.2 \times 0.1$  mm) and have the symmetry of space group *R*32. The crystals appear after a few days and grow to maximum dimensions within three weeks. There are likely to be two molecules of the protein in the asymmetric unit [assuming two molecules, the Matthews coefficient  $V_M$  (Matthews, 1968) and the solvent content are  $2.0 \text{ \AA}^3 \text{ Da}^{-1}$  and 38%, respectively]. The crystals diffract X-rays using the laboratory source to  $2.4 \text{ \AA}$  resolution. The data-collection statistics are shown in Table 1.

Size-exclusion chromatography suggests the protein expressed from our construct exists in a dimeric form (data not shown), which is consistent with known 4HBT domain structures (Dillon & Bateman, 2004). Likewise, the 43 kDa isoform showed a molecular weight of 100 kDa using gel filtration (Yamada *et al.*, 1996, 1999).

Determination of the crystal structure of this protein will provide important insights into the catalytic mechanism and regulation of the mammalian long-chain acyl-CoA thioesterases. The full-length protein contains two 4HBT domains fused as part of the same polypeptide (the N- and C-terminal domains share 27% sequence identity based on alignment of residues 18–171 and 180–330, respectively); it is currently unknown what the distinct activities of the two domains are and how they cooperate within the full-length proteins. Based on a spectrophotometric assay (Yamada *et al.*, 1994), our construct of the C-terminal 4HBT domain shows some palmitoyl-CoA hydrolyzing activity (J. K. Forwood, W. N. Meng, R. Serek and B. Kobe, unpublished results). While both 4HBT domains are predicted to have the structure corresponding to the Hotdog fold, no structures are currently available of any eukaryotic proteins containing this structural motif.

We thank Pawel Listwan for help with cloning and for critically reading the manuscript. This work was supported by the Australian Research Council (ARC; to JLM and BK); BK is a National Health

and Medical Research Council of Australia (NHMRC) Research Fellow and ARC Federation Fellow.

## References

- Anderson, A. D. & Erwin, V. G. (1971). *J. Neurochem.* **18**, 1179–1186.
- Benning, M. M., Wesenberg, G., Liu, R., Taylor, K. L., Dunaway-Mariano, D. & Holden, H. M. (1998). *J. Biol. Chem.* **273**, 33572–33579.
- Broustas, C. G. & Hajra, A. K. (1995). *J. Neurochem.* **64**, 2345–2353.
- Broustas, C. G., Larkins, L. K., Uhler, M. D. & Hajra, A. K. (1996). *J. Biol. Chem.* **271**, 10470–10476.
- Dillon, S. C. & Bateman, A. (2004). *BMC Bioinformatics*, **5**, 109.
- Engberg, S. T., Aoyama, T., Alexson, S. E., Hashimoto, T. & Svensson, L. T. (1997). *Biochem. J.* **323**, 525–531.
- Faergeman, N. J. & Knudsen, J. (1997). *Biochem. J.* **323**, 1–12.
- Hunt, M. C. & Alexson, S. E. (2002). *Prog. Lipid Res.* **41**, 99–130.
- Hunt, M. C., Yamada, J., Maltais, L. J., Wright, M. W., Podesta, E. J. & Alexson, S. E. (2005). *J. Lipid Res.* **46**, 2029–2032.
- Jancarik, J. & Kim, S.-H. (1991). *J. Appl. Cryst.* **24**, 409–411.
- Kuramochi, Y., Takagi-Sakuma, M., Kitahara, M., Emori, R., Asaba, Y., Sakaguchi, R., Watanabe, T., Kuroda, J., Hiratsuka, K., Nagae, Y., Suga, T. & Yamada, J. (2002). *Brain Res. Mol. Brain Res.* **98**, 81–92.
- Leesong, M., Henderson, B. S., Gillig, J. R., Schwab, J. M. & Smith, J. L. (1996). *Structure*, **4**, 253–264.
- McPherson, A. (1982). *Preparation and Analysis of Protein Crystals*. New York: John Wiley & Sons.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- Takagi, M., Kawabe, K., Suga, T. & Yamada, J. (2004). *Arch. Biochem. Biophys.* **429**, 100–105.
- Wells, C. A., Ravasi, T., Faulkner, G. J., Carninci, P., Okazaki, Y., Hayashizaki, Y., Sweet, M., Wainwright, B. J. & Hume, D. A. (2003). *BMC Immunol.* **4**, 5.
- Yamada, J. (2005). *Amino Acids*, **28**, 273–278.
- Yamada, J., Furihata, T., Iida, N., Watanabe, T., Hosokawa, M., Satoh, T., Someya, A., Nagaoka, I. & Suga, T. (1997). *Biochem. Biophys. Res. Commun.* **232**, 198–203.
- Yamada, J., Furihata, T., Tamura, H., Watanabe, T. & Suga, T. (1996). *Arch. Biochem. Biophys.* **326**, 106–114.
- Yamada, J., Kuramochi, Y., Takagi, M., Watanabe, T. & Suga, T. (2002). *Biochem. Biophys. Res. Commun.* **299**, 49–56.
- Yamada, J., Kurata, A., Hirata, M., Taniguchi, T., Takama, H., Furihata, T., Shiratori, K., Iida, N., Takagi-Sakuma, M., Watanabe, T., Kurosaki, K., Endo, T. & Suga, T. (1999). *J. Biochem. (Tokyo)*, **126**, 1013–1019.
- Yamada, J., Matsumoto, I., Furihata, T., Sakuma, M. & Suga, T. (1994). *Arch. Biochem. Biophys.* **308**, 118–125.