

Subhasree Roy Choudhury,<sup>a</sup>  
Aparna Gomes,<sup>b</sup> Antony  
Gomes,<sup>a</sup> Jiban K. Dattagupta<sup>c</sup>  
and Udayaditya Sen<sup>c\*</sup>

<sup>a</sup>Department of Physiology, University of Calcutta, 92 Acharya Prafulla Chandra Road, Kolkata 700 009, India, <sup>b</sup>New Drug Development Division, Indian Institute Of Chemical Biology, 4 Raja S. C. Mallick Road, Kolkata 700 032, India, and <sup>c</sup>Crystallography and Molecular Biology Division, Saha Institute of Nuclear Physics, 1/AF Bidhan Nagar, Kolkata 700 064, India

Correspondence e-mail:  
udayaditya.sen@saha.ac.in

Received 21 December 2005  
Accepted 17 February 2006

## Purification, crystallization and preliminary X-ray structural studies of a 7.2 kDa cytotoxin isolated from the venom of *Daboia russelli russelli* of the Viperidae family

A cytotoxin (MW 7.2 kDa) from Indian Russell's viper (*Daboia russelli russelli*) venom possessing antiproliferative activity, cardiotoxicity, neurotoxicity and myotoxicity has been purified, characterized and crystallized. The crystals belong to the tetragonal space group  $P4_1$ , with unit-cell parameters  $a = b = 47.94$ ,  $c = 50.2$  Å. Larger crystals, which diffracted to 1.5 Å, were found to be twinned; diffraction data were therefore collected to 2.93 Å resolution using a smaller crystal. Molecular-replacement calculations identified two molecules of the protein in the asymmetric unit, which is in accordance with the calculated  $V_M$  value.


### 1. Introduction

Snake-venom cytotoxins are typically small water-soluble proteins (60–74 amino-acid residues) that possess a diverse range of pharmacological activities including haemolysis, cardiotoxicity, neurotoxicity, depolarization of the membranes of excitable cells, membrane fusion and selective killing of certain type of tumour cells. Three-dimensional structure determinations of cytotoxins isolated from different venoms, mainly from those of cobras and kraits (Chen *et al.*, 2005; Dubovskii *et al.*, 2005; Lee *et al.*, 2005; Paaventhana *et al.*, 2003), have indicated that they all possess a common structure, popularly known as the three-fingered motif, which consists of  $\beta$ -sheets connected by four or five disulfide bridges with no helical content (Menez, 1998; Tsetlin, 1999). Despite the similarities in their backbone architecture, they show different biological activities even when isolated from the same venom source, which is believed to arise from subtle variations in their three-dimensional structures.

Although the functional profiles of cytotoxins from the venoms of *Daboia russelli siamensis*, *Echis coloratus*, *Cerastes cerastes* *etc.* from the Viperidae family have been studied, mostly on Yoshida sarcoma cells, KB cells, HeLa cells and EAC cell lines (Tu & Giltner, 1974; Maung-Maung *et al.*, 1995; Mady, 2002; Abu-Sinna *et al.*, 2003), no three-dimensional structure is presently available for any cytolytic and/or antitumour toxin isolated from a venom from the Viperidae family. Russell's viper (*D. russelli russelli*) is one of the most dangerous snakes prevalent in eastern India, accounting for thousands of deaths each year. Envenomation by Russell's viper causes oedema, haemolysis, haemorrhage, neurotoxicity, cardiotoxicity and kidney failure. Recently, we have purified and sequenced (20 N-terminal residues) a cytotoxin (hereafter referred to as *drCT-I*) from the venom of *D. russelli russelli*; its molecular weight was found to be 7.2 kDa. Together with cytotoxicity, *drCT-I* shows neurotoxicity, cardiotoxicity and myotoxicity. In addition, it exhibits antiproliferative activity in EAC and human cell lines. We have crystallized *drCT-I* and initiated its three-dimensional structure determination in order to address the diverse pharmacological functions of this protein. In this paper, we present the purification, crystallization and preliminary X-ray structural studies of *drCT-I*.

### 2. Purification

200 mg lyophilized venom was dissolved in 5 ml distilled water and after heat treatment at 353 K for 30 min to precipitate out the high-




© 2006 International Union of Crystallography  
All rights reserved


molecular-weight proteins present in the venom, the solution was centrifuged at 900g for 20 min and the supernatant was loaded onto a CM-cellulose (150 × 25 mm) column equilibrated with 0.02 M phosphate buffer pH 7.2. Fractions were eluted at a flow rate of 25 ml h<sup>-1</sup> with a stepwise gradient of NaCl (0.02, 0.05, 0.1, 0.2 and 0.5 M in 0.02 M phosphate buffer pH 7.2). For each NaCl gradient, 30 ml of solution was used and the desired toxin (*drCT-I*) was obtained at an NaCl concentration of 0.2 M (Fig. 1*a*). The purified *drCT-I* protein constitutes about 3.75% of the crude venom. The homogeneity of the protein was assessed by re-chromatography (Fig. 1*b*) on a C-18 Nova Pak column (7.2 × 300 mm) equilibrated with 100 mM phosphate buffer pH 7.0 at a flow rate of 0.5 ml min<sup>-1</sup> and also by SDS-PAGE (Fig. 2).

### 3. Crystallization


For crystallization, the protein was dialyzed against 10 mM Tris-HCl pH 7.2 containing 0.5 mM CaCl<sub>2</sub> and concentrated with Amicon Centriprep centrifugal filtration units (3000 Da molecular-weight cutoff) to 6 mg ml<sup>-1</sup>. Crystallization was performed by the hanging-




**Figure 1**  
(*a*) CM-cellulose chromatogram for the purification of *drCT-I* using a stepwise gradient of NaCl (0.02, 0.05, 0.1, 0.2 and 0.5 M). The peak containing *drCT-I* is indicated in the figure and is eluted at an NaCl concentration of 0.2 M. (*b*) HPLC profile of *drCT-I* in a C-18 Nova Pak column.



**Figure 2**  
15% SDS-PAGE of *drCT-I* samples. Lane 1, purified from a CM-cellulose column; lane 3, crystals dissolved in SDS-PAGE sample buffer; lane 2, molecular-weight markers (kDa).



(*a*)



(*b*)

**Figure 3**  
(*a*) Crystal of *drCT-I* (0.5 × 0.4 × 0.35 mm) grown in PEG 4000 at 277 K. (*b*) Diffraction image of the crystal of *drCT-I*; the arrow indicates reflections at 1.5 Å. Inset: a portion of the image (marked in square) is enlarged to show that the reflections are split (shown by arrow).

drop vapour-diffusion method using 24-well tissue-culture plates and initial trials for crystallizations were carried out using Crystal Screens I and II from Hampton Research. Typically, 2  $\mu\text{l}$  protein solution at 6  $\text{mg ml}^{-1}$  concentration was mixed with an equal volume of the screening solution and equilibrated over 700  $\mu\text{l}$  of the latter as reservoir solution. The best crystals (Fig. 3*a*) were obtained when 2  $\mu\text{l}$  protein solution was mixed with an equal volume of reservoir solution consisting of 0.1 *M* sodium acetate pH 4.6, 30% PEG 4000, 0.2 *M* ammonium sulfate and 6% glycerol and equilibrated at 277 K for 10 d.

## 4. Data collection and processing

Crystals of *drCT-I* grown in the presence of cryoprotectant were fished out from the crystallization drops and immediately flash-frozen in a stream of nitrogen (Oxford Cryosystem) at 100 K. X-ray diffraction data were collected to 2.93  $\text{\AA}$  resolution using a MAR Research 345 image-plate detector with Cu  $K\alpha$  radiation generated by a Bruker–Nonius FR591 rotating-anode generator equipped with Osmic MaxFlux confocal optics running at 50 kV and 90 mA. The larger crystals, which diffracted to 1.5  $\text{\AA}$ , showed a high degree of twinning (Fig. 3*b*), with twin fractions of between 0.25 and 0.35 as calculated using the online test for determining twinning in protein crystals (<http://nihserver.mbi.ucla.edu/Twinning>; Yeates, 1997). The smaller crystals diffracted weakly but were found to be less twinned, with twin fractions of  $\sim 0.1$ . The present data set was collected from such a smaller crystal with a low twin fraction. A total of 72 frames were collected with a crystal-to-detector distance of 290 mm. The exposure time for each image was 2 min and the oscillation range was maintained at 1°. Data were processed and scaled using the program suite *AUTOMAR* (<http://www.marresearch.com/automar/run/htm>). Data-collection and processing statistics are given in Table 1.

## 5. Results

The crystals belong to the tetragonal space group  $P4_1$ , with unit-cell parameters  $a = b = 47.94$ ,  $c = 50.2$   $\text{\AA}$ . The results of molecular-weight determination of *drCT-I* by MALDI–TOF and ESI–MS consistently produce a molecular weight of  $\sim 7.2$  kDa (see supplementary material<sup>1</sup>). Packing considerations, based on the molecular weight of 7.2 kDa, indicate the presence of two molecules in the asymmetric unit, corresponding to a Matthews coefficient ( $V_M$ ) of 1.99  $\text{\AA}^3 \text{Da}^{-1}$  and a solvent content of 37%. The N-terminal sequence of the protein (20 amino acids only) has been determined and shows highest homology with cobra cardiotoxin A3 (PDB code 1xt3; Lee *et al.*, 2005). Molecular-replacement calculations were performed by *AMoRe* (Collaborative Computational Project, Number 4, 1994) using the coordinates of cobra cardiotoxin A3 (PDB code 1xt3; Lee *et al.*, 2005) as the search model. Two molecules in the asymmetric unit produced a correlation coefficient of 67.5% and an *R* factor of 44.9%,

<sup>1</sup> Supplementary material has been deposited in the IUCr electronic archive (Reference: PU5127).

**Table 1**

Data-collection and data-processing parameters.

Values in parentheses are for the highest resolution shell.

Space group	$P4_1$
Unit-cell parameters ( $\text{\AA}$ )	$a = b = 47.94$ , $c = 50.2$
Oscillation range ( $^\circ$ )	1.0
Resolution ( $\text{\AA}$ )	2.93 (3.03–2.93)
No. of molecules per ASU	2
Matthews coefficient $V_M$ ( $\text{\AA}^3 \text{Da}^{-1}$ )	1.99
Solvent content (%)	37.0
No. of observations	8576
No. of unique reflections	2276
Mosaicity	0.38
Completeness (%)	95.2 (89.8)
$R_{\text{merge}}^\dagger$ (%)	10.8 (24.2)
$I/\sigma(I)$ for highest shell	2.1

<sup>†</sup>  $R_{\text{merge}} = \frac{\sum[\sum |I(h)_i - I(h)|]}{\sum I(h)}$ , where  $I(h)_i$  is the observed intensity of the  $i$ th measurement of reflection  $h$  and  $I(h)$  is the mean intensity of reflection  $h$  calculated after scaling.

with data in the resolution range 10–3.8  $\text{\AA}$ . Subsequent rigid-body refinement followed by 20 cycles of positional refinement using *CNS* v.1.0 (Brünger *et al.*, 1998) further reduced the *R* factor to 42.3% ( $R_{\text{free}} = 44.3\%$ ) with a correlation coefficient of 69.8% using data in the resolution range 10–3.0  $\text{\AA}$ . Further refinement of this structure and model building using the program *O* (Jones *et al.*, 1991) are under way.

SRC and US thank Dr J. Dasgupta and S. Khamrui for their help during crystallization and data collection.

## References

- Abu-Sinna, G., Esmat, A. Y., Al-Zahaby, A.-A. S., Soliman, N. A. & Ibrahim, T. M. (2003). *Toxicon*, **42**, 207–215.
- Brünger, A. T., Adams, P. D., Clore, G. M., DeLano, W. L., Gros, P., Grosse-Kunstleve, R. W., Jiang, J.-S., Kuszewski, J., Nilges, M., Pannu, N. S., Read, R. J., Rice, L. M., Simonson, T. & Warren, G. L. (1998). *Acta Cryst.* **D54**, 905–921.
- Chen, T. S., Chung, F. Y., Tjong, S. C., Goh, K. S., Huang, W. N., Chien, K. Y., Wu, P. L., Lin, H. C., Chen, C. J. & Wu, W. G. (2005). *Biochemistry*, **44**, 7414–7426.
- Collaborative Computational Project, Number 4 (1994). *Acta Cryst.* **D50**, 760–763.
- Dubovskii, P. V., Lesovoy, D. M., Dubinnyi, M. A., Konshina, A. G., Utkin, Y. N., Efremov, R. G. & Arseniev, A. S. (2005). *Biochem. J.* **387**, 807–815.
- Jones, T. A., Zou, J. Y., Cowan, S. W. & Kjeldgaard, M. (1991). *Acta Cryst.* **A47**, 110–119.
- Lee, S. C., Guan, H. H., Wang, C. H., Huang, W. N., Tjong, S. C., Chen, C. J. & Wu, W. G. (2005). *J. Biol. Chem.* **280**, 9567–9577.
- Mady, E. A. (2002). *J. Venom. Anim. Toxins*, **8**, 283–296.
- Maung-Maung, T., Gopallakrishnakone, P., Yuen, R. & Tan, C. H. (1995). *Toxicon*, **33**, 63–76.
- Menez, A. (1998). *Toxicon*, **36**, 1557–1572.
- Paaventhana, P., Joseph, J. S., Nirthanan, S., Rajaseger, G., Gopalakrishnakone, P., Kini, M. R. & Kolatkar, P. R. (2003). *Acta Cryst.* **D59**, 584–586.
- Tsetlin, V. I. (1999). *Eur. J. Biochem.* **264**, 281–286.
- Tu, A. & Giltner, J. B. (1974). *Res. Commun. Chem. Pathol. Pharmacol.* **9**, 783–786.
- Yeates, T. O. (1997). *Methods Enzymol.* **276**, 344–358.