

Acta Cryst. (1952). **5**, 285

Preliminary X-ray studies of crocidolite and amosite. By R. I. GARROD and C. S. RANN, *Defence Research Laboratories, Victoria, Australia*

(Received 8 November 1951)

During the course of an investigation into the properties of some fibrous silicates, it became necessary, in the absence of any published information at the time, to determine the unit-cell dimensions and space-group symmetry of crocidolite (ex West Australia) and amosite (ex Transvaal, South Africa), kindly supplied to us by James Hardie & Co. Pty. Ltd. In each material, fibres from the samples were found to consist of bundles of single crystals (individually too small for single-crystal techniques), oriented parallel or nearly so to the common fibre axis, but with random orientation in other directions. The degree of parallelism was found to be much closer for amosite than for crocidolite, which had a larger crystal size. Stationary X-ray photographs, with the incident beam normal to the axis of the fibres in the specimen, therefore approximate to a single-crystal rotation pattern, as found by Warren & Bragg for chrysotile (1930).

The crystal class was found to be monoclinic for both minerals; and, with the aid of a Bunn chart and appropriate reciprocal-lattice constructions, the unit cell dimensions were found to be:

Crocidolite: $a = 9.6_5$, $b = 18.3$, $c = 5.3_0$ Å, $\beta \approx 70^\circ$;
axial ratios 0.53:1:0.29.

Amosite: $a = 9.89$, $b = 18.26$, $c = 5.30$ Å, $\beta = 70^\circ$;
axial ratios 0.542:1:0.290.

(Owing to the poor parallelism of the crocidolite fibres the parameters for this mineral are approximate only.)

Both structures therefore belong to the monoclinic amphibole group. Absent reflexions are in accordance with the space group $I2/m$.

Shortly after we obtained the results for the West Australian crocidolite, the paper by Whittaker (1949) on the structure of Bolivian crocidolite reached us. In view of the similarity of the results obtained for the unit-cell dimensions, no further analysis on Australian crocidolite has been done.

Subsequent work on amosite has shown the unit cell to contain two molecules, with a formula corresponding to $(Ca Na Mn Fe Mg)_{6.09} (Si Al)_{7.99} O_{22}(OH)_2$ (probable error ± 0.02). Preliminary Fourier projections indicate a structure similar to (but differing in certain interesting details from) that proposed by Warren (1930) for tremolite and by Whittaker (1949) for crocidolite. Further refinement is at present in progress.

The authors are indebted to the Chief Scientist, Department of Supply, for permission to publish this communication.

References

- WARREN, B. E. (1930). *Z. Kristallogr.* **72**, 42.
WARREN, B. E. & BRAGG, W. L. (1930). *Z. Kristallogr.* **76**, 201.
WHITTAKER, E. J. W. (1949). *Acta Cryst.* **2**, 312.

Acta Cryst. (1952). **5**, 285

The crystal structure of kojic acid. By H. A. MCKINSTRY, P. F. EILAND and R. PEPINSKY, *The Pennsylvania State College, State College, Pennsylvania, U.S.A.*

(Received 6 September 1951 and in revised form 30 November 1951)

The structure of kojic acid, 2 hydroxy 5 hydroxymethyl 1,4 γ -pyrone, has been examined, in continuation of a program for X-ray investigation of sugars and related compounds. Weissenberg photographs were taken with Cu $K\alpha$ radiation. The cell is monoclinic, space-group $P2_1/n$, with

$$a = 3.85, \quad b = 18.4, \quad c = 8.55 \text{ \AA}, \quad \beta = 84^\circ.$$

The measured density of 1.58 g.cm.^{-3} gave $3.98 \approx 4$ molecules/cell. These results are in accord with previous measurements by Fox (1945). The molecule contains no asymmetric carbon atoms.

The a -axis periodicity indicated no that overlapping would occur in a projection on (100). The structure was consequently considered a good test subject for several recently-developed analytical methods.

The X-RAC method of approximation via positive kernels, as suggested by one of us (Pepinsky, 1951; Pepinsky & Cochran, 1951) for reduction of terms in the Fourier sum, was successful in giving the orientation of the molecule and its general position in the cell. The approximation technique was not successful in the next stages of the analysis. The application of the non-

negativity criterion on X-RAC was useful in all refinements (Eiland & Pepinsky, 1950).

The Harker-Kasper inequalities were inapplicable because of the low values of the unitary structure factors. There were sets of related factors large enough, and the cell was sufficiently thin, to suggest that the squared-crystal method (Sayre, 1951, 1952), and, in particular, the




Fig. 1. Kojic acid: projection on (100).