

Full-length article

Gene expression profile of amyloid beta protein-injected mouse model for Alzheimer diseaseLing-na KONG, Ping-ping ZUO¹, Liang MU, Yan-yong LIU, Nan YANG*Department of Pharmacology, School of Basic Medicine, Peking Union Medical College and Chinese Academy of Medical Sciences, Beijing 100005, China***Key words**

Alzheimer disease; dementia; amyloid beta-protein; cDNA microarray

¹ Correspondence to Prof Ping-ping ZUO.
Phn/Fax 86-10-6529-6404.
E-mail zuopp@public3.bta.net.cn

Received 2004-11-25

Accepted 2005-03-14

doi: 10.1111/j.1745-7254.2005.00129.x

Abstract

Aim: To investigate the gene expression profile changes in the cerebral cortex of mice injected icv with amyloid beta-protein (A β) fragment 25-35 using cDNA microarray. **Methods:** Balb/c mice were randomly divided into a control group and A β -treated group. The Morris water maze test was performed to detect the effect of A β -injection on the learning and memory of mice. Atlas Mouse 1.2 Expression Arrays containing 1176 genes were used to investigate the gene expression pattern of each group. **Results:** The gene expression profiles showed that 19 genes including *TBX1*, *NF- κ B*, *AP-1/c-Jun*, *cadherin*, *integrin*, *erb-B2*, and *FGFR1* were up-regulated after 2 weeks of icv administration of A β ; while 12 genes were down-regulated, including *NGF*, *glucose phosphate isomerase 1*, *AT motif binding factor 1*, *Na⁺/K⁺-ATPase*, and *Akt*. **Conclusions:** The results provide important leads for pursuing a more complete understanding of the molecular events of A β -injection into mice with Alzheimer disease.

Introduction

Alzheimer disease (AD) is the most common cause of dementia in the elderly, affecting 7%–10% of individuals over 65 years of age and approximately 40% of persons over 80 years old^[1]. The disease is characterized behaviorally by global cognitive decline, and defined histologically by two distinguishing pathologies: amyloid plaques, which are extracellular deposits consisting mainly of aggregated amyloid beta-protein (A β), and neurofibrillary tangles (NFT), which are intracellular deposits consisting predominantly of hyperphosphorylated tau protein^[2]. Epidemiological and molecular studies suggest that AD has multiple etiologies, including genetic mutations, susceptibility genes and environmental factors that promote formation and accumulation of insoluble A β and hyperphosphorylated tau. An animal model that mimics the progression of AD was developed using icv injection of A β into mice^[3–5]. This was accompanied by impairment in learning and memory in addition to biochemical changes and neuronal degeneration.

Recently, a high-throughput technique was developed using cDNA microarrays, in which labeled RNA hybridizes

to DNA molecules attached at specific locations on the surface of microarray membrane^[6]. One of the most attractive applications of microarrays is in the study of differential gene expression in disease and animal models. Microarrays are extensively employed to study global changes in gene expression in post-mortem tissues, cultured cells, animal models, and in response to drug treatment. The expression pattern of a gene provides indirect information about function, drug target and cause of a disease. In this study we used cDNA microarrays (Clontech Laboratories, Palo Alto, CA, USA) to investigate gene expression patterns in an icv A β mouse model for AD in the cerebral cortex. This approach provided an additional insight into gene processes, which occur not only in AD animal model but also in other neurodegenerative disorders. The knowledge of specific cascades of events leading and causing neurodegeneration will be the key factor in developing and using neuroprotective drugs.

Materials and methods

Reagents A β ₂₅₋₃₅ (Sigma Chemical, St Louis, MO, USA)

was dissolved in sterile normal saline at a concentration of 1 g/L and aggregated by incubation at 37 °C for 72 h. Sodium Dodecyl Sulfate (SDS) was also the product of Sigma. Trizol Reagent was purchased from Gibco-BRL (New York, NY, USA). [α - 32 P]-dATP was from Amersham Pharmacia Biotech (Piscataway, NJ, USA). MMLV reverse transcriptase was from Promega (Madison, WI, USA). Polyclonal integrin α M, c-Jun, and phosphor-c-Jun antibody were purchased from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Atlas Mouse 1.2 Expression Arrays and Atlas cDNA Expression Array Kit were provided by Clontech Laboratories. On each array membrane there are 1176 gene cDNA fragments with lengths of 200–600 bp. The name, coordinate, Gene-Bank Number, and other related information for these genes are available on Clontech's homepage (www.clontech.com).

Animals Male Balb/c mice weighing 22–24 g (Institute of Experimental Animal of Chinese Academy of Medical Sciences, Certificate No SCXK11-00-0006) were maintained in a climate-controlled room at 22±1 °C on a 12 h light/dark cycle with free access to food and water. They were acclimatized to laboratory conditions for 5 d before experiments.

Aggregated A β ₂₅₋₃₅ was icv injected into mice according to the method described by Maurice *et al*^[3]. In brief, a 28-gauge stainless-steel needle was inserted unilaterally 1 mm to the right of the midline point, equidistant from each eye, at an equal distance between the eyes and the ears and perpendicular to the plane of the skull. A β ₂₅₋₃₅ or sterile normal saline at a volume of 5 μ L was delivered gradually within 15 s.

Water maze task The Morris water maze test began on d 11 after A β -injection as described^[7]. The experimental apparatus consisted of a circular water tank (diameter=80 cm) filled to a depth of 15 cm with water maintained at 23±1 °C. The water was made opaque by adding black ink. A platform (diameter=8 cm) was submerged 0.5 cm below the water surface and placed at the midpoint of one quadrant. Two training trials each day were conducted for 4 consecutive days. On each trial, the mice were placed in the pool at one starting position. They were allowed to swim freely or until they found the platform. The time required to escape onto the hidden platform was recorded. Mice that found the platform were allowed to remain on the platform for 10 s and were then returned to the home cage. If a mouse did not reach the platform within 120 s, it was gently guided to platform by the experimenter, where it remained for 10 s. Statistic comparison of the water maze test between the A β -treated and control group was analyzed by two-way analysis of variance (ANOVA).

RNA preparation Mice were killed after the water maze test. Total RNA was extracted from the cerebral cortex with

Trizol according to the manufacturer's instructions. RNA extracts from 10 mice of each group were pooled to provide sufficient material for both cDNA microarray and reverse transcription-PCR analysis. Total RNA was treated with RNase-free DNase at 37 °C for 20 min to avoid contamination of genomic DNA. The RNA quality and concentration were assessed using agarose gel electrophoresis and spectrophotometric reading.

Probe preparation and hybridization Total RNA (5 μ g) was reverse transcribed using reagents provided in the Atlas cDNA Expression Array Kit and radiolabeled with [α - 32 P]-dATP. The array membranes were prehybridized for 1 h at 68 °C in ExpressHyb solution containing 100 mg/L of heat-denatured salmon testis DNA. The denatured 32 P-labeled cDNA was added to Express Hyb hybridization solution at a final concentration of 4.0 MBq/L and array membranes were hybridized with the labeled cDNA overnight at 68 °C. The next day, the membranes were washed three times for 30 min with pre-warmed (68 °C) washing solution 1 [2×SSC (1×SSC is 15 mmol/L sodium chloride, 15 mmol/L sodium citrate), 1% SDS] and once for 30 min with pre-warmed (68 °C) washing solution 2 (0.1×SSC, 0.5% SDS) with continuous agitation at 68 °C. After a final wash with 2×SSC at room temperature for 10 min, the membrane was immediately wrapped in plastic film and exposed to X-ray film at -70 °C.

Detection and analysis Gene expression levels were assessed according to autoradiographic intensity of hybridization signals by software. Only those autoradiograms with similar background and identical signal intensity of housekeeping genes were comparable. Genes that showed an increase or decrease of 2.0-fold or greater were considered as differentially expressed genes. Each group sample was carried out three times, those genes that were all differentially expressed in the three hybridizations were reported as mean±SEM.

Semi-quantitative RT-PCR Expressions of 6 genes were randomly selected to be further confirmed by RT-PCR. Primers were designed using DNA Star software (Perkin Elmer, Norwalk, Conn, USA) and synthesized by Sbs Bio Inc (Beijing, China). Reverse transcription reaction was catalysed by MMLV. PCR was performed for 30 s at 94 °C, 30 s at 55 °C, and 1 min at 70 °C for a cycle, 32 cycles were completed (25 cycles for *GAPDH* gene), and finally 10 min at 72 °C. The RT-PCR products were subjected to electrophoresis in 1% agarose gels and quantitative analysis was carried out using the YLN 2000 Gel Analysis System (Yalien, Beijing, China). The gene expression levels were assessed according to the gray ratio value of the target genes and housekeeping gene, *GAPDH*. The PCR primers used were as follows: *GAPDH*,

forward primer, 5'-CAG CAA CTC CCA CTC TTGCC CCT CCT GTT ATT AT-3'; *AP-1*, forward primer, 5'-AGGCAGAGAGGAAGCGCAT-3', reverse primer, 5'-TGGCACCCACTGTTAACGTG-3'; *NF- κ B*: forward primer, 5'-GCAGCCTATCACCAACTCT-3', reverse primer, 5'-TACTCCTTCTTCTCCACCA-3'; *integrin α M*, forward primer, 5'-TTAATGACTCTGCGTTTGCCC-3', reverse primer, 5'-TCATGTCTTGTACTGCCGCT-3'; *ATPase*, forward primer, 5'-CCATCTTCTGCACCATCGTTC-3', reverse primer, 5'-AGACGCCCTGCCTCTTTCAG-3'; *nerve growth factor α (NGF α)*, forward primer, 5'-ACTGGTGTCTCACAGCTGCC-3', reverse primer, 5'-CGCAGCAGCATCAGGTCAT-3'; *Akt*, forward primer, 5'-CCTGTCTCGAGAGCGTGTGTT-3', reverse primer, 5'-CATAGTGGCACCGTCCTTGA-3'.

Western blot analysis After decapitation, the cerebral cortex from 5 mice were immediately homogenized in 5 volumes of homogenization buffer [Tris-HCl 50 mmol/L (pH 7.5), NaCl 150 mmol/L, 1 % Triton-X 100, 1 % sodium deoxycholate, 0.1 % SDS]. After sonication and centrifugation at 4 °C (10 min, 10 000 \times g), the supernatant was used for determination of protein concentrations, and the equal amounts of total solubilized proteins were eluted by heating with SDS-PAGE sample buffer and separated by SDS-polyacrylamide gel electrophoresis. Proteins were then transferred to the nitrocellulose membrane. The membrane was blocked at room temperature with 5 % non-fat dry milk in Tris-buffered saline with 0.1 % (v/v) Tween-20 (TBST) and incubated with polyclonal integrin α M, c-Jun and phosphor-c-Jun antibody (1:1000) overnight at 4 °C. After washing, the membrane was incubated with horseradish-labeled secondary antibody for 1 h. After five additional washes in TBST, the membrane was incubated with an enhanced chemiluminescence detecting reagent according to the manufacturer's protocols and exposed to X-ray film.

Results

Effects of icv injection of $A\beta_{25-35}$ on performance of the Morris water maze task Effect of $A\beta$ -injection on learning and memory revealed a significant difference between the two groups. $A\beta$ -injected mice exhibited an obvious learning deficit compared to the saline-injected mice. ANOVA analysis showed a significant decrease in the latency time for the control group during the 8 training trials, but not for the experimental group (Figure 1).

Gene expression analysis of mice cerebral cortex after icv injection of $A\beta_{25-35}$ The human cerebral cortex serves to control functions such as speech, memory, logical and emotional response, as well as consciousness, interpretation of




Figure 1. $A\beta_{25-35}$ induced impairment on learning and memory in mice in the Morris water maze task. Ten days after icv administration of the sterile normal saline (Ctr) or the aggregated $A\beta_{25-35}$, mice were subjected to the Morris water maze test. The training trials were carried out on d 11–14. $n=10$. Mean \pm SEM. ^b $P<0.05$, ^c $P<0.01$ vs Ctr group.

sensation, coordination of voluntary movement, and these are all greatly involved and damaged in AD. In our study, we used cDNA microarray to investigate the molecular events of neuronal toxicity induced by $A\beta_{25-35}$ in the cerebral cortex of mice. We compared the gene expression levels in the cerebral cortex of $A\beta$ -injected mice and saline-injected mice. Of 1176 known mouse genes represented on the array, we detected 31 genes differentially expressed in the mice cerebral cortex. Among them 19 genes showed up-regulation in expression after injection of $A\beta$, including *TBX1*, *NF- κ B*, *AP-1/c-Jun*, *cadherin*, *integrin*, *erb-B2*, and *FGFR1*; while expression of 12 genes were down-regulated, including *NGF*, *glucose phosphate isomerase 1*, *AT motif binding factor 1*, *Na⁺/K⁺-ATPase*, and *Akt*. All differentially expressed genes were functionally characterized as a diverse spectrum of the biologic process, including transcription factor, cell cycle protein, cytokines, signal transduction, energy metabolism, neurotrophic factor, and so on (Table 1).

Semi-quantitative RT-PCR To confirm the results of cDNA array, RT-PCR analysis was performed to examine the expression level of 6 genes that were up-regulated or down-regulated in the cerebral cortex of mice. The RT-PCR results indicated that 6 genes showed identical results to that of the microarray (Figure 2).

Western blot analysis The expression of integrin α M and activated AP-1/c-Jun of the cerebral cortex were investigated by Western blot analysis. Although total c-Jun was unchanged in intensity between the two groups, the increased expression of integrin α M and phosphorylated c-Jun were found in the cerebral cortex of $A\beta$ -treated mice (Figure 3).

Table 1. Differentially expressed genes in the cerebral cortex of Aβ-treated mice. *n*=3. Mean±SEM.

| Gene code | Protein/gene | Gene classification | Ratio |
|-----------|--|--|------------|
| U57327 | T-box 1 (TBX 1) | Basic transcription factors | 11.73±1.84 |
| AB009453 | Transcription factor 21 | Basic transcription factors | 5.80±0.56 |
| M61909 | NF-κB p65 | Transcription activators and repressors | 3.17±0.86 |
| J04115 | Transcription factor AP-1; c-Jun | Transcription activators and repressors | 2.78±0.34 |
| X83974 | Transcription termination factor 1 | Oncogenes and tumor suppressors | 2.35±0.29 |
| M83749 | Cyclin D2753 | Cyclins | 2.58±0.11 |
| X07640 | Integrin alpha M | Cell-cell adhesion receptors | 6.27±0.99 |
| M31131 | Cadherin 2; neural-cadherin (NCAD) | Cell-cell adhesion receptors | 3.14±0.72 |
| M60778 | Integrin alpha L | Intracellular transducers /modulators | 2.74±0.66 |
| L47239 | erb-B2 proto-oncogene;neuro/glioblastoma derived oncogene homolog | Growth factor & chemokine receptors | 4.37±1.01 |
| M28998 | Fibroblast growth factor receptor 1 | Growth factor & chemokine receptors | 2.04±0.21 |
| AB010833 | Patched homolog 2 (PTC2; PTC2) | Intracellular transducers/modulators | 5.00±0.38 |
| U43320 | Frizzled homolog 7 (FZD7; FZ7) | Intracellular transducers/modulators | 3.62±1.40 |
| AF080215 | Coagulation factor II receptor-like 3 | Intracellular transducers/modulators | 3.24±0.55 |
| U43205 | Frizzled homolog 3 (FZD3; FZ3) | Intracellular transducers/modulators | 2.94±0.18 |
| X54924 | Neurofibromatosis 1 | Intracellular transducers/modulators | 2.02±0.23 |
| J05154 | Lecithin cholesterol acyltransferase | Extracellular transport/carrier proteins | 2.19±0.72 |
| X15830 | Secretory granule neuroendocrine protein 1, 7B2 protein | Neuropeptides | 6.29±1.66 |
| X02165 | Neurofilament, light polypeptide | Cytoskeleton/motility proteins | 2.58±0.48 |
| L12703 | Engrailed homeobox protein 1 | Transcription activators and repressors | 0.20±0.06 |
| AF005772 | Paired-like homeodomain transcription factor 3 | Basic transcription factors | 0.37±0.14 |
| U43788 | POU domain class 2-associating factor 1; Oct-1 | Transcription activators and repressors | 0.44±0.08 |
| D26046 | AT motif binding factor 1 | Telomere-associated proteins | 0.49±0.17 |
| AF053471 | Sodium/potassium-transporting ATPase isoform 2β polypeptide 2 (Na ⁺ /K ⁺ -ATPase 2β2) | Membrane channels and transporters | 0.23±0.05 |
| M11434 | Nerve growth factor α (NGF α) | Growth factors, cytokines | 0.35±0.11 |
| M14220 | Glucose phosphate isomerase 1 | Growth factors, cytokines | 0.41±0.19 |
| U15159 | LIM-domain containing protein kinase 1 | Intracellular kinase network members | 0.29±0.06 |
| M33960 | Serine (or cysteine) proteinase inhibitor, clade E, member 1 | Inhibitors of proteases | 0.26±0.10 |
| D89076 | Transthyretin | Extracellular transport/carrier proteins | 0.45±0.16 |
| M94335 | Akt proto-oncogene; protein kinase B (PKB) | Apoptosis associated proteins | 0.38±0.04 |
| M29855 | Interleukin 3 receptor | Interleukin and interferon receptors | 0.32±0.14 |




Figure 2. RT-PCR analysis of 6 gene expressions of the cerebral cortex of control (Ctr) or Aβ-treated mice (Aβ). (A) Agarose electrophoresis of PCR products; (B) Comparison of differential expression of 6 genes obtained from microarray and RT-PCR. The levels of gene expression were standardized against that of GAPDH as an internal control. The positive or negative ratios indicated the increase or decrease of gene expression in Aβ-treated mice compared with the control.




Figure 3. The expression of integrin α M and activated c-Jun in the cerebral cortex of control (Ctr) or A β -injected mice (A β). The experiment shown is a representative of three independent experiments showing similar results.

Discussion

Ample experimentation indicates that icv injection of A β to rodents can induce learning and memory impairments as well as neurodegeneration in brain areas related to cognitive function^[3-5]. This model of A β exposure is a useful model *in vivo* for AD. In the present study, we used the Morris water maze test to detect the effect of aggregated A β_{25-35} injection on the learning and memory of mice. Our results corresponded to previous reports that A β_{25-35} could significantly impair the spacial learning ability and memory of mice.

Microarray provides a promising approach to explore novel molecular and cellular mechanisms that may contribute to some diseases. Gene expression changes in the mice cerebral cortex between icv injection of A β_{25-35} and sterile normal saline were analyzed by using cDNA microarray. In total, 31 genes were differentially expressed, of which 19 genes were increased and 12 were decreased. This implies that only a small fraction of the surveyed genes were affected by A β_{25-35} , while most stayed at a stable expression level. The up-regulation or down-regulation of these genes may be responsible for A β -induced memory impairment in mice.

In the present study, we found that several genes encoding transcription and chromatin-modifying factors were differentially expressed after icv administration of A β_{25-35} . For example, *TBX1*, *transcription factor 21*, *transcription factor AP-1/c-Jun*, and *NF- κ B* showed an increased expression level, while *engrailed homeobox protein 1* and *Oct-1* were decreased in expression. *AP-1* complex is composed of a heterodimer *Fos* and *Jun*, the latter is involved in p38/c-Jun N-terminal MAPK pathway (JNK) and considered as a major regulator of both neuronal death and regeneration^[8,9]. A number of reports indicate that *NF- κ B* regulates the expression of various genes that play critical roles in apoptosis, viral replication, tumorigenesis, various autoimmune

diseases, and inflammation^[10,11]. Studies of postmortem brain tissue from patients with AD have revealed increased NF- κ B activity in cells involved in the neurodegenerative process. This is consistent with increased p65 immunoreactivity in neurons and astrocytes in the immediate vicinity of amyloid plaques in brain sections from AD patients^[12]. It is reported that A β can activate *NF- κ B* in the cultured neurons^[13], suggesting a molecular mechanism by which A β may act during AD pathogenesis. We also noticed that the most dramatic change in gene expression was for the transcript encoding for *TBX1*, a member of the T-box binding gene family^[14]. There are at least 21 members of this gene family. The T-box region, which mediates DNA binding, is highly conserved from *Drosophila* to humans. Human and mice *TBX1* proteins share 98% amino acid identity overall and are identical except for 2 residues within the T-box domain. Expression of human *TBX1* in adult and fetal tissues, as determined by Northern blot analysis, were similar to that found in the mice^[15]. These transcription factors are involved in the regulation of developmental processes, where they seem to be required for tissue-specific development. In particular, the *TBX1* gene is a major genetic determinant in Del22q11.2 syndrome in human beings, which is characterized by a 3-Mb deletion on chromosome 22q11.2, cardiac abnormalities, T-cell deficits, cleft palate facial anomalies, and hypocalcemia^[16]. However, there is little known about specific target genes for *TBX1* proteins and the relationship between this gene and AD. The function of the most up-regulated gene, *TBX1*, in the A β -injection mice should be further studied.

Integrin α M and *integrin α L*, two cell adhesion receptors with an A β binding site and involved in the uptake of A β in neurons and glial cells^[17], were observed to be up-regulated in the A β -treated mice. Adhesion proteins usually transduce the extracellular signal into cells and change the cell cytoskeleton. It is very possible that A β binds to integrin proteins and activates the focal adhesion (FA) proteins, paxillin and focal adhesion kinase (FAK). In this regard, fibrillar A β could promote dystrophy by aberrantly activating FA signal transduction cascades that would be involved in A β -induced neuronal dystrophy or cell death^[18,19]. Moreover, the binding of A β to integrin proteins would induce sustained activation of MAPK pathways followed by neurite degeneration and hyperphosphorylation of tau proteins^[20]. N-cadherin, another cell-adhesion protein, also showed an increase in expression after A β -injection. It is a cell surface glycoprotein that mediates calcium-dependent adhesion between neural cells and affects a wide range of cellular processes, including cell adhesion, cell morphogenesis and cell migration.

Two protein tyrosine kinases, *erb-B2*, and *FGF receptor 1*, were increased in expression after A β -injection. Erb-B2 is the receptor of neuregulin, and overexpression has been demonstrated to activate mitogenic signal transduction and enhance proliferative, prosurvival, and metastatic signals in cell lines. The activity of the MAPK pathway is up-regulated in cells overexpressing *erb-B2*^[21,22]. However, the understanding of the function of *erb-B2* in neural cells is very limited. This basic growth factor has pleiotropic effects and plays a regulatory role in angiogenesis, smooth muscle cell growth and wound healing through fibroblast growth factor receptors (FGFR)^[23]. FGFR1 activation is followed by intracellular signaling cascades involving Ca²⁺ channels, phospholipase C, src family kinases, mitogen-activated kinases and the Grb2/Sos signaling complex^[24,25]. An increase of FGFR2 in brain neurons and retinal pigment epithelial cells can lead to the activation of L-type channels of neuroendocrine and induce gene expression through the phosphorylation of CREB (cyclic AMP-responsive element binding protein)^[26]. Whether the up-regulation of neuroendocrine protein 1 resulted from the increase of FGFR 1 in our study still needs to be studied further.

It is well known that A β can impair mitochondria energy metabolism and selectively inhibit Na⁺/K⁺-ATPase activity both *in vivo* and *in vitro*. From our present results, we found that two Na⁺/K⁺-ATPase related genes, *AT motif-binding factor 1 (ATBF1)*, and *Na⁺/K⁺-ATPase 2 β 2*, were down-regulated. This is consistent with previous studies. ATBF1 is a large transcription factor protein containing four homeodomains and 23 zinc finger motifs. It also includes several motifs that were found in ATPases^[27]. The ATPase activity has been reported to be associated with ATBF1 activity in a unique DNA/RNA-dependent manner that requires both homeodomain and zinc finger motifs^[28].

Interleukin 3 receptor has been reported to have a neurotrophic effect on central cholinergic neurons^[29]. Down-regulation of this gene may be responsible for cholinergic dysfunction induced by A β . Another two growth factors, *NGF* and *glucose phosphate isomerase 1*, were also down-regulated. A large number of reports have shown that these neurotrophic factors have beneficial effects on neural cell survival both *in vivo* and *in vitro*.

The expression of an apoptosis associated protein, *Akt*, decreased in the A β -injected mice. This protein can protect cells from a variety of death-promoting insults in several different cell types, including A β toxicity^[30]. Activated Akt would lead to phosphorylation of the proapoptotic protein Bad, stimulate Bcl-2 and Bax expression, decrease the interaction of Bad with Bcl-xL, increase the Bcl-xL/Bad ratio, even-

tually promoting cell survival^[31].

In summary, we observed extensive activation of transcription factors and the signal transduction system, which may result in neuron apoptosis, energy metabolism and calcium ion homeostasis dysfunction, cell adhesion and neuronal dystrophy. A number of the candidate genes we describe have not been linked to AD in previous publications, and define novel injury pathways warranting further studies. The genes found from the cDNA microarray provide some clues for pursuing a more complete understanding of the molecular mechanism of A β -induced neural degeneration and memory impairment.

References

- 1 Price DL, Sisodia SS. Mutant genes in familial Alzheimer's disease and transgenic models. *Annu Rev Neurosci* 1998; 21: 479–505.
- 2 Cummings BJ, Pike CJ, Shankle R, Cotman CW. β -Amyloid deposition and other measures of neuropathology predict cognitive status in Alzheimer's disease. *Neurobiol Aging* 1996; 17: 921–33.
- 3 Maurice T, Lockhart BP, Privat A. Amnesia induced in mice by centrally administered β -amyloid peptides involves cholinergic dysfunction. *Brain Res* 1996; 706: 181–93.
- 4 Yamada K, Tanaka T, Han D, Senzaki K, Kameyama T, Nabeshima T. Protective effects of idebenone and α -tocopherol on β -amyloid-(1-42)-induced learning and memory deficits in rats: implication of oxidative stress in β -amyloid-induced neurotoxicity *in vivo*. *Eur J Neurosci* 1999; 11: 83–90.
- 5 Flood JF, Morley JE, Roberts E. Amnesic effects in mice of four synthetic peptides homologous to amyloid β protein from patients with Alzheimer disease. *Proc Natl Acad Sci USA* 1991; 88: 3363–6.
- 6 Lockhart DJ, Winzler EA. Genomics, gene expression and DNA arrays. *Nature* 2000; 405: 827–36
- 7 Morris R. Developments of a water-maze procedure for studying spatial learning in the rat. *J Neurosci Methods* 1984; 11: 47–60.
- 8 Behrens A, Sibilina M, Wagner EF. Amino-terminal phosphorylation of c-Jun regulates stress-induced apoptosis and cellular proliferation. *Nat Genet* 1999; 21: 326–9.
- 9 Crocker SJ, LambaWR, Smith PD, Callaghan SM, Slack RS, Anisman H, *et al*. c-Jun mediates axotomy-induced dopamine neuron death *in vivo*. *Proc Natl Acad Sci USA* 2001; 98: 13385–90.
- 10 Baeuerle PA, Baichwal VR. NF- κ B as a frequent target for immunosuppressive and anti-inflammatory molecules. *Adv Immunol* 1997; 65: 111–37.
- 11 Baichwal VR, Baeuerle PA. Activate NF- κ B or die? *Curr Biol* 1997; 7: R94–6.
- 12 O'Neill LA, Kaltschmidt C. NF- κ B: a crucial transcription factor for glial and neuronal cell function. *Trends Neurosci* 1997; 20: 252–8.
- 13 Barger SW, Horster D, Furukawa K, Goodman Y, Krieglstein J, Mattson MP. Tumor necrosis factors α and β protect neurons against amyloid β -peptide toxicity: evidence for involvement of

- a κ B-binding factor and attenuation of peroxide and Ca^{2+} accumulation. *Proc Natl Acad Sci USA* 1995; 92: 9328–32.
- 14 Papaioannou VE, Silver LM. The T-box gene family. *Bioessays* 1998; 20: 9–19.
 - 15 Chieffo C, Garvey N, Gong W, Roe B, Zhang G, Silver L, *et al*. Isolation and characterization of a gene from the DiGeorge chromosomal region homologous to the mouse Tbx1 gene. *Genomics* 1997; 43: 267–77.
 - 16 Yagi H, Furutani Y, Hamada H, Sasaki T, Asakawa S, Minoshima S, *et al*. Role of TBX1 in human del22q11.2 syndrome. *Lancet* 2003; 362: 1366–73.
 - 17 Bi X, Gall CM, Zhou J, Lynch G. Uptake and pathogenic effects of amyloid beta peptide 1-42 are enhanced by integrin antagonists and blocked by NMDA receptor antagonists. *Neuroscience* 2002; 112: 827–40.
 - 18 Berg MM, Krafft GA, Klein WL. Rapid impact of β -amyloid on paxillin in a neural cell line. *J Neurosci Res* 1997; 50: 979–89.
 - 19 Williamson R, Scales T, Clark BR, Gibb G, Reynolds CH, Kellie S, *et al*. Rapid tyrosine phosphorylation of neuronal proteins including tau and focal adhesion kinase in response to amyloid- β peptide exposure: involvement of Src family protein kinases. *J Neurosci* 2002; 22: 10–20.
 - 20 Anderson KL, Ferreira A. α 1 integrin activation: a link between β -amyloid deposition and neuronal death in aging hippocampal neurons. *J Neurosci Res* 2004; 75: 688–97.
 - 21 Wen D, Peles E, Cupples R, Suggs SV, Bacus SS, Luo Y, *et al*. Neu differentiation factor: a transmembrane glycoprotein containing an EGF domain and an immunoglobulin homology unit. *Cell* 1992; 69: 559–72.
 - 22 Tari AM, Lopez-Berestein G. Serum predominantly activates MAPK and Akt kinases in EGFR- and erbB2-over-expressing cells, respectively. *Int J Cancer* 2000; 86: 295–7.
 - 23 Bikfalvi A, Klein S, Pintucci G, Rifkin DB. Biological roles of fibroblast growth factor-2. *Endocr Rev* 1997; 18: 26–45.
 - 24 Koike H, Saito H, Matsuki N. Effect of fibroblast growth factors on calcium currents in acutely isolated neuronal cells from rat ventromedial hypothalamus. *Neurosci Lett* 1993; 150: 57–60.
 - 25 Williams EJ, Furness J, Walsh FS, Doherty P. Characterization of the second messenger pathway underlying neurite outgrowth stimulated by FGF. *Development* 1994; 120: 1685–93.
 - 26 Rosenthal R, Thieme H, Strauss O. Fibroblast growth factor receptor 2 (FGFR2) in brain neurons and retinal pigment epithelial cells act via stimulation of neuroendocrine L-type channels (Ca_v1.3). *FASEB J* 2001; 15: 970–7.
 - 27 Miura Y, Tam T, Ido A, Morinaga T, Miki T, Hashimoto T, *et al*. Cloning and characterization of an ATBF1 isoform that expresses in a neuronal differentiation-dependent manner. *J Biol Chem* 1995; 270: 26840–8.
 - 28 Kawaguchi M, Miura Y, Ido A, Morinaga T, Sakata N, Oya T, *et al*. DNA/RNA-dependent ATPase activity is associated with ATBF1, a multiple homeodomain-zinc finger protein. *Biochim Biophys Acta* 2001; 1550: 164–74.
 - 29 Tabira T, Chui DH, Fan JP, Shirabe T, Konishi Y. Interleukin-3 and interleukin-3 receptors in the brain. *Ann NY Acad Sci* 1998; 840: 107–16.
 - 30 Martin D, Salinas M, Lopez-Valdaliso R, Serrano E, Recuero M, Cuadrado A. Effect of the Alzheimer amyloid fragment A β (25–35) on Akt/PKB kinase and survival of PC12 cells. *J Neurochem* 2001; 78: 1000–8.
 - 31 Seo JH, Rah JC, Choi SH, Shin JK, Min K, Kim HS, *et al*. α -Synuclein regulates neuronal survival via Bcl-2 family expression and PI3/Akt kinase pathway. *FASEB J* 2002; 16: 1826–8.