

Full-length article

Identification and classification of all potential hemolysin encoding genes and their products from *Leptospira interrogans* serogroup Icterohaemorrhagiae serovar Lai¹

Yi-xuan ZHANG^{2,3,4,5}, Yan GENG^{2,3,4,5}, Bo BI^{2,4}, Jian-yong HE⁴, Chun-fu WU⁴, Xiao-kui GUO³, Guo-ping ZHAO^{2,4,6,7}

²Research Center of Biotechnology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200233, China;

³Department of Microbiology and Parasitology, Shanghai Second Medical University, Shanghai 200025, China; ⁴The School of Pharmaceutical Engineering, Shenyang Pharmaceutical University, Shenyang 110016, China; ⁵Chinese National Human Genome Center at Shanghai,

ZhangJiang High Tech Park, Shanghai 201203, China

Key words

hemolysin; *Leptospira interrogans*; sphingomyelinase

¹ Project supported by the National Natural Science Foundation of China (No 30370071) and the National High Technology Research and Development Program of China (863 Program, No 2003, AA223030).

⁵ The two authors contributed equally to this work.

⁷ Correspondence to Prof Guo-ping ZHAO. Phn 86-21-5080-1919. Fax 86-21-5080-1922. E-mail gpzhao@sibs.ac.cn

Received 2004-11-18

Accepted 2005-01-06

doi: 10.1111/j.1745-7254.2005.00075.x

Abstract

Aim: To identify and classify all potential hemolysin candidates of *Leptospira interrogans* serogroup Icterohaemorrhagiae serovar Lai. **Methods:** All of the potential hemolysin encoding genes were characterized *in silico*. These genes were cloned and expressed in *Escherichia coli*. The hemolytic activities of the expressed proteins were assayed observing the hemolysis on sheep blood agar plates. Sphingomyelinase activities of the hemolysin candidates were measured by thin-layer chromatography (TLC) and HPLC for sphingomyelin-hydrolysis. Expression and secretion of the hemolysins in *L. interrogans* were studied by reverse transcription polymerase chain reaction, Western blot, and enzyme-linked immunosorbent assays. **Results and Conclusion:** The hemolytic activities of hemolysin candidates (LA0327, LA0378, LA1027, LA1029, LA1650, LA3050, LA3937, LA4004) from *L. interrogans* strain Lai were confirmed. They were further divided into two groups, sphingomyelinase hemolysins and non-sphingomyelinase hemolysins, based on their ability to hydrolyze sphingomyelin. Most of these hemolysins were actually expressed in living *L. interrogans* and some of them were secreted into the environment. This study establishes an essential and complete basis for further studying the contribution of hemolysins to the pathogenesis of *L. interrogans*.

Introduction

Leptospirosis is a worldwide zoonosis caused by pathogenic species of *Leptospira*, particularly, *Leptospira interrogans* serovar Lai^[1]. Pathogenic *Leptospira* infection causes leptospirosis (Weil's syndrome), which manifests with jaundice and renal failure^[2–4] along with prominent respiratory symptoms. In some cases, severe pulmonary hemorrhages may happen, which may lead to sudden death. Although potential virulence factors, such as hemolysin, lipopolysaccharide (LPS), and heat shock proteins, are suggested for leptospiral infection, the pathogenetic mechanism of leptospirosis is yet to be clarified.

Among these suggested virulence factors, bacterial hemolysin has been demonstrated in several pathogenic bacteria^[5,6] and a limited number of them have been identified in *L. interrogans*^[7]. Annotation of the complete genomic sequence of *L. interrogans* serogroup Icterohaemorrhagiae serovar Lai^[8] indicated that there were ten putative hemolysin genes located on the large chromosome (CI, GB: AE010300), of which, one (LA3540) had been previously identified^[7]. Our current work characterized all the other genes, except LA0177 because of its extremely short nucleotide sequence. The classification of the hemolytic activities and their expression and secretion in *L. interrogans* were investigated.

Materials and methods

Bacterial strains and plasmids The virulent *L. interrogans* serogroup Icterohaemorrhagiae serovar Lai type strain 56601 used in this study was maintained by the Institute for Infectious Disease Control and Prevention, Beijing, China. The avirulent strain of *L. interrogans* serogroup Icterohaemorrhagiae serovar Lai (strain IPAV) was given as a gift by Dr PICARDEAU, M (Institute Pasteur, Paris, France). Strains were grown in liquid Ellinghausen-McCullough-Johnson-Harris (EMJH) medium^[1] at 28 °C under aerobic conditions and collected at a density of approximately 1×10^8 bacteria per mL. *Escherichia coli* DH5 α and BL21 (DH3) were used for the cloning and expression of hemolysin candidates, respectively. The pUCm-T and pET28b plasmids were served as the vectors for cloning and expression, respectively.

Characterization of hemolysin candidates First, coding sequences (CDSs) potentially encoding hemolysin candidates were identified during the genomic annotation^[8]. Second, amino acid sequences of those hemolysin candidates were analyzed with the SWISS-PROT/TrEMBL non-redundant databases^[9] to obtain homologous proteins, which were further compared with BioEdit (Tom Hall, North Carolina State University, Carolina, USA). Third, the domain structures of the hemolysin candidates were predicted by Pfam^[10], PROSITE^[11] or ProDom^[12]. The secondary structures of the proteins were predicted by Jpred^[13]. Finally, multiple sequence alignments were made for sphingomyelinase-like hemolysin homologous proteins using clustW, and Mega² (Sudhir Kumar, Arizona State University, Arizona, USA) was used to establish their phylogenetic tree.

Cloning, expression, and purification of the recombinant hemolysin candidates in *E. coli* Genomic DNA was isolated from *L. interrogans* strain 56601 cultivated in EMJH medium. The hemolysin candidate genes were obtained by polymerase chain reaction (PCR), ligated with pUCm-T vector and transformed into DH5 α cells. The confirmed recombinant plasmids were digested with corresponding enzymes, ligated with pET28b vector and then transformed into BL21 (DH3) cells. *E. coli* cells were grown in Luria-Bertani medium supplemented with kanamycin at 50 mg/L at 37 °C. Protein expression was induced at A_{600} of 0.6 by addition of isopropyl-beta-D-thiogalactopyranoside (IPTG) at 0.6 mmol/L for 3 h. The harvested cells were suspended in Tris-HCl buffer (20 mmol/L Tris-HCl, pH 7.9) and lysed by sonication. The insoluble inclusion bodies were dissolved in the buffer (Tris-HCl 20 mmol/L, pH 7.9, NaCl 0.5 mol/L, 10% glycerol, guanidium HCl 6 mol/L) and then centrifuged at 20 000 \times g for

20 min. The soluble supernatant was applied to Ni-NTA His-Bind resin and the His-tag fusion protein was eluted by an imidazole gradient from 10 mmol/L to 1000 mmol/L in the elution buffer (20 mmol/L Tris-HCl pH 7.9, 0.5 mol/L NaCl, 10% glycerin). The purified proteins were analyzed by sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) with coomassie brilliant stain.

Hemolytic activity assay on sheep blood agar plates Sheep blood plates, 10% (v/v) supplemented with 25 mmol/L MgCl₂ and 100 mg/L kanamycin, were used to measure hemolytic activities. Lysate of *E. coli* cells expressing hemolysin candidate proteins (100g/L) was dropped onto the plates, and the plates were incubated at 37 °C for 16 h, and then placed at 4 °C for 30 min. Sphingomyelinase C (0.1 U, Sigma, St Louis, MO, USA) and lysate of *E. coli* cells harboring pET28b (100 g/L) were used as positive and negative controls, respectively.

Sphingomyelinase assay by thin-layer chromatography and HPLC The biphasic system consisted of an ether: methanol (9:1 v/v) organic phase containing sphingomyelin (2 g/L) and a water phase (25 mmol/L MgCl₂) containing the lysate of *E. coli* cells expressing hemolysin candidate proteins (100 g/L). This biphasic solution was shaken at 37 °C for 4 h. After that, the organic phase (15 μ L) was applied on a silica gel-60 coated glass plate. The chromatogram was developed with a mobile phase (chloroform:methanol:water: 25% ammonia 58.0:35.0:3.5:3.5 v/v). Lipids were visualized by spraying on a plate with 30% H₂SO₄ at 110 °C for 10 min. For HPLC, sphingomyelinase activity was determined in a biphasic system as described above, except for the organic phase containing sphingomyelin (1 g/L). The organic phase (7.5 μ L) was applied to HPLC (YWG C₁₈ 5 μ m, 200 mm \times 4.6 mm, China) and the elutes were monitored by absorption at 207 nm. The mobile phase was a mixture of acetonitrile: methanol:water (154:45:81 v/v). The flow rate was 1 mL/min. Sphingomyelinase C (0.1 U) and lysate of *E. coli* cells harboring pET28b (100 g/L) were used as positive and negative controls, respectively.

Validation of hemolysin encoding gene expressions in *Leptospira interrogans* strain Lai by reverse transcription RCR and Western blot According to the manufacturer's instructions, total RNA was extracted with TRIzol reagent (Invitrogen, Carlsbad, California, USA) from *L. interrogans* strain Lai cultivated in EMJH medium. Total RNA (1 μ g) from each sample was reverse-transcribed into cDNA according to the instructions provided with the cDNA Synthesis Kit (Invitrogen, Carlsbad, California, USA). Equal amounts of the product of the reverse transcription reaction were subjected to PCR amplification. The primers and related infor-

Table 1. Primers of hemolysin candidate genes in reverse transcription polymerase chain reaction (RT-PCR) assay and the product information. T_m, temperature.

Gene ID	RT-PCR sense primer (5'-3')	RT-PCR antisense primer (5'-3')	Product length (bp)	T _m (°C)
LA1027	CAT ATG TAC CCA TAC CAA ACC AAC	CCG TAT ACC GGA TAA CGA TCT GAAA	665	50
LA1029	TGG GAC ACG ATG AAA GAG C	ATC CAG ATA CGG TCC AAG TTT	750	52
LA4004	GTT TAC GCC GAC CCT TCC ACT CC	CTA ACG ATA AAT TAG ATC CTT GCT CCA	569	52
LA3050	CAT ATG TCT AAA ACG ACA CGA AAA	AAA GTT TCG GTT GGT GGT	501	50
LA0327	CAT ATG CTT CTT TTT GAAAGG GGA T	ATC CGA CTT AAC TCT ATA AGT TCC A	763	54
LA3937	CAT ATG CTT TTG AAG CTG ATT GGA	AAG AAG ACC CGC AAC TCC ACC	495	54
LA1650	AGAGGT GCT GAT TGT TGG G	CTA GAG CAAATT AGA TTT GTC TGG	504	52
LA0378	AAG ATC CTG AAA ATG TAA AA	AAC AAT GCG ATT TGG TTT TGT GGA G	659	50

mation are shown in Table 1. After amplification, 5 µL of each PCR reaction product was electrophoresed on a 1.5% (w/v) agarose gel containing ethidium bromide (0.5 mg/L). For Western blot analysis, *L. interrogans* strains cultivated in EMJH medium were harvested by centrifugation at 14 600×g for 10 min, electrophoresed on 10% SDS-PAGE gels and electrotransferred to nitrocellulose membrane under a constant voltage of 5 mA/cm² for 1 h. The blot was first masked by the blotting buffer (10% skim milk in 10 mmol/L Tris-HCl buffer with NaCl 150 mmol/L and 0.1% Tween-20) for 2 h and then incubated with rabbit anti-hemolysin antibody (1:5000) for 2 h. After being washed with TBST 3 times, the blot was incubated for 1 h with alkaline phosphatase (AP)-conjugated goat anti-rabbit IgG antibody (1:7500) and the color was developed by addition of bromo-chloro-indoryl phosphate/nitroblue tetrazolium (BCIP/NBT) for AP reaction.

Detection of hemolysin secretion in *L. interrogans*
Briefly, the culture supernatant (100 ng/well) of *L. interrogans* in Korthof^{f1} was coated onto 96-well ELISA plates by incubation at 37 °C for 2 h. After the removal of blocking solution and washing 3–5 times with phosphate-buffered saline buffer (PBS) containing 0.1% Tween (PBS-Tween), anti-hemolysin antibodies were incubated in the plate at 37 °C for 2 h. The plate was washed 3–5 times with PBS-Tween and then incubated with AP-conjugated goat anti-rabbit IgG antibody (1:1500) at 37 °C for 1 h. The plate was washed 3–5 times with PBS-Tween and incubated with the BCIP/NBT for 20 min in the dark. Absorbance at 492 nm was recorded using an automated ELISA microplate reader.

Results

The *in silico* structural analysis indicated that there were at least two kinds of hemolysin candidates from *L. interrogans* Genomic annotation of *L. interrogans* strain Lai

indicated that there were 10 genes putatively encoding proteins highly similar to hemolysins reported in the NCBI/Genbank and SwissProt/TrEMBL databases (Table 2).

Domain structure analysis of the candidate proteins indicated that 5 of them had barring of the characteristic domains of the phosphatase family. Specifically, the Pfam-based domain analysis indicated that all of LA1027, LA1029, LA4004, and LA3050 barred a conserved domain PF03372 of the phosphatase family. The ProDom-based analysis showed that all of the proteins, except LA3050, were highly similar to *L. interrogans* serovar hardjo sphingomyelinase-like hemolysin (SP17627/SwissProt), barring phosphatase domains of PD011673, PD447657, and PD041204 in similar regions of the proteins. Although LA3050 was reasonably similar to SP17627 (67%) and also had the PD041204 domain, it lacks the other two domains mainly because of its short primary peptide sequence. Although LA3540 had PD011673, PD041204, and the phosphatase conserved domain PF03372, LA3540 had previously been identified as a pore-forming hemolysin^[5].

Amino acid sequence analysis of these sphingomyelinase-like hemolysin candidates indicated that the conserved Mg²⁺-complexing glutamic acid and asparagine involved in substrate binding were identified in LA1029, Glu²⁰⁰, and Asn³⁴³, corresponding to the Glu¹³¹ and Asn²⁷⁴ of SP17627, respectively. For LA4004, only Asn²⁶⁷ was identified as corresponding to the conserved Asn²⁷⁴ of SP17627. Protein secondary structures were predicted by Jpred², and their similarities are shown in Figure 1.

The remaining 5 candidate proteins have little sequence similarity to those of sphingomyelinase-hemolysin proteins. Based on the *in silico* analysis, they do not have any phosphatase family domains or share any common domains. Therefore, these hemolysin candidates, if they are active, may function based on the mechanism of pore formation and




Figure 1. Primary and secondary structure comparisons of SP17627 and *Leptospira interrogans* sphingomyelinase-like hemolysins. Multiple amino acid sequence alignment was performed using BioEdit. Identical amino acids are shaded in red. Similar amino acids are shaded in yellow. The Mg²⁺-complexing glutamic acid and asparagine involved in substrate binding are indicated by arrows; these were Glu¹³¹ and Asn²⁷⁴ in SP17627. Correspondingly, the conserved sites were found at Glu²⁰⁰ and Asn³⁴³ in LA1029, and Asn²⁶⁷ in LA4004. Protein secondary structures were predicted by Jpred², and only the results of SP17627 (up) and LA1029 (down) are shown (the grey belts are alpha-helices and the black belts are beta-sheets).

Table 2. Hemolysin candidate genes of *Leptospira interrogans*.

Groups	Gene identity	Gene name	Gene length (bp)	Transcription orientation	Product	Molecular weight (Da)	Isoelectric point
Sphingomyelinase	LA1027	<i>sph1</i>	1794	Reverse (R)	Sphingomyelinase C precursor	68 192	8.15
	LA1029	<i>sph2</i>	1872	R	Sphingomyelinase C precursor	71 030	5.95
	LA4004	<i>sph3</i>	1665	Forward (F)	Sphingomyelinase C precursor	64 433	7.94
	LA3050	<i>sph4</i>	720	R	Sphingomyelinase C precursor	27 926	8.73
Non-sphingomyelinase	LA3540	<i>sphH</i>	1677	R	Sphingomyelinase C precursor	65 336	8.84
	LA0327	<i>tlyA</i>	831	F	Hemolysin	31 670	9.54
	LA0378	<i>hlyX</i>	1179	R	TPR-repeat-containing protein	44 956	6.17
	LA1650	<i>hlpA</i>	942	F	Hemolysin hemolytic protein	36 534	8.52
	LA3937	<i>hlyC</i>	1335	R	Similar to hemolysin	50 431	5.04
	LA0177	<i>hlyA</i>	225	F	Alpha-hemolysin	8 590	9.64

other mechanisms.

Phylogenetic analysis was performed on the sphingomyelinase-hemolysin proteins. An neighbor-joining unrooted phylogenetic tree was established including sequences of SP17627, LA1027, LA1029, LA4004, LA3050, and other similar proteins, with a cut-off amino acid identity of 53% and similarity of 67%. Hemolysins from *Staphylococcus aureus*, *Bacillus cereus*, *Listeria ivanovii*, and *Pseudomonas* sp strain TK4 made up one branch of the tree, and the sphingomyelinase-like proteins of *L interrogans* made up another branch (Figure 2).




Figure 2. Comparison of sphingomyelinase-hemolysin protein sequences by neighbor-joining method shown in unrooted tree. STAAU, *Staphylococcus aureus*; BACCE, *Bacillus cereus*; LISIV, *Listeria ivanovii*; PSESP, *Pseudomonas* sp strain TK4; LEPIN, *Leptospira interrogans*.

The hemolytic activities of the recombinant hemolysin candidates from *L interrogans* were determined Among the 10 hemolysin candidates from *L interrogans*, LA3540

had previously been identified^[7], whereas LA0177 is extremely short in its sequence and has no detectable phosphatase domain. Therefore, only the remaining 8 hemolysin candidate genes were cloned into *E coli* and the recombinant proteins were purified to homogeneity determined by




Figure 3. Purification of recombinant hemolysins. Protein markers: 97 400 Da, 66 200 Da, 43 000 Da, 31 000 Da, 20 100 Da, 14 400 Da.

SDS-PAGE followed by coomassie brilliant stain (Figure 3).

Crude cell lysates of *E coli* expressing cloned hemolysin candidates were spotted onto sheep blood agar plates and, after incubation at 37 °C for 16 h, clear hemolytic zones were observed for all the candidates via a cold-warm hemolytic procedure (Figure 4). The hemolytic zone of LA1029 appeared at first and the area of hemolysis was the biggest. Clear hemolytic zones of LA1027 and LA4004 appeared a little later. Hemolysis caused by LA3050 appeared last and the area of the hemolytic zone was the smallest. For those predicted to be in the non-sphingomyelinase-hemolysin family, hemolytic abilities were also different: LA3937 was the highest and LA0327, LA0378, and LA1650 were second




Figure 4. Hemolysis of recombinant hemolysin candidates on sheep blood planes.

highest, third highest, and lowest, respectively.

The sphingomyelinase activities of hemolysin candidates were detected The sphingomyelinase activities of recombinant hemolysin candidates were detected by TLC assay. Results showed that only LA1027, LA1029, LA4004, and LA3050 could hydrolyze sphingomyelin. These results were confirmed by HPLC assay. The peak area of sphingomyelin was diminished after hemolysin treatment and the percentages of hydrolyzed sphingomyelin are displayed in Table 3. The percentages of hydrolyzed sphingomyelin of sphingomyelinase (SphC), LA1029, LA1027, LA4004, and LA3050 were 73.99%, 68.87%, 61.10%, 56.52%, and 48.04%, respectively. LA3937, LA0327, LA0378, and LA1650 could not hydrolyze sphingomyelin under the same condition.

Hemolysin gene expression and hemolysin secretion in *L interrogans* RT-PCR analysis indicated that all the 8 candidates, LA0327, LA0378, LA1027, LA1029, LA1650, LA3050, LA3937, and LA4004, were transcribed in *L interrogans* strain Lai and strain IPAV cultivated in EMJH or Korthof culture medium (Figure 5). The transcription levels of some hemolysin encoding genes, LA0378, LA0327, and LA3050, were higher in *L interrogans* cultivated in Korthof medium than those in the EMJH medium. Western blot can detect hemolysin proteins in strain Lai cell crude extracts for all the candidates except LA1027 (Table 4). In addition, we failed to detect this protein by Western blot in the strain IPAV cultured in EMJH medium. In addition, we failed to detect LA3050 under the same condition, whereas the other 6 hemolysin candidates were all detected. LA1029, LA4004, LA3050, LA1650, and LA3937 were secreted into the environment, as determined by ELISA, and the secretion levels were higher in strain Lai than those in strain IPAV (Figure 6). LA0327 was secreted into the environment both in strain Lai and in strain IPAV. LA1027 was not secreted either in strain Lai or in strain IPAV.

Discussion

In 1956, Alexander *et al*^[14], for the first time, discovered that *L interrogans* had a hemolysin-like substance that could hemolyze red blood cells of ruminants, such as sheep, cows,

Table 3. Percentage of hydrolyzed sphingomyelin after reaction with hemolysins for 4 h.

	SMaseC	LA1029	LA1027	LA4004	LA3050
Peak area before reaction	317 274	317 274	317 274	317 274	317 274
Peak area after reaction	82 498	98 754	123 410	137 946	164 836
Hydrolyzed sphingomyelin (%)	73.99	68.87	61.10	56.52	48.04




Figure 5. Transcription of hemolysin candidate genes encoded by *L interrogans* chromosome by reverse transcription polymerase chain reaction detection. 1, virulent strain Lai cultivated in Korthof (KV); 2, avirulent strain IPAV cultivated in Korthof (KNV); 3, avirulent strain IPAV cultivated in EMJH medium (ENV); 4, virulent strain Lai cultivated in EMJH medium (EV). DNA markers: 400 bp, 600 bp, 800 bp, 1200 bp, 1600 bp, 2000 bp.

Table 4. Hemolysin candidate gene expression in *L interrogans*. ELISA, enzyme-linked immunosorbent assay; NV, avirulent strain IPAV; V, virulent strain Lai.

Gene ID	Western blot assay	
	V	NV
LA1027	-	-
LA1029	+	+
LA4004	+	+
LA3050	+	-
LA0327	+	+
LA0378	+	+
LA1650	+	+
LA3937	+	+

and goats. Its hemolytic activity was completely lost after being heated at 56 °C for 5 min, indicating its heat-labile character. Later, hemolysins were detected in many pathogenic *L interrogans*^[15].

L interrogans gene encoding hemolysin was first cloned from the pathogenic serovar hardjo. This protein obviously possessed both hemolytic and sphingomyelinase C activi-

ties^[16]. The only hemolysin gene cloned from serovar lai before our efforts was CDS LA3540^[7]. Although this gene encoded a hemolytically active protein with a phosphatase-like domain, the purified protein did not have any detectable sphingomyelinase activity. Therefore, it was proposed that it was a transmembrane pore formation protein. The concept of transmembrane pore formation by bacterial protein toxins was first brought forward by Fussle *et al*^[17] to explain the mechanism of action of staphylococcal alpha-toxin and it is apparent today that the majority of medically relevant pathogens produce pore-forming proteins^[18]. Many of these toxins have been designated as hemolysins because of their lytic action on red blood cells.

Our studies have proven experimentally that, besides the previously characterized LA3540 and the very short hemolysin candidate gene, LA0177, the recombinant proteins encoded by the candidate genes from *L interrogans* strain Lai (LA0327, LA0378, LA1027, LA1029, LA1650, LA3050, LA3937, LA4004) were hemolysin. According to the *in silico* structure analysis, LA1027, LA1029, LA4004, and LA3050 are highly similar to SP17627, the hemolytic sphingomyelinase C from serovar hardjo, with respect to domain organizations, amino acid sequences, and the predicted secondary structures. These predictions were further confirmed by sphingomyelinase assays. Other candidates, LA0327, LA0378, LA1650, and LA3937, were confirmed to be non-sphingomyelinase hemolysins. They are likely either to be pore-forming proteins or to have other mechanisms of hemolytic activity.

Spirochetes are evolutionarily primitive and *L interrogans* is a facultative free-living pathogen possessing a large number of different kinds of hemolysins. Therefore, questions about the possibilities of convergence, divergence, or other models of evolution should be addressed in the future. In this study, the phylogenetic analysis of sphingomyelinase from *L interrogans*, both serovar lai and serovar hardjo, with those from other bacteria indicated that the sphingomyelinase hemolysins of *L interrogans* are closely related but distantly different from those of the others. Thus, this group of hemolysins is more likely to be evolved from a common ancestor with a divergence mechanism. For the other group of hemolysins, because of their high level of diversity, the possibility of horizontal gene transfer cannot be excluded.

Almost all of these hemolysins were expressed *in vivo* under normal culture conditions. Among them, LA1029, LA4004, LA1650, and LA3937 were secreted into the environment and the secretion level was significantly higher ($P < 0.01$) in the virulent strain Lai than in the avirulent strain IPAV (Figure 6). LA0327 was secreted into the environment both in strain Lai and IPAV without significant difference.




Figure 6. Secretion of hemolysin candidates in *Leptospira interrogans* by enzyme-linked immunosorbent assay. C, control (blank Korthof medium); NV, avirulent strain IPAV; V, virulent strain Lai. ^b $P < 0.05$; ^c $P < 0.01$ vs control. ^e $P < 0.05$, ^f $P < 0.01$ vs NV.

LA1027 was not expressed or secreted in either strain Lai or IPAV, which was testified by ELISA and Western blot assay (Figure 6, Table 4). The significant difference in hemolysin expression and secretion between the virulent strain Lai and the avirulent strain IPAV may suggest that hemolysins might play an important role in the pathogenesis of *Leptospira*.

Acknowledgment

We thank Shuang-xi REN for suggestions and comments, and Bao-yu HU and Xiu-gao JIANG for help in bacterial cul-

ture preparation.

References

- 1 Faine S. *Leptospira* and leptospirosis. New York: CRC Press; 1994.
- 2 Kmety E, Dikken H. Classification of the species *Leptospira interrogans* and history of its serovars. Groningen: University Press; 1993.
- 3 Ramadass P, Jarvis BD, Corner RJ, Penny D, Marshall RB. Genetic characterization of pathogenic *Leptospira species* by DNA hybridization. *Int J Syst Bacteriol* 1992; 42: 215–9.
- 4 Ozdemir V, Erol E. Leptospirosis in Turkey. *Vet Rec* 2002; 150:

- 248–9.
- 5 Lee SH, Kim S, Park SC, Kim MJ. Cytotoxic activities of *Leptospira interrogans* hemolysin SphH as a pore-forming protein on mammalian cells. *Infect Immun* 2002; 70: 315–22.
 - 6 Sakurai J, Honda T, Jinguji Y, Arita M, Miwatani T. Cytotoxic effect of the thermostable direct hemolysin produced by *Vibrio parahaemolyticus* on FL cells. *Infect Immun* 1976; 13: 876–83.
 - 7 Lee SH, Kim KA, Park YG, Seong IW, Kim MJ, Lee YJ. Identification and partial characterization of a novel hemolysin from *Leptospira interrogans* serovar lai. *Gene* 2000; 254: 19–28.
 - 8 Ren SX, Fu G, Jiang XG, Zeng R, Miao YG, Xu H, *et al*. Unique physiological and pathogenic features of *Leptospira interrogans* revealed by whole genome sequencing. *Nature* 2003; 422: 888–93.
 - 9 SWISS-PROT/TrEMBL databases. Europe: Swiss Institute of Bioinformatics. c2004. Available from: <http://www.expasy.org/sprot> and <http://www.expasy.org/TrEMBL>
 - 10 Pfam database. St Louis: Washington University in St Louis. c2004. Available from: <http://pfam.wustl.edu/hmmsearch.shtml>
 - 11 PROSITE database. Europe: Swiss Institute of Bioinformatics. c2004. Available from: <http://us.expasy.org/prosite>
 - 12 ProDom database. Europe: Laboratoire des Interactions Plantes Microorganismes. c2004. Available from: <http://prodes.toulouse.inra.fr/prodom/current/html/home.php>
 - 13 Jpred² database. Scotland: University of Dundee. c2004. Available from: <http://www.compbio.dundee.ac.uk/~www-jpred/submit.html>
 - 14 Alexander AD, Smith OH, Hiatt CW, Gleiser CA. Presence of hemolysin in cultures of pathogenic leptospire. *Proc Soc Exp Biol Med* 1956; 91: 205–11.
 - 15 Bauer DC, Morse EV. Variation and hemolysin production in relation to virulence of *Leptospira pomona*. *Proc Soc Exp Biol Med* 1958; 98: 505–8.
 - 16 Cloning of a hemolysin gene from *Leptospira interrogans* serovar hardjo. *Infect Immun* 1989; 57: 2588–90.
 - 17 Fussle R, Bhakdi S, Sziegoleit A, Trantum-Jensen J, Kranz T, Wellensiek HJ. On the mechanism of membrane damage by *Staphylococcus aureus* alpha-toxin. *J Cell Biol* 1981; 91: 83–94.
 - 18 Bhakdi S, Trantum-Jensen J. Damage to cell membranes by pore-forming bacterial cytolysins. *Prog Allergy* 1988; 40: 1–43.