

Full-length article

Effects of huperzine A on secretion of nerve growth factor in cultured rat cortical astrocytes and neurite outgrowth in rat PC12 cells¹

Li-li TANG, Rui WANG, Xi-can TANG²

State Key Laboratory of Drug Research, Shanghai Institute of Materia Medica, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 201203, China

Key words

huperzine A; nerve growth factor; neurite outgrowth; acetylcholinesterase inhibitors

¹ Project supported by the National Natural Science Foundation of China (No 30123005) and the Ministry of Science and Technology of China (No 2004CB518907).

² Correspondence to Prof Xi-can TANG.
Phn 86-21-5492-0510
Fax 86-21-5080-7088
E-mail xctang@mail.shnc.ac.cn

Received 2004-12-27
Accepted 2005-03-15

doi: 10.1111/j.1745-7254.2005.00130.x

Abstract

Aim: To study the effects of huperzine A (HupA) on neurotogenic activity and the expression of nerve growth factor (NGF). **Methods:** After being treated with 10 $\mu\text{mol/L}$ HupA, neurite outgrowth of PC12 cells was observed and counted under phase-contrast microscopy. Mitogenic activity was assayed by [³H]thymidine incorporation. Cell cytotoxicity was evaluated by lactate dehydrogenase (LDH) release. AChE activity, mRNA and protein expression were measured by the Ellman's method, RT-PCR, and Western blot, respectively. NGF mRNA and protein levels were determined by RT-PCR and ELISA assays. **Results:** Treatment of PC12 cells with 10 $\mu\text{mol/L}$ HupA for 48 h markedly increased the number of neurite-bearing cells, but caused no significant alteration in cell viability or other signs of cytotoxicity. In addition to inhibiting AChE activity, 10 $\mu\text{mol/L}$ HupA also increased the mRNA and protein levels of this enzyme. In addition, following 2 h exposure of the astrocytes to 10 $\mu\text{mol/L}$ HupA, there was a significant up-regulation of mRNA for NGF and P75 low-affinity NGF receptor. The protein level of NGF was also increased after 24 h treatment with HupA. **Conclusion:** Our findings demonstrate for the first time that HupA has a direct or indirect neurotrophic activity, which might be beneficial in treatment of neurodegenerative disorders such as Alzheimer disease.

Introduction

Nerve growth factor (NGF) is a member of the neurotrophin family that promotes the survival and outgrowth of central cholinergic neurons^[1]. The decrease in trophic support for the neurons in the aging brain is associated with neuronal death and appearance of neurodegenerative disorders such as Alzheimer disease (AD)^[2]. A case study showed that topical application of NGF into the brain of one AD patient relieved symptoms of dementia^[3]. However, the clinical utility of NGF is limited by an inability to cross the blood-brain barrier, necessitating invasive neurosurgical procedures for administration. In contrast, pharmacological stimulation of endogenous NGF synthesis or mimicking of NGF activity by compounds that penetrate membrane barriers may provide an alternative means to provide equivalent trophic actions in the central nervous system.

Accumulated data suggest that cholinergic mechanisms are involved in the regulation of NGF synthesis and release. It has been suggested that cholinergic activity in basal forebrain neurons may stimulate NGF synthesis in appropriate target areas during early postnatal development^[4]. Additionally, substances that augment cholinergic function, such as nicotine, muscarinic receptor agonists, and acetylcholine-releasing agents, can increase NGF expression and synthesis^[5,6]. Although there have been no reports that acetylcholinesterase (AChE) inhibitors elevate NGF production, some AChE inhibitors are known to exert NGF-like activity by potentiating the neurotogenic effect of NGF^[7] or increasing choline acetyltransferase (ChAT) activity^[8]. These actions appear to be independent of AChE inhibition. Such findings led us to suspect that the neurotrophic effects of huperzine A (HupA), a novel acetylcholinesterase inhibitor isolated from the Chinese herb *Huperzia serrata*, involve

NGF-like activity, directly or indirectly.

HupA has been proved to be one of the most promising new agents for AD therapy^[9]. Our previous studies showed that, in addition to its potent inhibition of AChE, HupA exhibited neuroprotective effects both *in vivo* and *in vitro*^[10,11]. For example, just as NGF can ameliorate neuronal degeneration in rat cerebral cortex and hippocampus after ischemic insult^[12], HupA will attenuate ischemic damage from transient global ischemia in gerbils and hypoxic-ischemia (HI) in neonatal rats^[13,14]. A multicenter, randomized, double-blind, placebo-controlled clinical trial in China proved that HupA markedly improved the cognitive function of vascular dementia (VD)^[15-17]. In view of these findings it is natural to ask if HupA might influence intrinsic neurotrophic factors. The aim of this study was to examine the effects of HupA on neurogenesis in rat pheochromocytoma cells and on the expression and secretion of NGF in primary cultures of rat cortical astrocytes.

Materials and methods

Materials HupA, provided by the Department of Phytochemistry at Shanghai Institute of Materia Medica is a colorless powder with mp 230 °C, and purity >99%. It was dissolved and diluted in phosphate-buffered saline (PBS). RPMI-1640 medium, DMEM/F12 medium, fetal bovine serum, horse serum, and N-2 supplement were purchased from Gibco (CA, USA). Cell cytotoxicity Kit, NGF E_{max}[®] Immunoassay System, Reverse Transcription System were purchased from Promega (Madison, WI, USA). Mouse monoclonal anti-AChE antibody was purchased from BD Biosciences (CA, USA). TRIzol reagent was purchased from Invitrogen (CA, USA). ECL kit was purchased from Pierce (Rockford, USA).

PC12 cell culture and neurite outgrowth assay Undifferentiated PC12 rat pheochromocytoma cells were obtained from ATCC. The cells were maintained in RPMI-1640 media containing 10% FCS, 5% HS, 100 kU/L penicillin, and 100 kU/L streptomycin in a humidified atmosphere of 95% air and 5% CO₂ at 37 °C. For each experiment, cells (3.5×10⁴ cells/well) were seeded into a 12-well collagen-coated plate and cultured overnight. Subsequently the cells were washed once with PBS and transferred to fresh serum-free DMEM containing an N-2 supplement. After a 2-h incubation with serum-free DMEM, NGF or HupA was added to the cultures, which were incubated for a further 48 h. Neurite formation was examined under a phase-contrast microscope, and processed longer than one cell diameter were scored as neurites. The percentage of neurite-bearing cells in relation to the total number of cells was examined in four fields from

each of the eight culture wells per treatment condition.

Mitogenic activity and cytotoxicity assay To analyze the effects of HupA on mitogenic activity, PC12 cells were seeded into 96-well plate at a density of 1×10⁴ cell/well. After incubation of the cells with 10 μmol/L HupA for 48 h at 37 °C, [³H]thymidine (37 000 Bq) was added to the cultures and incubation was continued for another 8 h. The cells were harvested onto filters using a cell harvester and the retained radioactivity was determined in a scintillation counter. HupA cytotoxicity was evaluated by an assay kit, following the manufacture's protocol to detect lactate dehydrogenase (LDH) release.

AChE activity assay AChE activity was measured by a standard spectrophotometric method^[18].

Western blot analysis Changes in AChE levels were assessed by Western blot. For this purpose, cultured cells harvested 48 h after treatment with HupA were lysed in 1×SDS PAGE gel loading buffer [Tris-HCl 50 mmol/L pH 6.8, DTT 100 mmol/L, 2% (w/v) SDS, 10% (w/v) glycerol, 0.1% Bromophenol Blue] and boiled in a water bath for 10 min. Equal amounts of protein (40 μg) were loaded in each gel lane and, after electrophoresis, proteins were transferred to a nitrocellulose membrane. The membranes were blocked with TBST containing 5% non-fat milk and then were incubated at 4 °C overnight with primary antibody (a mouse monoclonal antibody raised against human AChE and crossreactive with rat AChE). Finally the blots were incubated with HRP-conjugated anti-mouse IgG at 37 °C for 2 h and target protein bands were detected by the ECL method according to the manufacture's instruction.

Astrocytes cultures and experimental treatment For studies of NGF expression, primary cultures of astrocytes were prepared from 1-day-old neonatal Sprague-Dawley rats. Cerebral cortices were stripped of meninges and dissected in Ca²⁺-, Mg²⁺-free D-Hanks' solution. Next the samples were trypsinized and passed through a monofilament mesh (pore size 80 μm). Cells were collected by centrifugation and resuspended in DMEM/F12 containing 10% FCS, 100 kU/L penicillin and 100 kU/L streptomycin for culture in a humidified atmosphere (95% air and 5% CO₂) at 37 °C. Astrocytes were grown to confluence and oligodendrocyte and microglial cells were removed by shaking and washing with cold D-Hanks' solution. The astrocytes were then trypsinized and plated onto 24-well plates at a density of 1×10⁴ cell/cm², washed with D-Hanks' solution, and exposed for 24 h to serum-free medium. After this incubation, culture medium was collected and centrifuged to remove cell debris.

NGF content assay Released NGF was measured in the supernatant by a two-site enzyme-linked immunosorbent

assay (ELISA) using the NGF E_{max}[®] Immunoassay system.

RT-PCR analysis RT-PCR was performed to determine the expression of mRNA for AChE in PC12 cells and for NGF and P75^{NTR} in astrocytes. Total cellular RNA was isolated using TRIzol reagent following the manufacturer's protocols and quantified by absorbance at 260 nm. RNA purity was determined using the A₂₆₀/A₂₈₀ ratio (average ratio > 1.85). Total RNA of each sample was reverse-transcribed into cDNA using Reverse Transcription System. The cDNAs were amplified with the following specific primers. AChE: 5'-TCTTTGCTCAGCGACTTA-3' (upstream), 5'-GTCACAGG-TCTGAGCATCT-3' (downstream); NGF: 5'-CTTCAGCATT-CCCTTGACAC-3' (upstream), 5'-AGCCTTCCTGCTGAGCA-CACA-3' (downstream); P75^{NTR}: 5'-AGCCAACCAGACCG-TGTGTG-3' (upstream), 5'-TTGCA GCTGTTCCACCTCTT-3' (downstream); β-actin: 5'-CCTGCGTCTGGACCTGGCTG-3' (upstream), 5'-CTCAGGAGGAGCAATGATCT-3' (downstream). Amplifications were performed as follows: AChE: 30 cycles, 94 °C for 45 s, 56 °C for 45 s, 72 °C for 45 s; NGF: 24 cycles, 94 °C for 30 s, 60 °C for 30 s, and 72 °C for 60 s; P75^{NTR}: 25 cycles, 94 °C for 45 s, 55 °C for 30 s, and 72 °C for 90 s; β-actin: 30 cycles, 94 °C for 45 s, 57 °C for 45 s, and 72 °C for 60 s. The PCR products were normalized in relation to standards of β-actin mRNA.

Statistical analysis Data were expressed as mean±SEM. Statistical analysis was performed by one-way analysis of variance (ANOVA) followed by Duncan's multiple-range test, with P<0.05 as the significant level.

Results

Mitogenic activity and cytotoxicity of HupA on PC12 cells

There was no effect on cell proliferation during a 48-h exposure of PC12 cells to HupA 10 μmol/L. Likewise, measures of LDH release showed that HupA 10 μmol/L did not induce any cytotoxic effects (Figure 1).

Effects of HupA on neurite outgrowth Most PC12 cells displayed a flat, polygonal, undifferentiated morphology after incubation under control conditions for 48 h, and only 5.5% of them exhibited neurites. However, after incubation with HupA 10 μmol/L for 48 h more cells differentiated and nearly twice as many (10.4%) were found to bear neuritis. Moreover, 14.2% of PC12 cells exhibited neuritis after incubation with 2 μg/L NGF (Figure 2).

Effects of HupA on activity, expression, and protein levels of AChE In PC12 cells the activity of AChE was inhibited substantially by HupA treatment. The expression of AChE was also affected. AChE mRNA was increased at 6 h and 10 h. Consistent with the late rise in mRNA, AChE protein levels rose after a 48-h incubation with HupA (Figure 3).




Figure 1. Effects of huperzine A on PC12 cell proliferation and viability. (A) Proliferation was measured by [³H]thymidine incorporation. The [³H]thymidine incorporation of control cells was 62690±1242 cpm per 1×10⁴ cells. n=33. (B) Cytotoxicity was assessed by the release of LDH. The average absorbance of control cell supernatant was 0.214±0.009; of total control cell lysate was 2.68±0.11. n=18. Mean±SEM.

Effects of HupA on the expression and secretion of NGF

Effects of HupA on NGF and P75^{NTR} mRNA were determined by RT-PCR in primary rat cortical astrocytes, a cell type known to express the corresponding proteins. Figure 4A represents an example of RT-PCR products visualized after electrophoresis in 1.5% agarose gel containing ethidium bromide. The results suggested that NGF mRNA transcripts were slightly upregulated at 2, 4, and 6 h, while the levels of mRNA for P75^{NTR} appeared greatly increased. Moreover, an ELISA for NGF protein showed a large and statistically significant increase in cultures incubated with HupA 10 μmol/L for 24 h when compared to controls incubated for the same length of time in normal medium (Figure 4B).

Discussion

PC12 cells cannot secrete NGF, but they have NGF receptors and can respond to NGF. In the presence of NGF, PC12 cells cease cell division, differentiate into sympathetic neuron-like cells, and extend neuritis. So it is a putative model to




Figure 2. Effects of huperzine A on neurite outgrowth in PC12 cells. Top panels, representative phase contrast micrographs ($\times 200$). PC12 cells were incubated for 48 h under control conditions (A), with 1 $\mu\text{mol/L}$ HupA (B), with 10 $\mu\text{mol/L}$ HupA (C), and with 2 $\mu\text{g/L}$ NGF (D). Bottom panels, quantitative effects of HupA on neurite outgrowth. Three independent experiments were carried out. $n=8$ for each independent experiment. Mean \pm SEM. $^{\circ}P<0.01$ vs control.

determine the neurotogenic activity^[19]. Different from PC12 cells, astrocytes can secrete NGF. In the injured brain, for example, astrocytes play an important role in neurotrophic support^[20]. So the two different cells were used to determine the direct or indirect neurotrophic activity of HupA. In the present study, HupA was demonstrated to increase neurite outgrowth from undifferentiated PC12 cells and to enhance the expression and secretion of NGF in primary astrocytes. These findings indicate that HupA is not only an effective AChE inhibitor, but it also possesses NGF-inducing activity and an ability to induce certain NGF-like effects. The question arises whether these effects are closely linked, and what




Figure 3. Effects of huperzine A on AChE mRNA expression and immunoreactive protein levels and inhibition of AChE activity. (A) Representative RT-PCR pattern. Lane 1, control; lanes 2–5: 10 $\mu\text{mol/L}$ HupA incubated for 2, 4, 6 and 10 h, respectively. (B) Representative Western blot pattern. Lane 1: control; lane 2: treated with 10 $\mu\text{mol/L}$ HupA for 48 h. (C) Inhibition of AChE activity in PC12 cells after treatment with 10 $\mu\text{mol/L}$ huperzine A for 48 h. AChE activity of control was (1.80 ± 0.15) OD_{440} values per mg protein. $n=6$. Mean \pm SEM. Note that the true level of inhibition during incubation is probably underestimated, owing to the reversible nature of HupA interaction with AChE. $^{\circ}P<0.01$ vs control.

kind of relation they have to AChE. It has been reported that AChE plays an important role in neuronal proliferation and differentiation during early development of the central and peripheral nervous system. Neural AChE typically appears while axons are growing and before synaptic connections form^[21]. Accumulating evidence indicates that AChE may influence neurite outgrowth through a non-catalytic mechanism such as cell-cell or cell-substratum adhesion^[22]. It has been observed that the trophic activity of AChE is blocked by inhibitors that interact with the peripheral anionic site (PAS) but not by inhibitors that interact only with the active site, deep within the catalytic gorge^[23,24]. Such findings sug-




Figure 4. Effects of huperzine A on NGF and P75^{NTR} mRNA expression and NGF secretion in rat cortical astrocytes. (A) Representative RT-PCR pattern. Lane 1: control; Lanes 2–4: 10 μmol/L HupA incubated for 2, 4, and 6 h, respectively. The PCR products were normalized with reference to β-actin mRNA. (B) Huperzine A enhances NGF protein level in rat cortical astrocytes. Astrocytes serum deprived for 24 h were incubated with vehicle or 10 μmol/L HupA for another 24 h. NGF protein level was determined directly in the culture medium by NGF-ELISA. *n*=9. Mean±SEM. ^b*P*<0.05 vs control.

gest strongly that surface features of AChE rather than catalytic activity are responsible for its trophic effect. Hence, it is not contradictory to find that HupA affected neurite outgrowth in a positive rather than negative manner despite causing marked AChE inhibition. In fact, the differentiation-promoting effect of HupA may well reflect increased amounts of AChE protein. In other cell lines and to some extent *in vivo*, increased neurite outgrowth has been seen to parallel the level of AChE expression^[25–27]. Conversely, decreasing expression of AChE expression, using antisense techniques, reduced outgrowth^[28]. Our present results showed that AChE mRNA expression and protein levels were significantly up-regulated after treatment with HupA. This finding is consistent with other reports of increased AChE gene expression after exposure to AChE inhibitors^[29,30]. It remains to be determined whether feedback regulation of AChE synthesis is involved in these changes.

Previous observations suggest that NGF regulates the phenotype and survival of basal forebrain cholinergic neurons^[31] and protects hippocampal and cortical neurons against excitotoxic and ischemic damage^[32]. Neurons surviving from transient ischemia highly expressed P75^{NTR}, suggesting that this low affinity neurotrophin receptor could contribute to the cytoprotective effect of NGF^[33]. Our results showed that HupA enhanced the expression and secretion of NGF as well as increasing P75^{NTR} mRNA level in astrocytes. We conclude that these two responses may be key to the neuroprotective effects that have been observed *in vitro* and *in vivo* after treatment with HupA.

The exact mechanism by which HupA increases NGF secretion remains to be determined. There is evidence that cholinergic and adrenergic mechanisms as well as PKC activation all affect NGF gene expression in astrocytes. Our previous studies have demonstrated that HupA has enhancing effects on PKC^[34], and on cholinergic and adrenergic systems^[35]. Such effects may participate in the modulation of NGF synthesis. This provides another possible mechanism for HupA to promote survival of damaged neurons, which may interact synergistically with other pathways to exert the neuroprotective effect of this drug.

In the present study, HupA induced the NGF synthesis of cultured astrocytes and enhanced the neurite outgrowth of undifferentiated PC12 cells *in vitro*. These effects provide the possibility that HupA increase NGF-induced enhancement of neurons survival and their function improvement that was helpful in the rescue of injured neurons.

In summary, our study has demonstrated for the first time that HupA induces neurite outgrowth in PC12 cells and stimulates expression of NGF, P75 mRNA, and secretion of NGF in cultured rat cortical astrocytes. These effects might be helpful to restore and maintain neural cells in neurodegenerative disease.

Acknowledgement

The authors wish to thank Prof Stephen BRIMIJOIN (Mayo Clinic, USA) for helpful discussion and English revision to the manuscript.

References

- 1 Hefti F, Hartikka J, Knusel B. Function of neurotrophic factors in the adult and aging brain and their possible use in the treatment of neurodegenerative disease. *Neurobiol Aging* 1989; 10: 515–33.
- 2 Mufson EJ, Bothwell M, Kordower JH. Loss of nerve growth factor receptor-containing neurons in Alzheimer’s disease: a quantitative analysis across subregions of the basal forebrain. *Exp*

- Neurol 1989; 105: 221–32.
- 3 Olson L, Nordberg A, von Holst H. Nerve growth factor affects [¹⁴C]nicotine binding, blood flow, EEG, and verbal episodic memory in an Alzheimer patient (case report). *J Neural Transm (P-D Sect)* 1992; 4: 79–95.
 - 4 da Penha Berzaghi M, Cooper J, Castren E, Zafra F, Sofroniew M, Thoenen H, *et al*. Cholinergic regulation of brain-derived neurotrophic factor (BDNF) and nerve growth factor (NGF) but not neurotrophin-3 (NT-3) mRNA levels in the developing rat hippocampus. *J Neurosci* 1993; 13: 3818–26.
 - 5 French SJ, Humby T, Horner CH, Sofroniew MV, Rattray M. Hippocampal neurotrophin and trk receptor mRNA levels are altered by local administration of nicotine, carbachol and pilocarpine. *Brain Res Mol Brain Res* 1999; 67: 124–36.
 - 6 Ghelardini C, Galeotti N, Bartolini A, Furukawa S, Nitta A, Manetti D, *et al*. Memory facilitation and stimulation of endogenous nerve growth factor synthesis by the acetylcholine releaser PG-9. *Jpn J Pharmacol* 1998; 78: 245–51.
 - 7 Shigeta K, Ootaki K, Tatemoto H, Nakanishi T, Inada A, Muto N. Potentiation of nerve growth factor-induced neurite outgrowth in PC12 cells by a *Coptidis Rhizoma* extract and protoberberine alkaloids. *Biosci Biotech Biochem* 2002; 66: 2491–4.
 - 8 Kato K, Hayako H, Ishihara Y, Marui S, Iwane M, Miyamoto M. TAK-147, an acetylcholinesterase inhibitor, increases choline acetyltransferase activity in cultured rat septal cholinergic neurons. *Neurosci Lett* 1999; 260: 5–8.
 - 9 Tang XC, He XC, Bai DL. Huperzine A : a novel acetylcholinesterase inhibitor. *Drugs Fut* 1999; 24: 647–63.
 - 10 Xiao XQ, Wang R, Tang XC. Huperzine A and tacrine attenuate beta-amyloid peptide-induced oxidative injury. *J Neurosci Res* 2000; 61: 564–9.
 - 11 Wang R, Zhang HY, Tang XC. Huperzine A attenuates cognitive dysfunction and neuronal degeneration caused by beta-amyloid protein (1-40) in rat. *Eur J Pharmacol* 2001; 421: 149–56.
 - 12 Buchan AM, Williams L, Bruederlin B. Nerve growth factor: pretreatment ameliorates ischemic hippocampal neuronal injury. *Stroke* 1990; 21: 177.
 - 13 Zhou J, Zhang HY, Tang XC. HupA attenuates cognitive deficits and hippocampal neuronal damage after transient global ischemia in gerbils. *Neurosci Lett* 2001; 313: 137–40.
 - 14 Wang LS, Zhou J, Shao XM, Tang XC. Huperzine A attenuates cognitive deficits and brain injury in neonatal rats after hypoxia-ischemia. *Brain Res* 2002; 949: 162–70.
 - 15 Li HN, Min QY. Huperzine A improved the cognition of vascular dementia: a report of 30 patients in therapeutics. *Chin J Clin Rehabil* 2001; 5: 59.
 - 16 Ma YX, Zhu Y, Gu YD, Yu ZY, Y SM, Ye YZ. Double-blind trial of huperzine-A (HUP) on cognitive deterioration in 314 cases of benign senescent forgetfulness, vascular dementia, and Alzheimer's disease. *Ann NY Acad Sci* 1998; 854: 506–7.
 - 17 Pi X, Liu Y, Jiang ZY, Hu XQ, Zhu GZ. Clinical observation on treatment of light and moderate vascular dementia with meclofenoxate plus huperzine A. *Shanghai Med Pharm J* 2004; 255: 409–11.
 - 18 Wang YE, Yue DX, Tang XC. Anti-cholinesterase activity of huperzine A. *Acta Pharmacol Sin* 1986; 7: 110–3.
 - 19 Greene LA, Tischler AS. PC12 pheochromocytoma cultures in neurobiological research. *Adv Cell Neurobiol* 1982; 3: 373–414
 - 20 Lu B, Yokoyama M, Dreyfus CF, Black IB. NGF gene expression in actively growing brain glia. *J Neurosci* 1991; 11: 318–26.
 - 21 Drews U. Cholinesterase in embryonic development. *Prog Histochem Cytochem* 1975; 7: 1–53
 - 22 Brimijoin S, Koenigsberger C. Cholinesterases in neural development: new findings and toxicologic implications. *Environ Health Perspect* 1999; 107 (Suppl 1): 59–64.
 - 23 Munoz FJ, Aldunate R, Inestrosa NC. Peripheral binding site is involved in the neurotrophic activity of acetylcholinesterase. *Neuroreport* 1999; 10: 3621–5.
 - 24 Sharma KV, Koenigsberger C, Brimijoin S, Bigbee JW. Direct evidence for an adhesive function in the noncholinergic role of acetylcholinesterase in neurite outgrowth. *J Neurosci Res* 2001; 63: 165–75.
 - 25 Karpel R, Sternfeld M, Ginzberg D, Guhl E, Graessmann A, Soreq H. Overexpression of alternative human acetylcholinesterase forms modulates process extensions in cultured glioma cells. *J Neurochem* 1996; 66: 114–23.
 - 26 Koenigsberger C, Chiappa S, Brimijoin S. Neurite differentiation is modulated in neuroblastoma cells engineered for altered acetylcholinesterase expression. *J Neurochem* 1997; 69: 1389–97.
 - 27 Layer PG, Willbold E. Novel functions of cholinesterases in development, physiology and disease. *Prog Histochem Cytochem* 1995; 29: 1–94.
 - 28 Grifman M, Soreq H. Differentiation intensifies the susceptibility of pheochromocytoma cells to antisense oligodeoxynucleotide-dependent suppression of acetylcholinesterase activity. *Antisense Nucleic Acid Drug Dev* 1997; 7: 351–9.
 - 29 Kaufer D, Friedman A, Seidman S, Soreq H. Acute stress facilitates long-lasting changes in cholinergic gene expression. *Nature* 1998; 393: 373–7.
 - 30 Chiappa S, Padilla S, Koenigsberger C, Moser V, Brimijoin S. Slow accumulation of acetylcholinesterase in rat brain during enzyme inhibition by repeated dosing with chlorpyrifos. *Biochem Pharmacol* 1995; 49: 955–63.
 - 31 Whittemore SR, Seiger A. The expression, localization and functional significance of beta-nerve growth factor in the central nervous system. *Brain Res* 1987; 434: 439–64.
 - 32 Semkova I, Wolz P, Schilling M, Krieglstein J. Selegiline enhances NGF synthesis and protects central nervous system neurons from excitotoxic and ischemic damage. *Eur J Pharmacol* 1996; 315: 19–30.
 - 33 Lee TH, Abe K, Kogure K, Itoyama Y. Expressions of nerve growth factor and p75 low affinity receptor after transient forebrain ischemia in gerbil hippocampal CA1 neurons. *J Neurosci Res* 1995; 41: 684–95.
 - 34 Zhang HY, Yan H, Tang XC. HupA enhances the level of secretory amyloid precursor protein and protein kinase C-alpha in intracerebroventricular beta-amyloid-(1-40) infused rats and human embryonic kidney 293 Swedish mutant cells. *Neurosci Lett* 2004; 360: 21–4.
 - 35 Ou LY, Tang XC, Cai JX. Effect of HupA on working memory in reserpine- or yohimbine-treated monkeys. *Eur J Pharmacol* 2001; 433: 151-6.