

Full-length article

Peroxisome proliferator-activated receptor γ ligands induce cell cycle arrest and apoptosis in human renal carcinoma cell lines¹

Feng-guang YANG², Zhi-wen ZHANG³, Dian-qi XIN², Chang-jin SHI², Jie-ping WU², Ying-lu GUO^{2,4}, You-fei GUAN^{3,4}

²Department of Molecular Biology, Institute of Urology, Peking University First Hospital; ³Department of Physiology and Pathophysiology, Peking University Health Science Center, Beijing 100034, China

Key words

renal cell carcinoma; peroxisome proliferator-activated receptor γ ; cell cycle; apoptosis; cyclin; cdk

¹ Project partly supported by National Natural Science Foundation of China (No 30250002 to Yin-lu GUO and No 30271521, No 30340084 and No G2000056908 to You-fei GUAN) and a grant from Peking University "211" Program (to You-fei GUAN).

⁴ Correspondence to Dr You-fei GUAN and Ying-lu GUO.

Phn 86-10-8280-1447.

E-mail youfeiguan@bjmu.edu.cn

Received 2004-12-08

Accepted 2005-02-05

doi: 10.1111/j.1745.7254.2005.00103.x

Abstract

Aim: To study the effect of peroxisome proliferator-activated receptor γ (PPAR γ) ligands on cell proliferation and apoptosis in human renal carcinoma cell lines. **Methods:** The expression of PPAR γ was investigated by reverse transcriptase polymerase chain reaction (RT-PCR), Western blot and immunohistochemistry. The effect of thiazolidinedione (TZD) PPAR γ ligands on growth of renal cell carcinoma (RCC) cells was measured by MTT assay and flow cytometric analysis. Cell death ELISA, Hoechst 33342 fluorescent staining and DNA ladder assay were used to observe the effects of PPAR γ ligands on apoptosis. Regulatory proteins of cell cycle and apoptosis were detected by Western blot analysis. **Results:** PPAR γ was expressed at much higher levels in renal tumors than in the normal kidney (2.16 ± 0.85 vs 0.90 ± 0.73 ; $P < 0.01$). TZD PPAR γ ligands inhibited RCC cell growth in a dose-dependent manner with IC₅₀ values of 7.08 μ mol/L and 11.32 μ mol/L for pioglitazone, and 5.71 μ mol/L and 8.38 μ mol/L for troglitazone in 786-O and A498 cells, respectively. Cell cycle analysis showed a G₀/G₁ arrest in human RCC cells following 24-h exposure to TZD. Analysis of cell cycle regulatory proteins revealed that TZD decreased the protein levels of proliferating cell nuclear antigen, pRb, cyclin D₁, and Cdk4 but increased the levels of p21 and p27 in a time-dependent manner. Furthermore, high doses of TZD induced massive apoptosis in renal cancer cells, with increased Bax expression and decreased Bcl-2 expression. **Conclusion:** TZD PPAR γ ligands showed potent inhibitory effect on proliferation, and could induce apoptosis in RCC cells. These results suggest that ligands for PPAR γ have potential antitumor effects on renal carcinoma cells.

Introduction

Peroxisome proliferator-activated receptor γ (PPAR γ) belongs to the superfamily of nuclear hormone receptor transcription factors^[1–3]. PPAR γ forms a heterodimer with another nuclear receptor, retinoid X receptor α (RXR α)^[4]. In the presence of their ligands, PPAR γ /RXR α heterodimer binds to a specific DNA sequence designated peroxisome proliferating response element (PPRE) located in the promoter region of PPAR γ target genes and modulates their transcription^[5]. Thus far, hundreds of PPAR γ target genes have been identified and most of them are involved in adipogenesis^[6], glucose metabolism^[7] and angiogenesis^[8]. Since the dis-

covery of specific ligands for PPAR γ , including synthetic antidiabetic thiazolidinediones (TZD) and endogenous 15-deoxy-*D*^{12,14}-PGJ₂ (15dPGJ₂) and some polyunsaturated fatty acids, accumulating evidence using many experimental systems suggests that PPAR γ plays an important role in carcinogenesis in both adipose and nonadipose cells. It has been known that PPAR γ expression is up-regulated in many tumor tissues and its ligands, including pioglitazone, troglitazone and roglitazone, can induce apoptosis and exert antiproliferative effects in human colon cancer^[9], breast cancer^[10], pituitary adenomas^[11], gastric cancer^[12] and bladder cancer^[13]. Moreover, PPAR γ may play an important role in inducing cell differentiation or growth inhibition in patients

with liposarcoma^[14] and prostate cancer^[15]. Therefore activation of PPAR γ seems to be a potential approach for treatment of some malignant tumors.

Renal cell carcinoma (RCC) is the most common renal tumor and the third malignancy within urological oncology. It makes up approximately 2%–3% of all adult malignancies. At present, more than 50% of all RCC are found incidentally, which results in a high proportion of patients with progressive metastasis at the time of diagnosis^[16,17]. Surgery remains the only effective therapeutic option. Although combination of chemotherapy and/or radiotherapy might help in survival rate, it is considered to be of limited value for the treatment of RCC. Therefore, improvement of life span in patients with renal cell carcinoma greatly depends on the identification of novel treatment strategies. An alternative treatment may include activation of PPAR γ , as PPAR γ has been shown to be expressed in the kidney^[18], and its ligands can induce cell growth inhibition and terminal differentiation in many other malignancies^[9–14].

In the present study, we examined the expression of PPAR γ in human primary RCC and RCC cell lines, and determined the biological events of PPAR γ activation in inducing RCC cell cycle arrest and apoptosis. We have also explored the molecular mechanism through which PPAR γ agonists inhibit cell growth and induce cell death in RCC cell lines.

Materials and methods

Chemicals Pioglitazone and troglitazone were kindly gifted by Park-Davis Pharmaceutical Research (Ann Arbor, MI, USA) and Takeda Chemical Industries (Osaka, Japan). All other chemicals were purchased from Sigma (St Louis, MO, USA). Stock solutions of pioglitazone or troglitazone were made at 100 mmol/L concentration in dimethyl sulfoxide and added to the culture medium at the final concentration of less than 0.1%.

Cell lines and culture conditions Human RCC-derived cell lines 786-O and A498, human proximal tubular cell line (HK-2), and human mesangium cell line (HMCL) were cultured in Hepes-buffered RPMI-1640 medium (Gibco BRL, Grand Island, NY, USA) supplemented with 10% FBS, 100 kU/mL penicillin G, 100 g/L streptomycin, and 2 mmol/L *l*-glutamine in a humidified 5% CO₂ atmosphere at 37 °C. The media were changed every 3 d, and the cells were separated by trypsinization using trypsin/edetic acid when they reached 90% confluence.

Tumor specimens and immunohistochemistry Specimens were obtained from 120 patients with RCC and 20 patients with normal kidney (NK) who underwent total

nephroureterectomy as a result of ureteral cancer or trauma. All patients were treated at the Institute of Urology in Peking University between June 1999 and June 2001. No patients had received irradiation or chemotherapy prior to surgery. Tumor samples with the highest nuclear grade were selected. All tissues were preserved in 10% formalin serially sectioned onto microscope slides with a thickness of 4 μ m/L. Sections were immunostained with polyclonal antibody to human PPAR γ (1:100, Santa Cruz Biotechnology, Santa Cruz, CA, USA) using avidin-biotin-complex-peroxidase and counterstained with hematoxylin. Negative control slides were prepared by omitting the primary antibody. The intensity of immunostaining and the ratio of the positive cells were roughly graded into four scores (0, 1, 2, 3, and 4) by two observers who did not know the origin of the samples on two occasions. The score 4 was defined as maximum intensity of immunostaining throughout the section, while the score 0 implied that staining was absent throughout the specimen.

Measurement of PPAR γ mRNA by reverse transcription-polymerase chain reaction Total RNA from 15 RCC tissues, 10 NK tissues, RCC cell lines 786-O and A498, HK-2 cells and HMCC (normal kidney derived mesangium cell line). Cells was isolated using TRIzol reagent (Invitrogen, Carlsbad, CA, USA). Contaminated genomic DNA in total RNA was removed by RNase-free DNase I digestion. Reverse transcription was performed in a total volume of 25 μ L using 3 μ g of total RNA, oligo-dT primer and MMLV reverse transcriptase (Amersham, Buckinghamshire, UK). Primers used for the amplification of PPAR γ cDNA were synthesized at the SBS Gene Company (Beijing, China). The sense primer was 5'-AGAGATGCCATTCTGGCC-3' and antisense primer was 5'-GTGGAGTAGAAATGCTGGAGA-3'. Polymerase chain reaction (PCR) amplification yielded a PCR product of 130 bp. Primers used for amplifying a 500-bp of β -actin cDNA as an internal control were as follows: 5'-ACTGACTACCTCATGAAGATC-3' (sense) and 5'-CGTCATACTCCTGCTTGC-3' (anti-sense). The PCR condition was 40 s (denaturation) at 94 °C, 35 s (annealing) at 60 °C and 35 s (extension) at 72 °C for 33 cycles. PCR products were separated in 1.5% agarose gel and visualized after ethidium bromide staining.

Growth inhibition of RCC cell lines by PPAR γ agonists The effect of PPAR γ ligands on cell proliferation of RCC cells was determined using MTT assay^[13]. Briefly, cells of 0.5×10^4 cells/well were inoculated into a 96-well plate (Costar, Cambridge, MA, USA), treated with pioglitazone or troglitazone at various concentrations. After an incubation for 24 h, 20 μ L/well 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT, 5 g/L) was added to each well, the me-

dium was then removed, and 200 μL of 0.04 mol/L HCl in isopropanol was added to dissolve the reduced formazan product. The plate was read in a microplate reader (model 3550, BIO-RAD, Richmond, CA, USA) at 590 nm.

Flow cytometry assay RCC cell lines 786-O and A498 treated with pioglitazone (20 $\mu\text{mol/L}$) and troglitazone (30 $\mu\text{mol/L}$) for 24 h were harvested and washed with PBS. Cells of 1×10^6 were fixed in 75% ethanol at 4 °C for 24 h, washed in PBS, resuspended in 0.5–1.0 mL PBS containing 100 mg/L RNase A at 37 °C for 30 min, incubated in 10 mg/L propidium iodide in dark at 4 °C for 30 min, and sorted in a Coulter EPICS-XL Cytometer (Becton Dickinson and Beckman-Coulter, San Jose, CA, USA).

Apoptosis assay Cytosolic histon-bound DNA fragments caused by apoptosis were detected using a cell death ELISA kit (Roche Applied Science, Mannheim, Germany) according to manufacturer's protocol. Briefly, 1×10^4 cells were seeded in a 96-well microplate and incubated for 4 h at 37 °C. Pioglitazone 0–100 $\mu\text{mol/L}$ or troglitazone was added and incubated for 48 h. Cells were then lysed with lysis buffer for 30 minutes at 25 °C. Supernatant 20 μL was transferred into the streptavidin coated multiplate, 4 μL anti-histon-biotin and 4 μL anti-DNA-POD were added to each well, and the plate was shaken for 2 h. After being washed with incubation buffer, 100 μL ABTS (2,2'-Azino-di[3-ethylbenzthiazoline-sulfonate]) solution was pipetted into each well. The absorbance was read immediately in a microplate reader (BIO-RAD, model 3550) at 405 nm. Furthermore, apoptosis of RCC cell lines induced by 70 $\mu\text{mol/L}$ troglitazone or 80 $\mu\text{mol/L}$ pioglitazone was also assayed using two other methods. Morphological changes resulting from apoptosis were determined by Hoechst 33342 staining. Cells suspended in PBS were stained with 2 mg/L Hoechst33342 and observed under fluorescence microscope using a blue filter. Cells showing cytoplasmic and nuclear shrinkage and chromatin condensation or fragmentation were defined as apoptotic cells. DNA fragmentation manifested as laddering in agarose gel was also examined. After drug treatment, the cells pellet was lysed in 10 mmol/L Tris-HCl pH 8.0, 150 mol/L NaCl, 10 mmol/L edetic acid and 0.5% SDS on ice for 10 min, and treated with RNase A and proteinase K for 1 h. Following DNA precipitation with ethanol and ammonium acetate at -20 °C for 10 h, DNA was dissolved in TE buffer, electrophoresed in 1.8% agarose gel, and visualized under UV light.

Western blot analysis Cells before and after drug treatment were extracted with lysis buffer containing 50 mmol/L Tris-HCL, pH 8.0, 150 mmol/L sodium chloride, 1% TritonX-100, 0.02% sodium azide, 100 mg/L phenylmethylsulfonyl fluoride and 1 mg/L aprotinin. The protein content in cell lysate

was determined by bicichoninic acid assay using bovine serum albumin as the standard. Cell lysate containing 50–60 μg protein was resolved by SDS-polyacrylamide gel electrophoresis and electrophoretically transferred onto a polyvinylidene difluoride membrane (Millipore, Bedford, MA, USA). The membrane was blotted by antibodies against human PPAR γ (1:500), proliferating cell nuclear antigen (PCNA; 1:1500), pRb (1:1000), p21 (1:1000), p27 (1:1000), cyclinD $_1$ (1:1000), Cdk $_4$ (1:1000), Bcl-2 (1:1000) and Bax (1:800) (Santa Cruz Biotechnology). Blotted antibody was developed by horseradish peroxidase-conjugated secondary antibody and enhanced chemiluminescence detection system (Amersham Pharmacia Biotech, Uppsala, Sweden).

Statistical analysis The experimental results shown were repeated twice or three times, unless otherwise indicated. Results are expressed as mean \pm SD. Statistical analysis was carried out using Student's *t*-test and one-way ANOVA. Significance was set at $P < 0.05$. Statistical analyses were performed using SPSS10.0 (SPSS Inc, Chicago, IL, USA)

Results

PPAR γ expression in human RCC tissue and RCC cell lines Using RT-PCR and Western blot analysis PPAR γ mRNA expression was detected in 15 human RCC tissues, 10 normal kidney tissues, 2 RCC cell lines (786-O and A498) and 2 normal kidney cell lines (HK-2 and HMCL). PPAR γ expression at both mRNA (Figure 1A) and protein level (Figure 1B) was detected in all tissues and cell lines examined. Much higher PPAR γ expression levels were observed in human RCC tissues and cell lines (786-O and A498) compared to normal renal tissues and cell lines (Figure 1A, 1B). Immunostaining analysis further demonstrated that PPAR γ immunoreactivity was much higher over cancer cells in human RCC tissues compared to adjacent normal tissues and normal renal tissues (Figure 1C).

Strong staining of PPAR γ was observed in 118 out of 120 human RCC tissues. PPAR γ protein was predominantly localized to nucleus and cytoplasm areas of the cells. In contrast, although 16 out of the 20 normal renal samples showed PPAR γ protein expression, only weak PPAR γ expression was observed in cytoplasm of the medullary collecting duct cells. Statistically, the intensity of PPAR γ immunoreactivity was much higher in human RCC tissues than in normal renal tissues (2.16 ± 0.85 vs 0.90 ± 0.73 ; $P < 0.01$).

PPAR γ agonists inhibited proliferation of RCC cell lines To determine the effect of PPAR γ ligands on growth of human RCC cell lines, the exponentially growing 786-O and A498 cells were treated with 0–50 $\mu\text{mol/L}$ pioglitazone and




Figure 1. (A) RT-PCR analysis showing expression of PPAR γ mRNA in human RCC tissues, normal kidney (NK) tissues, RCC cell lines 786-O and A498, and normal renal cell lines HK-2 and HMCL. β -actin was used as an internal control for RNA loading. (B) Immunoblotting analysis showing that PPAR γ protein (55 kDa) was detected in all tissues and cell lines. β -Actin (47 kDa) was used as a protein loading control. Note: higher levels of PPAR γ mRNA and protein in human RCC tissues and cell lines. (C) Immunohistochemistry analysis of PPAR γ in human RCC tissues (a and c) and normal kidneys (b and d). The slides were counterstained with hematoxylin; a and b ($\times 100$) and c and d ($\times 400$). Note much higher staining of PPAR γ in human RCC tissues than normal kidneys and intense immunoreactivity was observed over malignant cells with little staining in surrounding non-tumor cells (a&c).

troglitazone for 24 h. Cell growth was determined by an MTT assay. Figure 2 showed that two TZD PPAR γ agonists pioglitazone and troglitazone significantly inhibited the cell growth of RCC in a dose-dependent manner, with the calculated IC₅₀ values of 7.08 and 11.32 $\mu\text{mol/L}$ for pioglitazone and 5.71 and 8.38 $\mu\text{mol/L}$ for troglitazone in 786-O and A498 cells, respectively.

Effect of PPAR γ agonists on cell cycle of RCC cell lines

Cell cycle analysis was performed in two RCC cell lines after exposure to 20–30 $\mu\text{mol/L}$ pioglitazone and troglitazone for 24 h. In both cell lines, pioglitazone and troglitazone caused




Figure 2. MTT assay showing effect of two TZD PPAR γ agonists, pioglitazone and troglitazone, on the growth of human RCC cell lines. RCC 786-O cells (A) and A498 cells (B) were cultured and treated with pioglitazone and troglitazone at various concentrations. Cell growth of both 786-O and A498 cells was determined by MTT assay. Both TZD compounds inhibited cell proliferation of RCC cells in a dose-dependent manner. $n=8$. Mean \pm SD. $^{\circ}P<0.01$ versus corresponding control group.

a dose-dependent increase in the G₀/G₁ phase and decrease in the S phase (Figure 3).

Effect of PPAR γ ligands on cell cycle protein expression in human RCC cell lines The cell cycle is tightly regulated through a complex network of positive and negative regulatory molecules including cyclin dependent kinases (Cdks), cyclins, and Cdk inhibitors (Cdkis). To elucidate the role of these molecules in the inhibition of cell cycle induced by pioglitazone and troglitazone in RCC cell lines, protein extract was prepared from the cells treated with 30 $\mu\text{mol/L}$ pioglitazone or troglitazone for 0, 12, and 24 h. Immunoblotting was performed using antibodies against human PCNA, pRb, cyclin D₁, p21, p27, and Cdk4. As shown in Figure 4, pioglitazone and troglitazone treatment time-dependently caused a marked decrease in PCNA, pRb, Cyclin




Figure 3. Flow cytometry analysis demonstrating the effect of pioglitazone and troglitazone on cell cycle of two RCC cell lines, 786-O cells (A–E) and A498 cells (F–J). (A) and (F), control cells; (B) and (G), cells treated with 20 μmol/L troglitazone; (C) and (H), cells treated with 30 μmol/L troglitazone; (D) and (I), cell treated with 20 μmol/L pioglitazone; (E) and (J), cells treated with 30 μmol/L pioglitazone. The two TZD PPAR γ ligands induced the G₀/G₁ phase proportion and decreased the S phase proportion in both RCC cell lines. *n*=3. Mean±SD. ^c*P*<0.01 vs control cells.




Figure 4. Western blot analysis showing the effect of PPAR γ activation on the levels of proteins involved in cell cycle regulation in RCC cells. 786-O and A498 cells were treated with 30 μmol/L pioglitazone and troglitazone for 0, 12, and 24 h. Equal amount of cell lysate containing 50 μg protein was separated and blotted by specific antibodies as described in “Materials and Methods”. Actin was used as a sample loading control.

D₁ and Cdk4 expression and a significant increase in p21 and p27 expression.

PPAR γ agonist induced apoptosis in human RCC cell lines The ability of pioglitazone and troglitazone to induce apoptosis in two RCC cell lines was initially determined by the cytosolic histon-bound DNA fragments and cell death

ELISA assay. We found that pioglitazone and troglitazone treatment for 48 h significantly increased apoptotic cell numbers in a dose-dependent manner, with IC₅₀ values of 67.73 μmol/L and 78.12 μmol/L for pioglitazone and 65.11 μmol/L and 63.91 μmol/L for troglitazone in 786-O and A498 cells, respectively (Figure 5). When 786-O and A498 cells were exposed to either 80 μmol/L pioglitazone or 70 μmol/L troglitazone for 48 h, they exhibited typical morphological changes of apoptosis including cytoplasmic and nuclear shrinkage, chromatin condensation and fragmentation after staining with Hoechst 33342 (Figure 6). The genomic DNA of treated cells exhibited a characteristic ladder pattern after electrophoresis in agarose gel (Figure 7). To further explore the mechanisms involved in PPAR γ agonist-induced apoptosis, the expression of proteins involved in the Bcl-2 apoptotic pathway was examined in RCC cell lines. Treatment of RCC cells with pioglitazone or troglitazone for 48 h decreased the levels of Bcl-2 expression in a time-dependent manner in both 786-O and A498 cells. In contrast, the expression of Bax was increased following TZD treatment (Figure 8).

Discussion

PPAR γ is a nuclear receptor transcription factor and plays an important role in many biological processes including adipogenesis, cell growth regulation, and cell differentiation. The synthetic TZD class of compounds, including pioglitazone and troglitazone, are originally developed as therapeutic




Figure 5. Cell death ELISA assay showing apoptosis of RCC cells induced by pioglitazone and troglitazone. (A) 786-O cells treated with two PPAR γ ligands at various concentrations; (B) A498 cells treated with pioglitazone and troglitazone. Note low concentrations of the ligands had no apoptotic effect on both 786-O and A498 cells, but at concentrations exceeding 50 $\mu\text{mol/L}$, pioglitazone and troglitazone markedly induced apoptosis in the cells in a dose dependent manner. Experiments were carried out in triplicate. $n=3$. Mean \pm SD. $^{\circ}P<0.01$ versus corresponding baseline (0 $\mu\text{mol/L}$).

tic agents for diabetes mellitus and have been recently found to be specific ligands for PPAR γ with high affinity. In the past decade, evidence began to emerge suggesting TZD PPAR γ agonists not only exert antidiabetic effect in type II diabetes mellitus but also induce cell growth arrest, apoptosis and terminal differentiation in many human malignant tumors including colon cancer, breast cancer, pituitary adenomas, pancreatic carcinoma and esophageal cancer^[9-11,19,20], suggesting PPAR γ may be a potential therapeutic target for treatment of certain human cancers. The present study provides evidence that high expression of PPAR γ was observed in human RCC tissues and cell lines, and activation of PPAR γ resulted in G₀/G₁ cell cycle arrest and apoptosis involved Bcl-2 pathway.

In the present study, we observed that PPAR γ was abun-




Figure 6. Fluorescence micrographs of 786-O and A498 cells stained with Hoechst 33342 ($\times 400$). Cells were treated with 80 $\mu\text{mol/L}$ pioglitazone or 70 $\mu\text{mol/L}$ troglitazone for 48 h. (A) and (D), control cells; (B) and (E), treated with 70 $\mu\text{mol/L}$ troglitazone; (C) and (F) treated with 80 $\mu\text{mol/L}$ pioglitazone. The PPAR γ ligand-induced apoptosis was characterized by nuclear condensation and fragmentation.




Figure 7. Agarose gel pictures showing DNA fragmentation in 786-O (A) and A498 cells (B) after treatment with 70–80 $\mu\text{mol/L}$ pioglitazone or troglitazone for 48 h. Lanes M1 and M2, DNA molecular standards; lanes 1, 2, 1', and 2', untreated cells; lanes 3 and 3', after treatment with 70 $\mu\text{mol/L}$ troglitazone, respectively; lanes 4 and 4', after treatment with 80 $\mu\text{mol/L}$ pioglitazone, respectively. Note DNA ladder was observed in lanes 3, 4, 3', and 4'.

dantly expressed in almost all high-grade human RCC specimens (98.3%), and two RCC cell lines were examined. Its intracellular distribution was mainly localized in nuclear area. In contrast, although PPAR γ immunostaining was also positive in most normal renal tissues, its expression was much lower than that in RCC tissues and limited in medullary collecting duct cells in the kidney. Furthermore, its subcellular localization was predominantly in the cytoplasm. The differ-




Figure 8. Western blot analysis showing Bcl-2 and Bax expression in 786-O and A498 RCC cells. Cells were exposed to 70 $\mu\text{mol/L}$ troglitazone or 80 $\mu\text{mol/L}$ pioglitazone for 0, 12, 24, and 48 h. Cell lysate containing 50 μg protein was separated and blotted with the antibodies. Actin was used as a sample loading control.

ences of PPAR γ in quantity and intracellular distribution between human RCC and normal kidney tissues may implicate an important role of PPAR γ in tumorigenesis of renal cell carcinoma.

Consistent with a previous study^[21], activation of PPAR γ in two human RCC cell lines resulted in inhibition of RCC cell growth as assessed using MTT assay and Flow Cytometry. As seen in other tumor cells, treatment of RCC cells lines with pioglitazone and troglitazone for 24 h caused G₀/G₁ phase arrest and blocked cells from entering the S phase^[13,19,22-24]. To explore the mechanism involved in PPAR γ -induced cell growth arrest, RCC cells were treated with pioglitazone and troglitazone and expression of cell cycle proteins was examined. We observed a marked reduction in PCNA, pRb, Cdk4 and Cyclin D1 expression and a dramatic increase in p21 and p27 expression in TZD-treated RCC cells. Down-regulation of PCNA, a nuclear protein essential for DNA replication and repair by DNA polymerase δ ^[25], is in agreement to the low proliferation rate observed in these treated RCC cells. Proliferation of eukaryotic cells is tightly regulated by expression and sequential activation of cell cycle-dependent cyclins, Cdks and CdkIs^[26]. pRb, a key regulator of G₁ cell cycle progression, is phosphorylated by a set of Cyclin-Cdk complexes, such as the complexes of Cyclin D and Cdk4/6 or cyclin E and Cdk2^[27,28]. In its dephosphorylated state, pRb binds to E2F and inhibits G₁-S phase transition. In contrast, phosphorylation of pRb causes breakdown of E2F/pRb complex, and initiates DNA synthesis. In the present study, Decreased expression of phosphorylated pRb was found in these treated cells. CdkIs are cell cycle regulatory molecules having negative effects on cell cycle machinery by binding to various cyclin-Cdk complexes and inhibiting their activities. In mammals, there are two structurally defined classes of

CdkIs, the INK4 family and the KIP/CIP family^[29]. The KIP/CIP family, including p21^{CIP}, p27^{KIP1}, and p57^{KIP2}, interacts with cyclin E-Cdk2, cyclin D-Cdk4, cyclin D-Cdk6, and cyclin A-Cdk2 complexes, and inhibits their activities^[28]. The decrease of cyclin D1 and Cdk4 and increased expression of CdkI proteins including p21 and p27 strongly suggest TZD PPAR γ agonists induced the overexpression of the members of KIP/CIP family which sequentially inhibited the activities of cyclin D1-Cdk4 complex, the phosphorylation of pRb, and the G₁-S phase transition. These findings are consistent with the observation that two TZD PPAR γ agonists decreased cell proliferation in MTT assay and arrested cells in G₀/G₁ phase by flow cytometry.

Cell cycle status and cell programmed death are usually closely associated^[13]. Besides the cell cycle arrest, the inhibition of cell growth observed in RCC cells treated with pioglitazone and troglitazone may also be a result of the increase in apoptosis^[30]. Cells failing to progress to mitosis phase are destined for apoptosis. TZD have been reported to be potent agents in inducing apoptosis in many human tumors including breast cancer, pituitary tumors and gastric cancer^[10-12]. In the present study, we clearly demonstrated that at high concentrations PPAR γ agonists pioglitazone and troglitazone caused marked apoptosis in human RCC cells as assessed by morphological change, DNA fragmentation and the cell death ELISA assay. These studies suggest that TZD may exert pro-apoptotic effect on human RCC cells.

Multiple pathways are involved in apoptosis including Bcl-2 system, fas/fasL pathway, and caspase cascade^[31-33]. To explore the molecular mechanism through which TZD PPAR γ agonist induces apoptosis in human RCC cells, we examined Bcl-2 and Bax expression following pioglitazone and troglitazone treatment in 786-O cells and A498 cells. We observed for the first time that activation of PPAR γ by both TZD markedly decreased Bcl-2 protein expression but increased Bax protein expression. Because Bcl-2 protects cells from apoptosis, while the increase of Bax induces apoptosis^[34,35], our findings suggest that decreased Bcl-2 expression and increased Bax expression may participate in TZD PPAR γ agonist-induced apoptosis in human RCC cells.

PPAR γ agonists, such as 15-dPGJ₂, troglitazone, have been shown to possess effects not involving PPAR γ ^[36,37]. For instance, the potencies for inhibition of cholesterol synthesis by troglitazone is mechanistically distinct from the transcriptional regulation by PPAR γ ^[38]. In the present study, low doses of TZD fail to induce RCC cell death. It has been reported that TZD promote T cell survival at doses that induce optimal PPAR γ transcriptional activity^[39], and cell deaths induced by TZD always use concentrations of TZD several

orders of magnitude higher than K_D for PPAR γ ^[39,40]. Moreover, troglitazone could induce apoptosis of rat hepatoma cells, but rosiglitazone, a potent PPAR γ agonist, did not have the same effect^[41], indicating that PPAR γ independent pathways, such as TZD-induced loss of mitochondrial membrane potential^[42], may be involved in the mechanisms of TZD induced cell apoptosis. Furthermore, it has also been reported that PPAR γ agonists could induce apoptosis in non-malignant cells, including isolated rat mesangial cells^[43]. Thus the mechanisms of anticancer effect of TZD need to be further explored.

In summary, the present study provides evidence that PPAR γ is highly expressed in human renal cell carcinoma cells. Activation of PPAR γ by TZD PPAR γ agonists results in inhibition of cell proliferation, likely by arresting the cells in G₀/G₁ cell cycle phase. In addition, PPAR γ agonists also induce marked apoptosis partly through decreased Bcl-2 and enhanced Bax protein expression in human RCC cells. Additional research will need to be performed to explore the mechanism of TZD which induced cell apoptosis, and the potential therapeutic value of PPAR γ agonists in renal cell carcinoma.

References

- 1 Issemann I, Green S. Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 1990; 347: 645–50.
- 2 Schoonjans K, Martin G, Staels B, Auwerx J. Peroxisome proliferator-activated receptors, orphans with ligands and functions. *Curr Opin Lipidol* 1997; 8: 159–66.
- 3 Kliewer SA, Umesono K, Noonan DJ, Heyman RA, Evans RM. Convergence of 9-*cis* retinoic acid and peroxisome proliferator signaling pathways through heterodimer formation of their receptors. *Nature* 1992; 358: 771–4.
- 4 Kliewer SA, Lehmann JM, Milburn MV, Willson TM. The PPARs and PXR: nuclear xenobiotic receptors that define novel hormone signaling pathways. *Recent Prog Horm Res* 1999; 54: 345–67.
- 5 Palmer CN, Hsu MH, Griffin HJ, Johnson EF. Novel sequence determinants in peroxisome proliferator signaling. *J Biol Chem* 1995; 270: 16114–21.
- 6 Camp HS, Ren D, Leff T. Adipogenesis and fat-cell function in obesity and diabetes. *Trends Mol Med* 2002; 8: 442–7.
- 7 Rangwala SM, Lazar MA. Peroxisome proliferator-activated receptor gamma in diabetes and metabolism. *Trends Pharmacol Sci* 2004; 25: 331–6.
- 8 Fauconnet S, Lascombe I, Chabannes E, Adessi GL, Desvergne B, Wahli W, *et al*. Differential regulation of vascular endothelial growth factor expression by peroxisome proliferator-activated receptors in bladder cancer cells. *J Biol Chem* 2002; 277: 23534–43.
- 9 Sarraf P, Mueller E, Jones D, King FJ, DeAngelo DJ, Partridge JB, *et al*. Differentiation and reversal of malignant changes in colon cancer through PPARgamma. *Nat Med* 1998; 4: 1046–52.
- 10 Elstner E, Muller C, Koshizuka K, Williamson EA, Park D, Asou H, *et al*. Ligands for peroxisome proliferator-activated receptor-gamma and retinoic acid receptor inhibit growth and induce apoptosis of human breast cancer cells *in vitro* and in BXN mice. *Proc Natl Acad Sci USA* 1998; 95: 8806–11.
- 11 Heaney AP, Fernando M, Melmed S. PPAR-gamma receptor ligands: novel therapy for pituitary adenomas. *J Clin Invest* 2003; 111: 1381–8.
- 12 Takahashi N, Okumura T, Motomura W, Fujimoto Y, Kawabata I, Kohgo Y. Activation of PPARgamma inhibits cell growth and induces apoptosis in human gastric cancer cells. *FEBS Lett* 1999; 455: 135–9.
- 13 Guan YF, Zhang YH, Breyer RM, Davis L, Breyer MD. Expression of peroxisome proliferator-activated receptor gamma (PPARgamma) in human transitional bladder cancer and its role in inducing cell death. *Neoplasia* 1999; 1: 330–9.
- 14 Demetri GD, Fletcher CD, Mueller E, Sarraf P, Naujoks R, Campbell N, *et al*. Induction of solid tumor differentiation by the peroxisome proliferator-activated receptor-gamma ligand troglitazone in patients with liposarcoma. *Proc Natl Acad Sci USA* 1999; 96: 3951–6.
- 15 Mueller E, Smith M, Sarraf P, Kroll T, Aiyer A, Kaufman DS, *et al*. Effects of ligand activation of peroxisome proliferator-activated receptor gamma in human prostate cancer. *Proc Natl Acad Sci USA* 2000; 97: 10990–5.
- 16 Golimbu M, Joshi P, Sperber A, Tessler A, Al-Askari S, Morales P. Renal cell carcinoma: survival and prognostic factors. *Urology* 1986; 27: 291–301.
- 17 Flanigan RC, Campbell SC, Clark JI, Picken MM. Metastatic renal cell carcinoma. *Curr Treat Options Oncol* 2003; 4: 385–90.
- 18 Guan Y, Zhang Y, Schneider A, Davis L, Breyer RM, Breyer MD. Peroxisome proliferator-activated receptor-gamma activity is associated with renal microvasculature. *Am J Physiol Renal Physiol* 2001; 281: 1036–46.
- 19 Motomura W, Okumura T, Takahashi N, Obara T, Kohgo Y. Activation of peroxisome proliferator-activated receptor gamma by troglitazone inhibits cell growth through the increase of p27Kip1 in human pancreatic carcinoma cells. *Cancer Res* 2000; 60: 5558–64.
- 20 Hashimoto Y, Shimada Y, Itami A, Ito T, Kawamura J, Kawabe A, *et al*. Growth inhibition through activation of peroxisome proliferator-activated receptor gamma in human oesophageal squamous cell carcinoma. *Eur J Cancer* 2003; 39: 2239–46.
- 21 Inoue K, Kawahito Y, Tsubouchi Y, Kohno M, Yoshimura R, Yoshikawa T, *et al*. Expression of peroxisome proliferator-activated receptor gamma in renal cell carcinoma and growth inhibition by its agonists. *Biochem Biophys Res Commun* 2001; 287: 727–32.
- 22 Heaney AP, Fernando M, Yong WH, Melmed S. Functional PPAR-gamma receptor is a novel therapeutic target for ACTH-secreting pituitary adenomas. *Nat Med* 2002; 8: 1281–7.
- 23 Mossner R, Schulz U, Kruger U, Middel P, Schinner S, Fuzesi L, *et al*. Agonists of peroxisome proliferator-activated receptor gamma inhibit cell growth in malignant melanoma. *J Invest Dermatol* 2002; 119: 576–82.
- 24 Brockman JA, Gupta RA, Dubois RN. Activation of PPARgamma leads to inhibition of anchorage-independent growth of human

- colorectal cancer cells. *Gastroenterology* 1998; 115: 1049–55.
- 25 Luo Y, Hurwitz J, Massague J. Cell-cycle inhibition by independent CDK and PCNA binding domains in p21Cip1. *Nature* 1995; 375: 159–61.
- 26 Pines J. Four-dimensional control of the cell cycle. *Nat Cell Biol* 1999; 1: 73–9.
- 27 Bartek J, Bartkova J, Lukas J. The retinoblastoma protein pathway in cell cycle control and cancer. *Exp Cell Res* 1997; 237: 1–6.
- 28 Sherr CJ. Mammalian G1 cyclins. *Cell* 1993; 73: 1059–65.
- 29 Aprelikova O, Xiong Y, Liu ET. Both p16 and p21 families of cyclin-dependent kinase (CDK) inhibitors block the phosphorylation of cyclin-dependent kinases by the CDK-activating kinase. *J Biol Chem* 1995; 270: 18195–7.
- 30 Shi C, Yu L, Yang F, Yan J, Zeng H. A novel organoselenium compound induces cell cycle arrest and apoptosis in prostate cancer cell lines. *Biochem Biophys Res Commun* 2003; 309: 578–83.
- 31 Zhang CL, Wu LJ, Tashiro S, Onodera S, Ikejima T. Oridonin induces apoptosis of HeLa cells via altering expression of Bcl-2/Bax and activating caspase-3/ICAD pathway. *Acta Pharmacol Sin* 2004; 25: 691–8.
- 32 Filippova M, Parkhurst L, Duerksen-Hughes PJ. The human papillomavirus 16 E6 protein binds to Fas-associated death domain and protects cells from Fas-triggered apoptosis. *J Biol Chem* 2004; 279: 25729–44.
- 33 Orrenius S. Mitochondrial regulation of apoptotic cell death. *Toxicol Lett* 2004; 149: 19–23.
- 34 Li XH, Li JJ, Zhang HW, Sun P, Zhang YL, Cai SH, *et al*. Nimesulide inhibits tumor growth in mice implanted hepatoma: overexpression of Bax over Bcl-2. *Acta Pharmacol Sin* 2003; 24: 1045–50.
- 35 Salomons GS, Brady HJ, Verwijs-Janssen M, Van Den Berg JD, Hart AA, Van Den Berg H, *et al*. The Bax alpha:Bcl-2 ratio modulates the response to dexamethasone in leukaemic cells and is highly variable in childhood acute leukaemia. *Int J Cancer* 1997; 71: 959–65.
- 36 Chawla A, Barak Y, Nagy L, Liao D, Tontonoz P, Evans RM. PPAR-gamma dependent and independent effects on macrophage-gene expression in lipid metabolism and inflammation. *Nat Med* 2001; 7: 48–52.
- 37 Davies GF, Khandelwal RL, Wu L, Juurlink BH, Roesler WJ. Inhibition of phosphoenolpyruvate carboxykinase (PEPCK) gene expression by troglitazone: a peroxisome proliferator-activated receptor-gamma (PPARgamma)-independent, antioxidant-related mechanism. *Biochem Pharmacol* 2001; 62: 1071–9.
- 38 Wang M, Wise SC, Leff T, Su TZ. Troglitazone, an antidiabetic agent, inhibits cholesterol biosynthesis through a mechanism independent of peroxisome proliferator-activated receptor-gamma. *Diabetes* 1999; 48: 254–60.
- 39 Wang YL, Frauwirth KA, Rangwala SM, Lazar MA, Thompson CB. Thiazolidinedione activation of peroxisome proliferator-activated receptor gamma can enhance mitochondrial potential and promote cell survival. *J Biol Chem* 2002; 277: 31781–8.
- 40 Yu C, Chen L, Luo H, Chen J, Cheng F, Gui C, *et al*. Binding analyses between human PPARgamma-LBD and ligands. *Eur J Biochem* 2004; 271: 386–97.
- 41 Bae MA, Rhee H, Song BJ. Troglitazone but not rosiglitazone induces G1 cell cycle arrest and apoptosis in human and rat hepatoma cell lines. *Toxicol Lett* 2003; 139: 67–75.
- 42 Bova MP, Tam D, McMahon G, Mattson MN. Troglitazone induces a rapid drop of mitochondrial membrane potential in liver HepG2 cells. *Toxicol Lett* 2005; 155: 41–50.
- 43 Tsuchiya T, Shimizu H, Shimomura K, Mori M. Troglitazone inhibits isolated cell proliferation, and induces apoptosis in isolated rat mesangial cells. *Am J Nephrol* 2003; 23: 222–8.