

Full-length article

Nitric oxide inhalation inhibits inducible nitric oxide synthase but not nitrotyrosine formation and cell apoptosis in rat lungs with meconium-induced injury¹Mei-ping LU, Li-zhong DU², Wei-zhong GU, Xiang-xiang CHEN*Neonatal Intensive Care Unit, Department of Pediatrics, Children's Hospital, Zhejiang University School of Medicine, Hangzhou 310003, China***Key words**

inhaled nitric oxide; meconium; acute lung injury; nitrotyrosine; apoptosis

¹ Project supported by the National Natural Science Foundation of China (No 30371498), the Commission of Science and Technology of Zhejiang Province (No 2004C33019) and the Bureau of Education of Zhejiang Province (No 20030304), partly supported by E-Institute of Shanghai Municipal Education Commission (Project No E-04010, China)

² Correspondence to Dr Li-zhong DU.

Phn 86-138-0572-2352.

Fax 86-571-8703-3296.

E-mail dulizhong@yahoo.com.cn

Received 2005-01-10

Accepted 2005-06-03

doi: 10.1111/j.1745-7254.2005.00153.x

Abstract

Aim: To investigate the effects of inhaled nitric oxide (NO) on pulmonary inflammation, apoptosis, peroxidation and protein nitration in a rat model of acute lung injury (ALI) induced by meconium. **Methods:** Twenty-four healthy male Sprague-Dawley rats were randomly divided into 3 groups ($n=8$): meconium-induced ALI with intratracheal instillation of 1 mL/kg saline (Mec/saline group), continuous inhalation of NO at 20 μ L/L (Mec/iNO), and the control group (control). Electromicroscopic examination was used to determine the extent of epithelial apoptosis. TUNEL was used to detect DNA fragmentation in pulmonary apoptotic cells, expressed as the apoptosis index (AI). Western blotting was used to detect pulmonary inducible NO synthase (iNOS) expression. RT-PCR was used to detect interleukin (IL)-1 β mRNA expression. Cell count in bronchoalveolar lavage (BAL), myeloperoxidase (MPO) activity, as well as malondialdehyde (MDA) and nitrotyrosine formation, the markers of toxic NO-superoxide pathway in rat lung parenchyma specimens, were also examined. **Results:** Expression of iNOS protein and IL-1 β mRNA were increased significantly in the Mec/saline group (both $P<0.01$) compared with the control group. BAL cell count, MPO activity, lung injury score, pulmonary AI, MDA level and nitrotyrosine formation were also increased significantly (all $P<0.01$). The meconium-induced iNOS protein and IL-1 β mRNA expression were inhibited significantly by NO inhalation when compared with the Mec/saline group (both $P<0.01$). BAL cell count, MPO activity and lung injury score were also decreased significantly ($P<0.01$ or $P<0.05$). However, there were no statistical differences in MDA level, nitrotyrosine formation or pulmonary AI between the Mec/saline and Mec/iNO groups. Electromicroscopic examination revealed a significant degree of epithelial apoptosis in both the Mec/saline and Mec/iNO groups. **Conclusions:** Early continuous inhalation of NO 20 μ L/L may protect the lungs from inflammatory injury, but does not decrease epithelial apoptosis or lung nitrotyrosine formation. Inhalation of NO alone is not associated with a detectable increase in oxidant stress.

Introduction

Meconium aspiration syndrome (MAS) remains a significant cause of morbidity and mortality in term newborn infants with intrauterine hyperoxia, despite improvements in obstetric care. The pathophysiology of lung injury in MAS

is very complicated. The complex nature of MAS contributes to its poor response to many therapeutic interventions^[1-4]. Clinical and experimental evidence suggest that inhaled nitric oxide (NO) reduces pulmonary hypertension, improves systemic oxygenation and inhibits the transendothelial migration of activated neutrophils in a variety of patho-

logical conditions of the lungs, including meconium aspiration^[3,5,6].

As a free radical, however, NO inhalation is potentially toxic. NO induces apoptosis in various cell lines^[7]. NO inhalation can cause damage to the lung tissue through its reaction with superoxide anions, resulting in the production of peroxynitrite anion (ONOO⁻). Nitrotyrosine is a marker of ONOO⁻ production, which provides evidence for therapy-induced iNO toxicity^[8]. A previous study showed that NO inhalation during the early phase of inflammation did not increase, but rather decreased nitrotyrosine formation^[8,9]. However, nitrotyrosine formation in meconium-induced acute lung injury (ALI) with or without inhalation of NO has not been investigated. The anti-inflammatory effect of iNO in meconium-induced ALI is still controversial^[3].

The aims of our present study were to explore the mechanism of anti-inflammatory effects and to investigate the possible toxic role of iNO in meconium-induced ALI. A rat model of ALI following meconium aspiration was established. The pulmonary inducible NO synthase (iNOS) protein and interleukin (IL)-1 β mRNA expression, nitrotyrosine formation, pulmonary apoptosis, as well as lung injury score, bronchoalveolar lavage (BAL) cell count, myeloperoxidase (MPO) activity and malondialdehyde (MDA) level were measured.

Materials and methods

Meconium preparation First-passed meconium was taken from urine-free diapers of healthy term neonates. The meconium was placed in a sterile jar and frozen for less than 7 d and then lyophilized, pooled and diluted with sterile saline to a final concentration of 20% suspension^[10]. This meconium slurry was frozen at -70 °C until use.

Animal management Twenty-four 30–45 d old Sprague-Dawley rats weighing 130–170 g were studied. A rat model of meconium-induced ALI was established as previously described^[10,11]. Briefly, rats were anesthetized with sodium pentobarbital (40 mg/kg, ip). After tracheotomy, an endotracheal tube was placed through the incision, and then stabilized. Prior to meconium instillation, rats were given vecuronium bromide (2 mg/kg, ip). After 15 min, the rats were randomized into 3 groups ($n=8$ each). Sixteen of the rats were instilled intratracheally with meconium (1 mL/kg) followed by a 3 mL bolus of air to disperse the meconium into the lung to establish the model of ALI. Half of the dose was given with the rat lying on one side and the other half with the rat lying on the other side and then randomized into the Mec/saline group, with intratracheal instillation of 1 mL/kg saline, or the Mec/iNO group, with continuous inhalation of

NO at 20 μ L/L. The remaining 8 rats were given 1 mL/kg sterile saline intratracheally followed by a 3 mL bolus of air as control. At 60 min after surgery, skin and tracheal incisions were closed with a 4–0 nylon suture. The rats in the Mec/saline group and in the control group were allowed to breath spontaneously in room air.

Nitric oxide exposure Immediately after the skin and tracheal incisions were closed, the rats in the Mec/iNO group were housed for 24 h in a sealed Plexiglas chamber that was flushed continuously with a precise dilution of NO gas (800 μ L/L) and medical air (4 mL/kg) to achieve a final NO concentration of 20 μ L/L NO concentration in the chamber was continuously monitored by a NO/NO₂ monitor confirming that the NO concentration was maintained at 20 \pm 1 μ L/L and that the NO₂ concentration was consistently <2 μ L/L.

Sample collection and storage The animals were killed 24 h after treatment. The chest was opened by a midline incision. The lungs were isolated and left lung lavage was carried out. Tissue samples from the right lung were obtained. Apoptosis of lung tissue cells was evaluated using electron microscopy and a terminal deoxynucleodityl transferase-mediated dUTP nick-end labeling (TUNEL) assay was carried out.

Bronchoalveolar lavage total cell count The left lung was lavaged using 31 mL aliquots of cold sterile saline following tracheotomy. Each lung was instilled and cleared 3 times. The recovered amount was always more than 90%. A total BAL cell count was obtained within 4 h.

Pulmonary myeloperoxidase activity and malondialdehyde assay Myeloperoxidase (MPO) activity was assessed as a measure of pulmonary neutrophil influx and activity. Pulmonary MPO activity and malondialdehyde (MDA) levels were assayed in duplicate on homogenized lung samples, according to the manufacturer's manual. One unit of MPO activity was expressed as units per gram of wet lung tissue. MDA levels were standardized for tissue protein concentration determined by the Bradford protein assay^[12].

Reverse transcription-polymerase chain reaction (PCR) analysis of pulmonary IL-1 β mRNA expression Total RNA was isolated with Trizol Reagent (Sangon, Shanghai, China) in accordance with the manufacturer's instructions. Complementary DNA (cDNA) was prepared by reverse transcription (RT) of 10 μ g total RNA using oligo dT18 and 200 u MmuLV reverse transcriptase (Sangon) at 37 °C for 90 min according to the manufacturer's manual. cDNA was used as a template in the PCR, which was run in *Tag*-polymerase buffer, supplemented with 1.5 mmol/L MgCl₂, 5 μ g/mL of each primer, 4 μ L of 10 mmol/L dNTPs, and 1 U of *Tag*-polymerase (TaKaRa, Dalian, China). The total reaction volume

was 50 μ L. The primers used for IL-1 β in RT-PCR were coding, 5'-GAAGCTGTGGCA GCTACCTATGTCT-3' and uncoding, 5'-CTCTGCCTTGAGAGGTGCTGATGTAC-3', which yielded a PCR product of 520 bp. The primers for β -actin (200 bp) were coding, 5'-ATG CCA ACA CAG TGC TGT CT-3' and uncoding, 5'-CTG CTT GCT GAT CCA CAT CT-3'. Amplification was carried out for 30 cycles at 94 °C for 1 min, 65 °C for 45 s, 72 °C for 45 s, followed by a 7 min extension at 72 °C. Gene expression was determined by the size of the PCR product in 1.6% agarose gels (in Tris-buffer, pH 8.0), stained with 0.1% μ g/mL ethidium bromide and observed under ultraviolet light. The data of the IL-1 β mRNA levels was normalized against the housekeeping gene β -actin. Molecular weight markers were run in parallel.

Western blot analysis of pulmonary nitrotyrosine formation and inducible nitric oxide synthase expression Frozen lung tissues were homogenized in protein extraction buffer. The homogenate was left on ice for 30 min before centrifugation at 10 000 \times g. Protein content was estimated^[12], and aliquots of 50 μ g homogenate protein were run on 10% (w/v) sodium dodecylsulfate (SDS)-polyacrylamide gel electrophoresis (PAGE) gels. Proteins were subsequently transferred to nitrocellulose membranes and stained with Coomassie blue to verify the equivalent transfer of samples. Membranes were blocked by incubation with 3% (w/v) fat dry milk in PBS for 1 h at room temperature, with shaking. Thereafter, the membranes were incubated overnight at 4 °C with either mouse monoclonal anti-nitrotyrosine (Calbiochem-Novabiochem, San Diego, CA, USA) or rabbit polyclonal anti-iNOS or rabbit polyclonal anti-actin (Santa Cruz, California, USA). The primary antibody was used at dilutions from 1:500 to 1:1000. The blots were washed 4 times (10 min each time) with Tris-buffered saline Tween-20 (TBST), comprising 50 mmol/L Tris-HCl (pH 7.6), 150 mmol/L NaCl and 0.1% Triton X-100, and subsequently incubated for 2 h at room temperature in Tris-buffered saline with horseradish peroxidase-conjugated anti-rabbit or anti-mouse IgG antibody (1:5000; Pierce, Rockford, IL, USA). Immunoreactive bands were visualized using the enhanced chemiluminescence system (Pierce) and quantified by densitometric analysis using Scion image software (Kodak, EDAS290 Imager Analysis). Results were calculated as the relative ratio of the specific band compared with actin.

Histological examination Pulmonary tissue samples from the right lower lobe were fixed in 10% buffered formalin, embedded in paraffin, and stained with hematoxylin and eosin for determination of the severity of lung injury. Samples were assessed by a blinded pathologist who was unaware of the group of the rats. A score from 0 to 4 represented the

percentage of affected area on the lung section (0=0%, 1=1%–25%, 2=26%–50%, 3=51%–75%, and 4=76%–100%) and was assigned for 3 different characteristics: (i) extension of pulmonary leukocyte infiltration; (ii) amount of intra-alveolar leukocytes; and (iii) amount of exudative debris and edema fluid. The calculated total injury score represents the sum of these scores^[3].

Quantification of apoptosis in lung tissues by TUNEL The semiquantification of apoptotic cells in lung tissue was carried out by in situ TUNEL using the Apop Tag Peroxidase *In Situ* Cell Death Detection Kit (Roche Diagnostic GmbH Mannheim, Germany) according to the manufacturer's protocol. The number of positive signals in each section was evaluated by 3 independent pathologists using light microscopy. The apoptosis index (AI) was calculated. AI is a measure of the number of positive cells in each 100 cells counted in 5 different blocks in the same section^[13].

Electron microscopy Small pieces of lung tissue (1 mm³) from the left lower lobe were obtained for ultrastructural analysis of apoptotic pulmonary cell death by electron microscopy.

Statistical analysis Data are expressed as mean \pm SD. Statistical significance among sample groups was analyzed with ANOVA. The Mann-Whitney test was used to analyze the data from lung injury score. A level of $P<0.05$ was considered statistically significant.

Results

The meconium-instilled group showed increased BAL cell count, pulmonary MPO activity and lung injury score when compared with the control group (all $P<0.01$; Table 1). The expression of IL- β mRNA and iNOS protein was also increased significantly (both $P<0.01$; Figures 1,2). Pulmonary nitrotyrosine, a marker of ONOO⁻ formation as an index of pulmonary peroxidation, and MDA, a marker of lipid peroxidation, were increased significantly as well (both $P<0.01$; Table 1; Figure 3). Pulmonary AI, demonstrated by TUNEL staining, was significantly elevated at 24 h after the treatment ($P<0.01$; Figure 4). This was confirmed by the electron microscopic finding that apoptotic change occurred in pulmonary epithelium in the meconium-instilled rat lungs (Figure 5).

Inhalation of NO (20 μ L/L) significantly inhibited meconium-induced iNOS protein and IL-1 β mRNA expression when compared with the Mec/saline group (both $P<0.01$; Figures 1, 2). BAL cell count, MPO activity, and lung injury score were also decreased ($P<0.01$, $P<0.01$, and $P<0.05$, respectively; Table 1). However, there were no statistical

Table 1. Bronchoalveolar lavage (BAL) cell count, pulmonary myeloperoxidase (MPO) activity, malondialdehyde (MDA) and lung injury score in rats with intratracheal saline (control) or meconium instillation followed by saline instillation (Mec/saline) or nitric oxide (NO) inhalation (Mec/iNO). $n=8$. Mean \pm SD. ^c $P<0.01$ vs control; ^e $P<0.05$, ^f $P<0.01$ vs Mec/saline.

Group	10 ⁶ ×Cell count (cells/mL)	MPO (U/g wet lung tissue)	MDA (nmol/mg protein)	Lung injury score
Control	0.53 \pm 0.19	0.62 \pm 0.16	1.40 \pm 0.35	2.25 \pm 1.04
Mec/saline	4.68 \pm 1.40 ^c	1.54 \pm 0.24 ^c	3.50 \pm 0.82 ^c	10.00 \pm 1.07 ^c
Mec/iNO	3.38 \pm 0.58 ^{cf}	1.24 \pm 0.14 ^{cf}	3.25 \pm 1.20 ^c	8.50 \pm 1.14 ^{ce}




Figure 1. (A) Representative reverse transcription-polymerase chain reaction analysis of pulmonary interleukin (IL)-1 β mRNA (520 bp) from control (1, control), acute lung injury with intratracheal-instilled saline (2, Mec/saline) or with 20 μ L/L nitric oxide (NO) inhalation (3, Mec/iNO). (B) NO inhalation inhibits the expression of pulmonary IL-1 β mRNA ($n=8$ in each group), and the intensity of the bands is greater in the Mec/saline and Mec/iNO lanes compared with the control. ^c $P<0.01$ vs control. ^f $P<0.01$ vs Mec/saline.

differences in MDA level, nitrotyrosine formation or pulmonary AI between the Mec/saline and Mec/iNO groups (Table 1; Figures 3, 4). Electron microscopy revealed a significant amount of epithelial apoptosis in rat lung from both




Figure 2. (A) Representative Western blot analysis of inducible nitric oxide synthase (iNOS) (~115 kDa) from control (1, control), acute lung injury with intratracheal instilled saline (2, Mec/saline) or with 20 μ L/L nitric oxide (NO) inhalation (3, Mec/iNO). (B) The intensity of the bands is greater in the Mec/saline and Mec/iNO lanes compared with the control. NO inhalation inhibits the expression of pulmonary iNOS protein ($n=8$ in each group). ^c $P<0.01$ vs control. ^f $P<0.01$ vs Mec/saline.




Figure 3. (A) Representative Western blot analysis of nitrotyrosine (~66 kDa) formation from control (1, control), acute lung injury with intratracheal instilled saline (2, Mec/saline) or with 20 μ L/L nitric oxide (NO) inhalation (3, Mec/iNO). (B) The intensity of the bands is greater in the Mec/saline and Mec/iNO lanes compared with the control ($n=8$ in each group). ^c $P<0.01$ vs control.

the Mec/saline and the Mec/iNO group (Figure 5).

Discussion

The present study shows that meconium aspiration in meconium-induced ALI, compared with NO inhalation, is associated with increased MPO activity, lung injury score,




Figure 4. Representative lung sections from (A) control, (B) acute lung injury with intratracheal instilled saline (Mec/saline) or (C) with 20 μ L/L. nitric oxide (NO) inhalation (Mec/iNO), showing stained brown nuclei of apoptotic cells with terminal deoxynucleotidyl transferase-mediated dUTP nick-end labeling. (D) Pulmonary apoptotic cells are more numerous and the pulmonary apoptosis index is significantly increased in the Mec/saline and Mec/iNO groups compared with the control ($n=8$). ^c $P<0.01$ vs control.

high numbers of neutrophils in BAL, and increased iNOS protein and IL-1 β mRNA expression. It demonstrates that early NO inhalation at a dose of 20 μ L/L. inhibits the pulmonary inflammatory injury, but does not increase pulmonary MDA and nitrotyrosine formation in meconium induced-ALI. These findings indicate that NO inhalation may protect the




Figure 5. Representative electron microscopic figures from (A) a control lung (control), (B) a lung with acute lung injury with intratracheal instilled saline (Mec/saline) or (C) 20 μ L/L. nitric oxide (NO) inhalation (Mec/iNO). Apoptotic change occurred in type II pneumocytes in both (B) and (C). The condensed chromatin represents apoptotic change.

lung tissue against the meconium-induced pulmonary inflammatory response and that at the present dosage, the oxidative effects of iNO are not clinically significant.

Meconium contains variable amounts of proinflammatory substances, such as IL-1 β , IL-6, IL-8, TNF- α , and heme, which may activate lung inflammatory cells to produce cytokines and elicit inflammatory reactions^[11,14]. This is also demonstrated in the rat lungs in our study by induction of pulmonary IL-1 β mRNA expression. As previously reported, high expression of iNOS might lead to overproduction of NO^[15,16]. NO is an important mediator in inflammation and high levels of NO produced by iNOS can mediate lung injury^[8]. During inflammatory conditions, meconium is able to activate macrophages to produce oxygen radicals, including superoxide anions^[4]. Reaction of NO with superoxide anions produces

peroxynitrite, which is a highly oxidative species and is capable of nitrating tyrosine residues of numerous proteins, leading to the formation of nitrotyrosine that may result in protein inactivation and lipid and DNA degradation. Nitrotyrosine is a product of the peroxynitrite pathway that has been used as a probe to detect NO-mediated oxidative reactions^[8]. Our findings show that meconium aspiration results in the expression of iNOS and the production of MDA and nitrotyrosine, indicating that meconium-induced ALI is associated with pulmonary inflammatory and oxidative/nitrosative injury.

Findings from earlier studies suggest that inhaled NO can decrease inflammatory reactions in neonatal and adult lungs^[6,17]. In the present study, we showed an attenuation in pulmonary inflammatory injury after NO treatment, similar to earlier findings. Evidence from our study shows that the mechanism of the anti-inflammatory effects of NO inhalation may be associated with inhibited expression of pulmonary iNOS protein and IL-1 β mRNA. On the other hand, during inflammatory conditions, inhaled NO might increase the reaction between superoxide and NO to enhance tissue nitration and exacerbate injury. Nitrotyrosine has been found in the lungs of patients with acute respiratory distress syndrome who have received inhaled nitric oxide^[18,19]. In newborn piglets the formation of peroxynitrite may be enhanced in lungs exposed to high doses of inhaled NO or hyperoxia^[20]. However, inhaled NO might have contradictory effects on lung tissue: first, an increase in nitration as a result of increased amounts of NO and nitrite, thereby enhancing the production of peroxynitrite and the reaction via peroxidase as a source of nitrite^[19]; and second, a decrease in nitration by inhibiting leukocyte accumulation in the lungs, thereby reducing the production of NO and superoxide^[9,21]. In the present study, inhaled NO at 20 μ L/L does not further increase pulmonary nitrotyrosine formation. Some studies have also shown that early inhalation of NO in severe respiratory failure is not associated with increased NO toxicity. In the rat model of lipopolysaccharide-induced ALI, NO inhalation during the early phase of inflammation did not increase, but rather decreased tyrosine nitration and chlorination, possibly by reducing neutrophil sequestration^[9]. Inhaled NO at 100 μ L/L was shown to increase the survival of rats with hyperoxia^[22]. Inhaled NO at 20 μ L/L with concurrent hyperoxia did not increase intracellular or intercellular nitrotyrosine in a mouse model^[23]. According to evidence from this and from previous studies, the oxidative effects of iNO are not clinically significant.

There is some indirect evidence suggesting that apoptosis may play an important role in ALI. Aspirated meconium

may cause apoptotic cell death of the airway epithelium, supposedly through the action of bile salt or pulmonary inflammatory cells. Increased release of reactive oxygen and nitrogen species by pulmonary inflammatory cells may be associated with apoptotic cell death in the lung. Previous studies indicate that NO may function as an inhibitor of the apoptotic death program. An earlier study showed that NO inhalation inhibited pulmonary apoptosis in porcine meconium aspiration^[3]. However, there was a bulk of evidence indicating that NO induced apoptosis in different cell lines^[7]. In this study we found that NO inhalation did not protect the lung from meconium-induced apoptosis, despite the positive effect on pulmonary inflammatory injury.

In summary, meconium-induced ALI is associated with inflammatory oxidant injury and epithelial apoptosis. Early inhalation of NO at a dose of 20 μ L/L does not seem to prevent epithelial apoptosis caused by meconium aspiration. Inhalation of NO at the present dose is not associated with any detectable increase in oxidative or nitrosative damage. The results presented here suggest a lack of toxicity of inhaled NO due to oxidant injury and further justify therapeutic trials on inhaled NO in clinical respiratory failure. Caution with regard to pulmonary toxicity is still required.

Acknowledgement

We thank Mr Zhong-sheng YU and Mr Yao-sheng DONG for excellent technical assistance.

References

- 1 Cleary GM, Wiswell TE. Meconium-stained amniotic fluid and the meconium aspiration syndrome: an update. *Pediatr Clin North Am* 1998; 45: 511–29.
- 2 MH, Bae CW. Inhibitory effect of meconium on pulmonary surfactant function tested *in vitro* using the stable microbubble test. *Eur J Pediatr* 2000; 159: 770–4.
- 3 Holopainen R, Aho H, Laine J, Halkola L, Kaapa P. Nitric oxide inhalation inhibits apoptosis but not inflammatory injury in porcine meconium aspiration. *Acta Pediatr* 1999; 8: 1147–55.
- 4 Soukka HR, Ahotupa M, Ruutu M, Kaapa PO. Meconium stimulates neutrophil oxidative burst. *Am J Perinatol* 2002; 19: 279–84.
- 5 Miao CH, Sun B, Jiang H, Xue ZG, Lindwall R. Pharmacodynamics and pharmacokinetics of inhaled nitric oxide in dogs with septic acute respiratory distress syndrome. *Acta Pharmacol Sin* 2002; 23: 278–84.
- 6 Kinsella JP, Parker TA, Galan H, Sheridan BC, Halbower AC, Abman SH. Effects of inhaled nitric oxide on pulmonary edema and lung neutrophil accumulation in severe experimental hyaline membrane disease. *Pediatr Res* 1997; 41 Suppl: 457–63.
- 7 Brune B, von Knethen A, Sandau KB. Nitric oxide and its role in apoptosis. *Eur J Pharmacol* 1998; 351: 261–72.

- 8 Lang JD, McArdle PJ, O'Reilly PJ, Matalon S. Oxidant-antioxidant balance in acute lung injury. *Chest* 2002; 122 Suppl: S314–20.
- 9 Hataishi R, Kobayashi H, Takahashi Y, Hirano S, Zapol WM, Jones RC. Myeloperoxidase-associated tyrosine nitration after intratracheal administration of lipopolysaccharide in rats. *Anesthesiology* 2002; 97: 887–95.
- 10 Lu MP, Du LZ, Yu ZZ, Chen XX. Protective effects and mechanisms of recombinant human superoxide dismutase in acute lung injury of rats following meconium aspiration. *Chin J Pediatr* 2004; 42: 777–81.
- 11 Zagariya A, Bhat R, Uhal B, Navale S, Freidine M, Vidyasagar D. Cell death and lung cell histology in meconium aspirated newborn rabbit lung. *Eur J Pediatr* 2000; 159: 819–26.
- 12 Bradford, M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of dye-binding. *Anal Biochem* 1976; 72: 248–54.
- 13 Li HP, Li X, He GJ, Yi XH, Kaplan AP. The influence of dexamethasone on the proliferation and apoptosis of pulmonary inflammatory cells in bleomycin-induced pulmonary fibrosis in rats. *Respirology* 2004; 9: 25–32.
- 14 de Beaufort AJ, Bakker AC, van Tol MJ, Poorthuis BJ, Schrama AJ, Berger HM. Meconium is a source of pro-inflammatory substances and can induce cytokine production in cultured A549 epithelial cells. *Pediatr Res* 2003; 54: 491–5.
- 15 Ding HL, Zhu HF, Dong JW, Zhu WZ, Yang WW, Yang HT, *et al*. Inducible nitric oxide synthase contributes to intermittent hypoxia against ischemia/reperfusion injury. *Acta Pharmacol Sin* 2005; 26: 315–22.
- 16 Li YH, Yan ZQ, Braune A, Tullus K. Meconium induces expression of inducible NO synthase and activation of NF-kappaB in rat alveolar macrophages. *Pediatr Res* 2001; 49: 820–5.
- 17 Du LZ, Long XY. Effects of inhaled nitric oxide on lung injury and neutrophil accumulation in rabbits with meconium aspiration. *Acta Pharmacol Sin* 2002; 23 (Suppl): 165–6.
- 18 Lamb NJ, Quinlan GJ, Westerman ST, Gutteridge JM, Evans TW. Nitration of proteins in bronchoalveolar lavage fluid from patients with acute respiratory distress syndrome receiving inhaled nitric oxide. *Am J Respir Crit Care Med* 1999; 160: 1031–4.
- 19 Weinberger B, Laskin DL, Heck DE, Laskin JD. The toxicology of inhaled nitric oxide. *Toxicol Sci* 2001; 59: 5–16.
- 20 Youssef JA, Thibeault DW, Rezaiekhalthigh MH, Mabry SM, Norberg MI, Truog WE. Influence of inhaled nitric oxide and hyperoxia on Na, K-ATPase expression and lung edema in newborn piglets. *Biol Neonate* 1999; 75: 199–209.
- 21 Honda K, Kobayashi H, Hataishi R, Hirano S, Fukuyama N, Nakazawa H, *et al*. Inhaled nitric oxide reduces tyrosine nitration after lipopolysaccharide instillation into lungs of rats. *Am J Respir Crit Care Med* 1999; 160: 678–88.
- 22 Sato Y, Walley KR, Klut ME, English D, D'yachkova Y, Hogg JC, *et al*. Nitric oxide reduces the sequestration of polymorphonuclear leukocytes in lung by changing deformability and CD18 expression. *Am J Respir Crit Care Med* 1999; 159: 1469–76.
- 23 Lorch SA, Foust R 3rd, Gow A, Arkovitz M, Salzman AL, Szabo C, *et al*. Immunohistochemical localization of protein 3-nitrotyrosine and S-nitrosocysteine in a murine model of inhaled nitric oxide therapy. *Pediatr Res* 2000; 47: 798–805.