

## Full-length article

## Effect of resveratrol on L-type calcium current in rat ventricular myocytes

Li-ping ZHANG, Jing-xiang YIN, Zheng LIU, Yi ZHANG, Qing-shan WANG<sup>1</sup>, Juan ZHAO

Department of Physiology, Institute of Basic Medicine, Hebei Medical University, Shijiazhuang 050017, China

### Key words

resveratrol; patch-clamp techniques; myocardium; L-type calcium channels

<sup>1</sup> Correspondence to Prof Qing-shan WANG.  
Phn 86-311-606-5829.  
E-mail WQS413926@163.com

Received 2005-08-07  
Accepted 2005-09-22

doi: 10.1111/j.1745-7254.2006.00250.x

### Abstract

**Aim:** To study the effect of resveratrol on L-type calcium current ( $I_{Ca-L}$ ) in isolated rat ventricular myocytes and the mechanisms underlying these effects. **Methods:**  $I_{Ca-L}$  was examined in isolated single rat ventricular myocytes by using the whole cell patch-clamp recording technique. **Results:** Resveratrol (10–40  $\mu\text{mol/L}$ ) reduced the peak amplitude of  $I_{Ca-L}$  and shifted the current-voltage ( $I-V$ ) curve upwards in a concentration-dependent manner. Resveratrol (10, 20, 40  $\mu\text{mol/L}$ ) decreased the peak amplitude of  $I_{Ca-L}$  from  $-14.2 \pm 1.5$  pA/pF to  $-10.5 \pm 1.5$  pA/pF ( $P < 0.05$ ),  $-7.5 \pm 2.4$  pA/pF ( $P < 0.01$ ), and  $-5.2 \pm 1.2$  pA/pF ( $P < 0.01$ ), respectively. Resveratrol (40  $\mu\text{mol/L}$ ) shifted the steady-state activation curve of  $I_{Ca-L}$  to the right and changed the half-activation potential ( $V_{0.5}$ ) from  $-19.4 \pm 0.4$  mV to  $-15.4 \pm 1.9$  mV ( $P < 0.05$ ). Resveratrol at a concentration of 40  $\mu\text{mol/L}$  did not affect the steady-state inactivation curve of  $I_{Ca-L}$ , but did markedly shift the time-dependent recovery curve of  $I_{Ca-L}$  to the right, and slow down the recovery of  $I_{Ca-L}$  from inactivation. Sodium orthovanadate ( $\text{Na}_3\text{VO}_4$ ; 1 mmol/L), a potent inhibitor of tyrosine phosphatase, significantly inhibited the effects of resveratrol ( $P < 0.01$ ). **Conclusion:** Resveratrol inhibited  $I_{Ca-L}$  mainly by inhibiting the activation of L-type calcium channels and slowing down the recovery of L-type calcium channels from inactivation. This inhibitory effect of resveratrol was mediated by the inhibition of protein tyrosine kinase in rat ventricular myocytes.

### Introduction

Resveratrol (trans-3,4',5-trihydroxystilbene) is a common phytoalexin that is found in some edible materials, including grape skins, peanuts, and red wine. It has been reported to have a variety of estrogenic, anti-inflammatory, anti-platelet, and anti-carcinogenic effects<sup>[1–4]</sup>. There is accumulating evidence that resveratrol might serve as a cardioprotective agent. It is possible that resveratrol might protect the heart from ischemia-reperfusion injury, decrease plasma triglyceride and cholesterol accumulation in the aorta and prevent atherosclerosis<sup>[5,6]</sup>. Moreover, resveratrol has been found to relax the coronary arteries<sup>[7,8]</sup>. Recently, we found that resveratrol decreased the amplitude of action potential (APA), overshoot (OS) and maximal velocity of phase 0 depolarization ( $V_{\text{max}}$ ) in partially depolarized papillary muscles<sup>[9]</sup>. Moreover, resveratrol inhibited delayed afterdepolarization (DAD) and triggered activity (TA) induced by ouabain in guinea pig papillary muscles<sup>[10]</sup>. These effects were likely due to a de-

crease in calcium influx. In addition, Dobrydyneva *et al* found that resveratrol could directly block calcium channels and inhibit calcium influx in human platelets<sup>[11]</sup>. However, no study regarding the effect of resveratrol on  $I_{Ca-L}$  in ventricular myocytes has been reported. The present study was undertaken to observe the effect of resveratrol on L-type calcium channels and to investigate the mechanism by which it exerts this effect in rat ventricular myocytes using the whole-cell patch-clamp recording technique.

### Materials and methods

**Isolation of ventricular myocytes** Single ventricular myocytes were obtained by using the modified enzymatic dissociation technique<sup>[12]</sup>. Male Sprague-Dawley rats (340  $\pm$  40 g, grade II, Certificate No 04036) were provided by the Experimental Animal Center of Hebei Province. From each animal, the heart together with 2–3 mm of aorta was removed and placed in oxygenated ice-cold  $\text{Ca}^{2+}$ -free

Tyrode's solution. Langendorff retrograde perfusion was performed through the aorta at a rate of 9 mL/min with Ca<sup>2+</sup>-free Tyrode's solution for 5 min and then with the same solution containing 34 μmol/L CaCl<sub>2</sub> and 360 mg/L collagenase (type II; Sigma, Raleigh, North Carolina, USA) for 9 min at 37 °C. The ventricles were incubated in Ca<sup>2+</sup>-free Tyrode's solution at room temperature for approximately 30 min. Afterwards, a piece of ventricle was cut out and teased into smaller pieces in Kraftbrühe solution (KB solution). Myocytes were harvested after filtration through a 200-μm nylon mesh and stored in KB solution for at least 1 h before the experiment. The Ca<sup>2+</sup> concentration of the solution was gradually increased to 1.5 mmol/L. All experiments were performed within 12 h after isolation.

**Measurement of Ca<sup>2+</sup> current** Isolated ventricular myocytes were placed in an experimental chamber (0.4 mL) mounted on the stage of an inverted microscope (CK2, Olympus). After being settled to the bottom of the chamber, cells were superfused with external solution for 10 min at a rate of 2–3 mL/min at 25 °C. Transmembrane currents were recorded with an Axopatch amplifier (200 B; Axon Instruments). Glass microelectrodes were made using a microelectrode puller (PB-7; Narishige, Japan) by 2-stage pulling and had a resistance of 2.0–4.0 MΩ when filled with electrode internal solution. Only rod-shaped cells with visible cross-striations were used for experiments. Liquid junction potential between the pipette solution and external solution was corrected after the pipette tip was placed into the external solution. After gigaseal formation, the membrane was ruptured with a gentle suction to obtain the whole cell voltage-clamp configuration. To minimize the duration of capacitive current, membrane capacitance and series resistance were compensated after membrane rupture. The external solution was changed to Na<sup>+</sup>-free solution in which Na<sup>+</sup> was replaced by equimolar tetraethylammonium chloride (TEA-Cl). Na<sup>+</sup> current was also inactivated at the holding potential ( $E_h$ ) of -50 mV and blocked by tetrodotoxin (TTX; 2×10<sup>-6</sup> mol/L). K<sup>+</sup> current was suppressed by substituting intracellular K<sup>+</sup> with Cs<sup>+</sup>. Computer-generated voltage pulses were programmed using pCLAMP 6.0 software (Axon Instruments). Online acquired data were stored on the hard disk of a microcomputer. All experiments were carried out at room temperature (22–24 °C).

**Solutions and drugs** Resveratrol, collagenase type II, taurine, tetraethylammonium chloride (TEA-Cl), *N*-2-hydroxyethylpiperazine-*N'*-2-ethanesulfonic acid (HEPES), egtazic acid, CsOH, CsCl, and MgATP were purchased from Sigma. TTX was purchased from the Hebei Ocean Product Institute of China (Qinhuangdao City, Hebei Province,

China).

The Ca<sup>2+</sup>-free Tyrode's solution contained (in mmol/L) NaCl 135, KCl 5.4, MgSO<sub>4</sub> 1.0, NaH<sub>2</sub>PO<sub>4</sub> 0.33, glucose 5, HEPES 10, the pH was adjusted to 7.4 with NaOH. The KB solution contained (in mmol/L) KOH 80, KCl 40, NaH<sub>2</sub>PO<sub>4</sub> 25, MgSO<sub>4</sub> 3, glutamic acid 50, taurine 20, ethyleneglycol-*bis*(β-amino-ethyl ether)-*N,N,N',N'*-tetraacetic acid (EGTA) 1, HEPES 10, and glucose 10, the pH was adjusted to 7.4 with KOH. The electrode internal solution for whole-cell recording was composed of (in mmol/L): MgATP 3, CsCl 140, HEPES 10, egtazic acid 10, the pH was adjusted to 7.2 with CsOH. The external solution was composed of (in mmol/L): TEA-Cl 140, MgCl<sub>2</sub> 2.0, CaCl<sub>2</sub> 1.5, glucose 10, HEPES 10, TTX 0.002; and was gassed with 100% O<sub>2</sub>, and the pH was adjusted to 7.3–7.4 with TEAOH. Resveratrol was dissolved in dimethylsulfoxide and diluted in the external solution to achieve concentrations of 10, 20, and 40 μmol/L. The same amount of dimethylsulfoxide was also added to the normal external solution as a control. No change was observed during perfusion with the control solution.


**Statistical analysis** Data are expressed as mean±SD. Statistical analysis was performed using the *t*-test and one-way ANOVA. *P*<0.05 was considered statistically significant.

## Results

**Effect of resveratrol on L-type calcium current** L-type Ca<sup>2+</sup> current in rat ventricular myocytes was evoked by using a depolarizing step pulse from a holding potential ( $E_h$ ) of -50 mV to 0 mV at a frequency of 0.1 Hz. The step pulse duration was 350 ms. The rundown of  $I_{Ca-L}$  was minimized by adding Mg-ATP (3 mmol/L) and egtazic acid (10 mmol/L)<sup>[13]</sup>. Resveratrol at concentrations of 10, 20, and 40 μmol/L inhibited the peak amplitude of  $I_{Ca-L}$  in a concentration-dependent manner (Figure 1). The inhibitory effect of resveratrol disappeared after 15 min washout with control solution.

**Effect of resveratrol on current-voltage relationship of  $I_{Ca-L}$**  The current-voltage (*I*-*V*) curve of L-type Ca<sup>2+</sup> current was obtained by using a number of depolarizing step pulses (350 ms) from an  $E_h$  of -50 mV to test potentials between -40 mV and 50 mV in 10 mV increments. The pulse frequency was 0.1 Hz.  $I_{Ca-L}$  was activated at -30 mV and the peak amplitude occurred at a potential of 0 mV. Resveratrol at concentrations of 10, 20, and 40 μmol/L shifted the *I*-*V* curve upwards, and decreased the peak amplitude of  $I_{Ca-L}$  from -14.2±1.5 pA/pF to -10.5±1.5 pA/pF (*n*=6 cells from 5 hearts; *P*<0.05), -7.5±2.4 pA/pF (*P*<0.01) and -5.2±1.2 pA/pF (*P*<0.01), respectively (Figure 1).


**Effects of resveratrol on activation of  $I_{Ca-L}$**  Steady-state



**Figure 1.** Effect of resveratrol on  $I_{Ca-L}$  in isolated rat ventricular myocytes. (A) Current traces of  $I_{Ca-L}$  were recorded during 350 ms depolarization from a holding potential of -50 mV to 0 mV. (B) Effect of resveratrol on the  $I-V$  curve of  $I_{Ca-L}$  in isolated rat ventricular myocytes.  $n=6$  cells from 5 hearts. Mean $\pm$ SD.

activation of L-type calcium channels was obtained by using depolarizing pulses from an  $E_h$  of -50 mV to the test potential of +50 mV in 10 mV increments for 350 ms. The pulse frequency was 0.1 Hz. The activation curves were fitted according to the Boltzmann equation:  $I/I_{max} = 1 / \{1 + \text{EXP}[(V - V_{0.5})/\kappa]\}$ .  $V_{0.5}$  is the midpoint voltage of the activation functions, and  $\kappa$  is the Boltzmann slope parameter for activation. Resveratrol at a concentration of 40  $\mu\text{mol/L}$  shifted the half-activation potential ( $V_{0.5}$ ) from  $-19.4 \pm 0.4$  mV to  $-15.4 \pm 1.9$  mV, and the slope parameter ( $\kappa$ ) from  $5.4 \pm 0.8$  mV to  $3.8 \pm 0.6$  mV ( $n=6$  cells from 5 hearts;  $P < 0.05$ ; Figure 2).

**Effects of resveratrol on inactivation of  $I_{Ca-L}$ .** Steady-state inactivation of L-type channels was measured by




**Figure 2.** Effects of resveratrol (40  $\mu\text{mol/L}$ ) on steady-state activation kinetics (A) and steady-state inactivation kinetics (B) of  $I_{Ca-L}$  in myocytes.  $n=6$  cells from 5 hearts. Mean $\pm$ SD.

using a double-pulse protocol<sup>[14]</sup>. Membrane potential was first stepped from -50 mV to +20 mV in 10 mV increments for 1000 ms and then to +10 mV for 400 ms (test pulse) and finally clamped back to the holding potential of -50 mV at a pulse frequency of 0.1 Hz. The peak current elicited by test pulses was normalized against the maximum current and plotted against the conditioning potential. The inactivation curves were also fitted according to the Boltzmann equation.  $V_{0.5}$  is the midpoint voltage of the inactivation functions, and  $\kappa$  is the Boltzmann slope parameter for inactivation. Resveratrol at a concentration of 40  $\mu\text{mol/L}$  shifted  $V_{0.5}$  from  $-29.9 \pm 2.3$  mV to  $-31.1 \pm 3.6$  mV and  $\kappa$  from  $6.3 \pm 1.8$  mV to  $6.8 \pm 3.6$  mV ( $P > 0.05$ ;  $n=6$  cells from 5 hearts; Figure 2).

**Effect of resveratrol on recovery of  $I_{Ca-L}$  from inactivation** The recovery of  $I_{Ca-L}$  from inactivation was studied by using a double-pulse protocol consisting of a 300 ms prepulse to +10 mV (P1) followed by a 300 ms test pulse to +10 mV (P2) after a variable P1–P2 coupling interval from 0 to 2500 ms at a holding potential of -60 mV. Double-pulse stimulation was repeated every 6 s. Resveratrol at a concentration of 40  $\mu\text{mol/L}$  markedly shifted the recovery curve of  $I_{Ca-L}$  to the right and changed the half-recovery time of the  $\text{Ca}^{2+}$  channel from  $169 \pm 19$  ms to  $197 \pm 19$  ms ( $n=6$  cells from 6 hearts;  $P < 0.05$ ), indicating slower recovery of  $I_{Ca-L}$  from inactivation (Figure 3).

**Effects of sodium orthovanadate on resveratrol-induced  $I_{Ca-L}$  change** To further assess the mechanism underlying




**Figure 3.** Effects of resveratrol (40  $\mu\text{mol/L}$ ) on time-dependent recovery of  $I_{\text{Ca-L}}$  from steady-state inactivation.  $n=6$  cells from 6 hearts. Mean $\pm$ SD.

the inhibitory effect of resveratrol on  $I_{\text{Ca-L}}$ , we observed the effect of sodium orthovanadate ( $\text{Na}_3\text{VO}_4$ ; 1 mmol/L), a potent inhibitor of protein tyrosine phosphatase, on the inhibitory effect of resveratrol. When cells were pretreated with  $\text{Na}_3\text{VO}_4$  (1 mmol/L), the resveratrol-induced reduction in  $I_{\text{Ca-L}}$  was significantly attenuated. ( $n=6$  cells from 6 hearts;  $P<0.01$ ; Figure 4).

## Discussion

In the present study, we found that resveratrol (10, 20, and 40  $\mu\text{mol/L}$ ) decreased  $I_{\text{Ca-L}}$  and shifted the  $I-V$  curve upward in a concentration-dependent manner. Furthermore, the inhibitory effect of resveratrol disappeared after 15 min washout. This indicates that resveratrol can block L-type calcium channels in rat ventricular myocytes, but the characteristic of  $I-V$  curve of  $I_{\text{Ca-L}}$  was not significantly alerted by resveratrol. Resveratrol (40  $\mu\text{mol/L}$ ) shifted the steady-state activation curve of  $I_{\text{Ca-L}}$  to the right, suggesting that it can inhibit the steady-state activation of  $I_{\text{Ca-L}}$ ; but the inactivation kinetics of  $I_{\text{Ca-L}}$  were not changed by resveratrol (40  $\mu\text{mol/L}$ ). In addition, resveratrol (40  $\mu\text{mol/L}$ ) markedly shifted the recovery curve of  $I_{\text{Ca-L}}$  to the right, indicating that resveratrol can slow down the recovery of  $I_{\text{Ca-L}}$  from inactivation. These findings indicate that resveratrol decreased  $I_{\text{Ca-L}}$  in rat ventricular myocytes by markedly slowing down the activation of L-type calcium channels and their recovery from inactivation. Therefore, the myocardial electrophysiological effects of resveratrol observed in our previous studies<sup>[9,10]</sup> might be attributed to its inhibitory effect on  $I_{\text{Ca-L}}$ .



**Figure 4.** Effects of sodium orthovanadate ( $\text{Na}_3\text{VO}_4$ ; 1 mmol/L) on resveratrol-induced  $I_{\text{Ca-L}}$  change in ventricular myocytes.  $n=6$  cells from 6 hearts. Mean $\pm$ SD.

Tyrosine kinase activation is thought to contribute to cell survival, proliferation, and differentiation in many cell types<sup>[15]</sup>. Furthermore, several lines of evidence show that phosphorylation of tyrosine kinase modulates ion channel activity<sup>[16,17]</sup>. Some researchers have reported that resveratrol inhibits tyrosine kinase activity, and that many of effects of the resveratrol are mediated by tyrosine kinase<sup>[18,19]</sup>. Bruder *et al* reported that resveratrol-induced cellular phenotype was dependent on intracellular calcium and tyrosine kinase activity in bovine pulmonary artery endothelial cells<sup>[20]</sup>. Furthermore, Conte *et al* reported that resveratrol could inhibit tyrosine kinase activity in PC12 cells<sup>[21]</sup>. Sodium orthovanadate, an inhibitor of tyrosine phosphatase, can enhance protein tyrosine phosphorylation<sup>[22]</sup>. In the present study, we observed the influence of sodium orthovanadate on the effect of resveratrol. We found that pretreatment with sodium orthovanadate markedly antagonized the inhibitory effects of resveratrol on L-type calcium current, suggesting that the tyrosine kinase pathway might be involved in the effects of resveratrol.

In conclusion, we found that resveratrol inhibited  $I_{\text{Ca-L}}$  in rat ventricular myocytes mainly by inhibiting the activation of L-type calcium channels and slowing down the recovery of calcium channels from inactivation. These effects of resveratrol might be mediated via the tyrosine kinase pathway. The inhibitory effects of resveratrol on  $I_{\text{Ca-L}}$  may contribute to its antiarrhythmic actions. This study provides an electrophysiological basis for the use of resveratrol in the

treatment of cardiovascular diseases.

## References

- 1 Gehm BD, McAndrews JM, Chien PY, Jameson JL. Resveratrol, a polyphenolic compound found in grapes and wine, is an agonist for the estrogen receptor. *Proc Natl Acad Sci USA* 1997; 94: 14138–43.
- 2 Lopez-Velez M, Martinez-Martinez F, Del Valle-Ribes C. The study of phenolic compounds as natural antioxidants in wine. *Crit Rev Food Sci Nutr* 2003; 43: 233–44.
- 3 Cho DI, Koo NY, Chung WJ, Kim TS, Ryu SY, Im SY, *et al*. Effects of resveratrol-related hydroxystilbenes on the nitric oxide production in macrophage cells: structural requirements and mechanism of action. *Life Sci* 2002; 71: 2071–82.
- 4 Gerhauser C, Klimo K, Heiss E, Neumann I, Gamal-Eldeen A, Knauff J, *et al*. Mechanism-based *in vitro* screening of potential cancer chemopreventive agents. *Mutat Res* 2003; 523: 163–72.
- 5 Hattori R, Otani H, Maulik N, Das DK. Pharmacological preconditioning with resveratrol: role of nitric oxide. *Am J Physiol Heart Circ Physiol* 2002; 282: H1988–95.
- 6 Zern TL, West KL, Fernandez ML. Grape polyphenols decrease plasma triglycerides and cholesterol accumulation in the aorta of ovariectomized guinea pigs. *J Nutr* 2003; 133: 2268–72.
- 7 Jager U, Nguyen-Duong H. Relaxant effect of trans-resveratrol on isolated porcine coronary arteries. *Arzneimittelforschung* 1999; 49: 207–11.
- 8 El-Mowafy AM. Resveratrol activates membrane-bound guanylyl cyclase in coronary arterial smooth muscle: a novel signaling mechanism in support of coronary protection. *Biochem Biophys Res Commun* 2002; 291: 1218–24.
- 9 Zhao J, Ma HJ, Dong JH, Zhang LP, Liu HL, Wang QS. Electrophysiological effects of resveratrol on guinea pig papillary muscles. *Acta Physiol Sin* 2004; 56: 708–12.
- 10 Zhang LP, Ma HJ, Zhao J, Wang QS. Effects of resveratrol on delayed afterdepolarization and triggered activity induced by ouabain in guinea pig papillary muscles. *Acta Physiol Sin* 2005; 57: 361–6.
- 11 Dobrydneva Y, Williams RL, Morris GZ, Blackmore PF. Dietary phytoestrogens and their synthetic structural analogues as calcium channel blockers in human platelets. *J Cardiovasc Pharmacol* 2002; 40: 399–410.
- 12 Wang Y, Gao J, Mathias RT, Cohen IS, Sun X, Baldo GJ.  $\alpha$ -Adrenergic effects on  $\text{Na}^+\text{-K}^+$  pump current in guinea pig ventricular myocytes. *J Physiol* 1998; 509: 116–28.
- 13 Belles B, Malecot CO, Hescheler J, Trautwein W. “Rundown” of the  $\text{Ca}^{2+}$  current during long whole-cell recordings in guinea pig heart cells: role of phosphorylation and intracellular calcium. *Pflugers Arch* 1988; 411: 353–60.
- 14 Sun L, Fan JS, Clark JW, Palade PT. A model of the L-type  $\text{Ca}^{2+}$  channel in rat ventricular myocytes: ion selectivity and inactivation mechanisms. *J Physiol* 2000; 529: 139–58.
- 15 Fantl WJ, Johnson DE, Williams LT. Signalling by receptor tyrosine kinases. *Annu Rev Biochem* 1993; 62: 453–81.
- 16 Inoue R, Waniishi Y, Yamada K, Ito Y. A possible role of tyrosine kinase in the regulation of muscarinic receptor-activated cation channels. *Biochem Biophys Res Commun* 1994; 203: 1392–7.
- 17 Wijetunge S, Aalkjaer C, Schachter M, Hughes AD. Tyrosine kinase inhibitors block calcium channel currents in vascular smooth muscle cells. *Biochem Biophys Res Commun* 1992; 198: 1620–3.
- 18 Jayatilake GS, Jayasuriya H, Lee ES, Koonchanok NM, Geahlen RL, Ashendel CL, *et al*. Kinase inhibitors from *Polygonum cuspidatum*. *J Nat Prod* 1993; 56: 1805–10.
- 19 Palmieri L, Mameli M, Ronca G. Effect of resveratrol and some other natural compounds on tyrosine kinase activity and on cytolysis. *Drugs Exp Clin Res* 1999; 25: 79–85.
- 20 Bruder JL, Hsieh T, Lerea KM, Olson SC, Wu JM. Induced cytoskeletal changes in bovine pulmonary artery endothelial cells by resveratrol and the accompanying modified responses to arterial shear stress. *BMC Cell Biol* 2001; 2: 1–11.
- 21 Conte A, Pellegrini S, Tagliazucchi D. Effect of resveratrol and catechin on PC12 tyrosine kinase activities and their synergistic protection from beta-amyloid toxicity. *Drugs Exp Clin Res* 2003; 29: 243–55.
- 22 Di Salvo J, Semenchuk LA, Lauer J. Vanadate-induced contraction of smooth muscle and enhanced protein tyrosine phosphorylation. *Arch Biochem Biophys* 1993; 304: 386–91.