

## Full-length article

# Isoprenaline enhances local Ca<sup>2+</sup> release in cardiac myocytes<sup>1</sup>

Jian-xin SHEN<sup>2,3,4</sup>

<sup>2</sup>Department of Physiology, Shantou University Medical College, Shantou 515041, China; <sup>3</sup>Department of Physiology, Medical School of Xi'an Jiaotong University, Xi'an 710061, China

## Key words

local calcium release; calcium signaling; cardiac myocytes; isoprenaline; confocal microscopy

<sup>1</sup> Project supported by the National Natural Science Foundation of China (No 30470900), Natural Science Foundation of Guangdong Province (No 04020253) and SRF for ROCS, SEM.

<sup>4</sup> Correspondence to Jian-xin SHEN, PhD. Phn 86-754-890-0451 (office). E-mail jxshen@stu.edu.cn

Received 2006-04-23

Accepted 2006-05-17

doi 10.1111/j.1745-7254.2006.00383.x

## Abstract

**Aim:** Contraction of cardiac myocytes is controlled by the generation and amplification of intracellular Ca<sup>2+</sup> signals. The key step of this process is the coupling between sarcolemma L-type Ca<sup>2+</sup> channels (LCCs) and ryanodine receptors (RyRs) in the sarcoplasmic reticulum (SR).  $\beta$ -Adrenergic stimulation is an important regulatory mechanism for this coupling process. But the details underlied the global level, which require local Ca<sup>2+</sup> release study are still unclear. The present study is to explore the effects of  $\beta$ -adrenergic stimulation on local Ca<sup>2+</sup> release. **Methods:** Using confocal microscopy combined with loose-seal patch-clamp approaches, effects of isoprenaline (1  $\mu\text{mol}\cdot\text{L}^{-1}$ ), a  $\beta$ -adrenergic agonist, on local SR Ca<sup>2+</sup> release triggered by Ca<sup>2+</sup> influx through LCCs in intact rat cardiac myocytes were investigated. **Results:** Isoprenaline increased the intensity of ensemble averaged local Ca<sup>2+</sup> transients, the peak of which displayed a typical bell-shaped voltage-dependence over the membrane voltages ranging from  $\sim -40\text{mV}$  to  $\sim +35\text{mV}$ . Further analysis showed that this enhancement could be explained by the increased coupling fidelity (which refers the increased probability of RyRs activation upon depolarization), and the increased amplitude of evoked Ca<sup>2+</sup> sparks (due to more Ca<sup>2+</sup> releases through local RyRs). In addition, isoprenaline decreased the first latency, which displayed a typical "U"-shaped voltage-dependence, showing the available acceleration and synchronization of  $\beta$ -adrenergic stimulation on intracellular calcium release. **Conclusions:** Isoprenaline enhances local Ca<sup>2+</sup> release in cardiac myocytes. These results underscore the importance of regulation of  $\beta$ -adrenergic stimulation on local intermolecular signals between LCCs and RyRs in heart cells.

## Introduction

Contraction of cardiac myocytes is controlled by the generation and amplification of intracellular Ca<sup>2+</sup> signals, which are composed of the elementary Ca<sup>2+</sup> release events or Ca<sup>2+</sup> sparks, emitting from the ryanodine receptors (RyRs) in the sarcoplasmic reticulum (SR)<sup>[1,2]</sup>. This process is so called the excitation-contraction (EC) coupling<sup>[3,4]</sup>. Upon depolarization, RyRs could be activated by Ca<sup>2+</sup> influx via sarcolemma L-type Ca<sup>2+</sup> channels (LCCs). Physiologically, temporal and spatial summation of Ca<sup>2+</sup> sparks through RyRs from SR are synchronized by the trigger Ca<sup>2+</sup> influx through LCCs ( $I_{\text{Ca,L}}$ ). The mechanism for this process is termed Ca<sup>2+</sup>-

induced Ca<sup>2+</sup> release (CICR)<sup>[3,4]</sup>. Activation of LCCs and RyRs are with a specific probability that may depend on some complicated regulatory mechanisms (eg,  $\beta$ -adrenergic stimulation) or diseased states (eg, heart failure)<sup>[2,5,6]</sup>.

$\beta$ -Adrenergic stimulation is an important physiological inotropic pathway which has been well characterized in cardiac myocytes<sup>[2,4,7,8]</sup>. Generally, in heart physiology,  $\beta$ -adrenergic stimulation increases muscle contractions (inotropy) and accelerates relaxation (lusitropy). First, stimulation of  $\beta$ -adrenergic receptors by agonists activates a GTP-binding protein ( $G_s$ ), which leads to activation of adenylate cyclase (AC). And then, the level of intracellular cyclic adenylic acid (cAMP) goes up, which in turn activates protein kinase A

(PKA). PKA phosphorylates several proteins that related to excitation-contraction coupling (eg, phospholamban, LCCs, RyRs, troponin I and myosin binding protein C, and *etc*). Many studies showed that  $\beta$ -adrenergic stimulation may enhance SR  $\text{Ca}^{2+}$  release by increasing the  $I_{\text{Ca,L}}$  trigger (due to phosphorylation of LCCs), or by increasing the SR  $\text{Ca}^{2+}$  load (due to phosphorylation of phospholamban), or by enhancing the cross signaling between LCCs and RyRs (due to phosphorylation of LCCs and RyRs). However, the precise mechanisms responsible for such alterations in the intracellular local  $\text{Ca}^{2+}$  release remain incompletely understood, partly due to lack of direct investigating approaches<sup>[9]</sup>.

In the present study, a new proper method, confocal microscopic imaging combined with loose-seal patch-clamp approaches (loose-patch method)<sup>[9,10]</sup> was employed to investigate effects of isoprenaline ( $1 \mu\text{mol}\cdot\text{L}^{-1}$ ), a  $\beta$ -adrenergic agonist, on local SR  $\text{Ca}^{2+}$  release triggered by the  $\text{Ca}^{2+}$  influx through LCCs in intact rat cardiac myocytes.

## Materials and methods

**Single heart cell preparation** Enzymatically isolated ventricular myocytes from adult Sprague-Dawley rats (age, 2–3 months; weight, 225–300 g) were loaded with  $\text{Ca}^{2+}$  indicator Fluo-4-AM ( $15 \mu\text{mol}\cdot\text{L}^{-1}$ ) (Molecular Probes, Eugene, OR) for 5–8 min, followed by a 10-min rest allowing for de-esterification of the indicator, as described previously<sup>[9]</sup>. The criteria for cell selection included rod shape, clear striation and clean cell surface, and lack of spontaneous contractions during a 1-min observation period.

**Solution and isoprenaline application** Isoprenaline (ISO) was freshly made (about 2 h before use) and perfused with extracellular fluid to act on cardiac myocytes 5 min before experiments and reached a final concentration of  $1 \mu\text{mol}\cdot\text{L}^{-1}$ . The extracellular and patch pipette filling solution contained (in  $\text{mmol}\cdot\text{L}^{-1}$ ): 137 NaCl, 1  $\text{CaCl}_2$ , 4.9 KCl, 1  $\text{MgCl}_2$ , 1.2  $\text{NaH}_2\text{PO}_4$ , 15 glucose, and 20 HEPES (pH 7.4, adjusted with NaOH).

**Loose-seal patch-clamp** Cell-attached patch-clamping was established using axopatch 200B amplifier (Axon Instruments, Foster City, CA) in loose-seal configuration, as described previously<sup>[9,10]</sup>. A glass pipette (3–5 M $\Omega$ , <1  $\mu\text{m}$  at the tip) was gently pressed onto the selected cell surface to form a low resistance seal (20–40 M $\Omega$ ). The patch membrane voltage (or potential) was determined according to the equation of  $V_{\text{PM}} = \text{RP} - V_{\text{com}} \cdot \text{Rs} / (\text{Rs} + \text{Rp})$ , where  $V_{\text{PM}}$  refers to the patch membrane voltage, RP the resting potentials (approximately -80 mV),  $V_{\text{com}}$  the command voltage applied, Rs and Rp the seal resistance and pipette resistance, respectively.


**Confocal  $\text{Ca}^{2+}$  imaging**  $\text{Ca}^{2+}$  images were acquired by

using a Zeiss LSM510 confocal microscope equipped with an argon laser (488 nm) and a 40 $\times$ , 1.3 numerical aperture, oil-immersion objective, at sampling rates of 0.77 ms per line and 45 nm per pixel. Using loose-patch method, population of in-focus  $\text{Ca}^{2+}$  sparks could be evoked by repeated patch depolarization, with an interval of 6 s between two consecutive confocal microscopic images in line-scan mode. All experiments were performed at room temperature (23–25  $^{\circ}\text{C}$ ).

**Data analysis and statistics**  $\text{Ca}^{2+}$  spark detection algorithm was almost the same as that described previously<sup>[11]</sup>, with some minor modifications. Computer programs for the spark detection and measurement were coded in Interactive Data Language (IDL, Research Systems, Boulder, CO).  $\text{Ca}^{2+}$  spark amplitudes were measured as  $\Delta R = \Delta F / F_0$ , where  $F$  refers to the present Fluo signal intensity,  $F_0$  the background Fluo signal intensity, and  $\Delta F / F_0$  the alteration of  $F / F_0$ . Data were expressed as mean  $\pm$  SEM, if not otherwise specified. Ensemble averaged single-couplon  $\text{Ca}^{2+}$  transients images were guided by the onsets of the depolarization pulse (Figure 1), while averaged line-scan images of  $\text{Ca}^{2+}$  sparks by the peak position of each spark (Figure 4). In this study, single-couplon refers to one elementary  $\text{Ca}^{2+}$  release unit, including one or few LCCs and some coupled RyRs. The significance of difference between means or ratios was determined, when appropriate, by using the Student  $t$  test or the nonparametric Kruskal-Wallis test.  $P < 0.05$  was considered statistically significant.

## Results

**Effect of isoprenaline on local intracellular  $\text{Ca}^{2+}$  release revealed by ensemble averaged single-couplon  $\text{Ca}^{2+}$  transients** To have a overall look at the effect of isoprenaline (ISO,  $1 \mu\text{mol}\cdot\text{L}^{-1}$ ) on local intracellular  $\text{Ca}^{2+}$  release in cardiac myocytes, depolarization pulse with a duration of 100 ms were applied under loose-seal patch-clamp configuration and confocal line-scan  $\text{Ca}^{2+}$  images were obtained from 62 patches (27 patches under the control and 35 patches with ISO treatment). Pixel-to-pixel averaging of confocal line-scan images guided by the onsets of the depolarization pulse was performed on all runs grouped by membrane potentials, with an increment of 15 mV (Figure 1A and Figure 1B). These are called the images of ensemble averaged single-couplon  $\text{Ca}^{2+}$  transients (Figure 1A left and Figure 1B left). At all membrane potential levels (ranging from approximately approximately -40 mV to approximately +35 mV), ensemble averaged single-couplon  $\text{Ca}^{2+}$  transients triggered by the depolarization pulse after ISO treatment were much stronger than that triggered under control condition (Figure 1A and Figure 1B) and displayed a typical bell-shaped voltage-dependence




**Figure 1.** Effect of isoprenaline on ensemble averaged single-couplon  $Ca^{2+}$  transients. (A) Results in control condition. Left, Spatiotemporal visualization of ensemble averaged single-couplon  $Ca^{2+}$  transients at six representative voltages. Pixel-to-pixel averaging of confocal images was performed on all runs grouped in 15-mV increments. Total runs for each group are from 29 to 44. Right, The line plots show time courses of the averaged local evoked  $Ca^{2+}$  transients. RP, resting potential. Sym  $\Delta F/F_0$ , refers to the alteration of Fluo-4 signal ratio to the background Fluo-4 signal. (B) Results after isoprenaline ( $1 \mu\text{mol}\cdot\text{L}^{-1}$ ). Total runs for each group are from 43 to 58. (C) Effect of isoprenaline on the peak amplitudes of ensemble averaged single-couplon  $Ca^{2+}$  transients. Con, peak of the averaged  $Ca^{2+}$  transients under the control condition. ISO, peak of the averaged  $Ca^{2+}$  transients under the isoprenaline condition.  $^bP < 0.05$  vs control.

(Figure 1C). These results showed that generally ISO enhanced the local intracellular  $Ca^{2+}$  release ( $P < 0.05$ ).

**Effect of ISO on coupling fidelity** Generally, enhancement of ensemble averaged single-couplon  $Ca^{2+}$  transients could be the result of the increased coupling fidelity and/or the increased amplitude of local  $Ca^{2+}$  release, both of which could be tested in the present study. Coupling fidelity of local  $Ca^{2+}$  release (including  $Ca^{2+}$  sparks) was analyzed first. In this study, coupling fidelity ( $\delta$ ) was determined by the number of active runs over the corresponding total number of all runs (including active runs and silent runs). Active runs refers to those images that recorded the triggered single-couplon  $Ca^{2+}$  release (including  $Ca^{2+}$  sparks). The peak of  $\Delta F/F_0$  for an active run is at least 0.25 (Figure 4A). For instance, under the control condition, when  $V_{PM}$  = approximately 5 mV, total runs of 29 were acquired and 27 of them were active runs. So the coupling fidelity is 0.93, meaning that the probability for the  $Ca^{2+}$  influx through LCCs to successfully trigger RyRs activation is 93%. Results showed

that at all membrane potential levels (ranging from  $\sim -40$  mV to  $\sim +35$  mV), coupling fidelity increased after ISO treatment (Figure 2) and displayed a left half bell-shaped voltage-dependence. These results explain one possible mechanism for enhancement of local intracellular  $Ca^{2+}$  release by ISO: to trigger more elementary  $Ca^{2+}$  release events. In other words, local  $Ca^{2+}$  release is easier to be activated with  $\beta$ -adrenergic stimulation.


**Effect of ISO on averaged  $Ca^{2+}$  spark amplitude** Next, effect of ISO on averaged  $Ca^{2+}$  spark amplitude was considered. Since the solitary  $Ca^{2+}$  sparks were difficult to separate from each other when the plasmic membrane was depolarized to higher voltage level (eg  $> 10$  mV), only those solitary  $Ca^{2+}$  sparks that were triggered around -30 mV or 0 mV were taken into account. Pixel-to-pixel averaging of confocal line-scan images of solitary  $Ca^{2+}$  sparks were guided by the peak position of each spark (Figure 3). Results showed that the evoked averaged  $Ca^{2+}$  sparks were stronger in ISO treatment condition than in control condition, at either ap-



**Figure 2.** Effect of isoprenaline on single-couplon excitation-contraction coupling fidelity ( $\delta_{sc}$ ).  $\delta_{sc}$  is shown as a function of patch membrane voltage ( $V_{PM}$ ).  $\delta_{sc}$  were determined by the numbers of active runs over the total numbers of runs at different voltage group in 15-mV increments. <sup>b</sup> $P < 0.05$  vs control.


proximately -30 mV or approximately 0 mV (Figure 3). That is,  $\beta$ -adrenergic stimulation enhances the intensity of single elementary  $Ca^{2+}$  release event.

**Effect of ISO on the first latency of couplon activation ( $L_{sc}$ )** In addition, the first latency of couplon activation was investigated, too. The first latency of couplon activation refers to the duration from the onset of the depolarization pulse to the onset of the first  $Ca^{2+}$  spark that is triggered



**Figure 3.** Effect of isoprenaline on averaged sparks at different voltages. (A) Averaged spark at  $\sim -30$  mV. From top to bottom: averaged line-scan images of sparks at peak, time courses of the corresponding line-scan images (spatially averaged over 1.1- $\mu$ m width); left: averaged spark under the control condition; right: averaged spark after isoprenaline ( $1 \mu\text{mol}\cdot\text{L}^{-1}$ ) treatment. (B) Averaged spark at  $\sim 0$  mV. <sup>b</sup> $P < 0.05$  vs control.

by this depolarization pulse (Figure 4A). The results showed that the first latency were shorter in ISO treatment condition than in control condition ( $P < 0.05$ ) (Figure 4B) and displayed



**Figure 4.** Effect of isoprenaline on the first latency of couplon activation ( $L_{sc}$ ). (A) A typical case to show the first latency measurement. From top to bottom: the voltage protocol, line-scan image of the evoked spark, its contour plot at  $\Delta F/F_0 = 0.25$  and its mass center (marked by the cross), and time course of local  $Ca^{2+}$  transient (spatially averaged over 1.1- $\mu$ m width). The mass center coordinates ( $X, T$ ) were determined by  $X = (\iint x \Delta F dx dt) / (\iint \Delta F dx dt)$  and  $T = (\iint t \Delta F dx dt) / (\iint \Delta F dx dt)$ . The mass center coordinates were used to determine whether two or more  $Ca^{2+}$  sparks in one image were activated from the same single couplon (see Ref 9). (B)  $L_{sc}$  is shown as a function of patch membrane voltage ( $V_{PM}$ ). <sup>b</sup> $P < 0.05$  vs control.

a typical “U”-shaped voltage-dependence over membrane potentials ranging from approximately -40 mV to approximately +35 mV. This suggests that after ISO treatment (ie  $\beta$ -adrenergic stimulation), single  $\text{Ca}^{2+}$  release unit (or single-couplon) is more sensitive to be activated.

## Discussion

Physiologically sympathetic nerve regulates heart muscles through  $\beta$ -adrenergic receptors to generate positive inotropy (increasing contractility), positive chronotropy (increasing heart rate), positive dromotropy (increasing conduction velocity) and lusitropy (relaxation acceleration)<sup>[12,13]</sup>. Inotropy and lusitropy are tightly related to the process of CICR. Inotropic effect is mediated by the combination of increased  $\text{Ca}^{2+}$  influx and greater availability of SR  $\text{Ca}^{2+}$ <sup>[2,4]</sup>. And lusitropic effect is mediated by phosphorylation of phospholamban and troponin I, which speed up SR  $\text{Ca}^{2+}$  reuptake and dissociation of  $\text{Ca}^{2+}$  from the myofilaments<sup>[2,4]</sup>. When CICR is activated in a synchronized way on a global cell level, the  $\text{Ca}^{2+}$  signals are predominantly governed by SR  $\text{Ca}^{2+}$  content, which defines the amplitude of the resulting  $\text{Ca}^{2+}$  transients<sup>[2,14]</sup>. However, when CICR is activated on a local level, the  $\text{Ca}^{2+}$  signals behave in a dramatically different way, which cannot be explained only by global changes in SR  $\text{Ca}^{2+}$  content<sup>[2]</sup>. Obviously, local  $\text{Ca}^{2+}$  release depends somewhat on the SR  $\text{Ca}^{2+}$  load. But since local SR  $\text{Ca}^{2+}$  release only occurs from one or a few functional units of the SR, local SR  $\text{Ca}^{2+}$  content depletion by local  $\text{Ca}^{2+}$  release can be refilled instantly from neighbouring SR network<sup>[2,15]</sup>. Enhancement of global  $\text{Ca}^{2+}$  release by ISO has been well addressed in some previous studies<sup>[2,4,7]</sup>. However, details of local  $\text{Ca}^{2+}$  release are still unclear. In the present study, confocal microscopic imaging combined with loose-seal patch-clamp approaches is a proper way to explore local intracellular  $\text{Ca}^{2+}$  release directly<sup>[9,10]</sup>.

**Enhancement of ensemble averaged single-couplon  $\text{Ca}^{2+}$  transients by ISO is consistent with previous works at global level** As stated in the results, ensemble averaged single-couplon  $\text{Ca}^{2+}$  transients were increased by ISO and their peak amplitudes displayed a typical bell-shaped voltage-dependence, just like the relationship between L-type  $\text{Ca}^{2+}$  currents ( $I_{\text{Ca,L}}$ ) and membrane voltages<sup>[7]</sup>. This confirmed the local  $\text{Ca}^{2+}$  signals were triggered by  $\text{Ca}^{2+}$  influx through L-type  $\text{Ca}^{2+}$  channels. These results were consistent with the observations that  $\beta$ -adrenergic stimulation can greatly enhance  $\text{Ca}^{2+}$  transients amplitude on global level<sup>[2,4,7]</sup>.

**Enhancement of local  $\text{Ca}^{2+}$  release by ISO could be explained by the increased coupling fidelity and the increased amplitude of evoked  $\text{Ca}^{2+}$  sparks** Results showed that both

coupling fidelity of single-couplon (or single  $\text{Ca}^{2+}$  release unit) and evoked  $\text{Ca}^{2+}$  spark amplitude were increased significantly by ISO treatment. This suggests that  $\beta$ -adrenergic stimulation makes more single-couplons of  $\text{Ca}^{2+}$  release ready to be activated; and in each activation, more  $\text{Ca}^{2+}$  releases through SR  $\text{Ca}^{2+}$  release channels (or RyRs) from SR. The latter may be due to increased RyRs open probability (one result of phosphorylation of RyRs by  $\beta$ -adrenergic stimulation)<sup>[16-18]</sup> and increased SR  $\text{Ca}^{2+}$  load (the result of phosphorylation of phospholamban, a relief to inhibit the SR  $\text{Ca}^{2+}$  pump uptake)<sup>[2,4,19]</sup>.

**Significance of the shortened first latency of local  $\text{Ca}^{2+}$  release by ISO** The present results also showed that ISO could shorten the first latency of local  $\text{Ca}^{2+}$  release and displayed a typical “U”-shaped voltage-dependence. First latency of local  $\text{Ca}^{2+}$  release indicates the shortest time to activate an elementary  $\text{Ca}^{2+}$  release. The results suggest that  $\beta$ -adrenergic stimulation accelerates local  $\text{Ca}^{2+}$  release activation and thus synchronizes intracellular  $\text{Ca}^{2+}$  release on global level<sup>[7]</sup>. This may be due to the bigger triggered  $\text{Ca}^{2+}$  influx (the result of phosphorylation of LCCs) and increased sensitivity of  $\text{Ca}^{2+}$  release unit to  $\text{Ca}^{2+}$  (the result of phosphorylation of RyRs and phospholamban)<sup>[2,4,7,20]</sup>.

In conclusion, using confocal microscopic imaging combined with loose-seal patch-clamp approaches, effects of isoprenaline (ISO, 1  $\mu\text{mol}\cdot\text{L}^{-1}$ ) on local SR  $\text{Ca}^{2+}$  release triggered by  $\text{Ca}^{2+}$  influx through LCCs in intact rat cardiac myocytes were investigated in the present study. ISO, acting as a  $\beta$ -adrenergic agonist, increased the intensity of ensemble averaged local  $\text{Ca}^{2+}$  transients and displayed a typical bell-shaped voltage-dependence. This enhancement could be explained by two aspects of data: the increased coupling fidelity (which refers the probability of RyRs activation upon depolarization), and the increased amplitude of evoked  $\text{Ca}^{2+}$  spark (due to more  $\text{Ca}^{2+}$  releases through local RyR activation from SR). In addition, ISO could decrease the first latency of local evoked  $\text{Ca}^{2+}$  signals and displayed a typical “U”-shaped voltage-dependence. All these data were consistent with some previous studies on global levels. Furthermore, confocal microscopic imaging combined with loose-seal patch-clamp approaches reveals more subcellular details, especially the  $\text{Ca}^{2+}$  spark amplitude and the first latency of local  $\text{Ca}^{2+}$  release, which could not be unraveled by other methods previously. All these results underscore the importance of regulation (eg phosphorylation) of  $\beta$ -adrenergic stimulation on LCCs and RyRs or some other related proteins to enhance local  $\text{Ca}^{2+}$  signals<sup>[2,4]</sup>. This may also provide new therapeutic avenues to recover impaired  $\text{Ca}^{2+}$

signaling during cardiac disease<sup>[5,6,12]</sup>.

## Acknowledgement

I would like to thank the valuable comments on the manuscript by Dr He-ping CHENG (the Institute of Molecular Medicine, Peking University, Beijing 100871, China) and Dr Tai-zhen HAN (Department of Physiology, Medical School of Xi'an Jiaotong University, Xi'an 710061, China); and the strong technology support and nice help from Dr Shi-qiang WANG (National Laboratory of Biomembrane and Membrane Biotechnology, Peking University, Beijing 100871, China).

## References

- Cheng H, Lederer WJ, Cannell MB. Calcium sparks: elementary events underlying excitation-contraction coupling in heart muscle. *Science* 1993; 262: 740–4.
- Lindegger N, Niggli E. Paradoxical SR Ca<sup>2+</sup> release in guinea-pig cardiac myocytes after beta-adrenergic stimulation revealed by two-photon photolysis of caged Ca<sup>2+</sup>. *J Physiol* 2005; 565: 801–13.
- Cheng H, Wang SQ. Calcium signaling between sarcolemmal calcium channels and ryanodine receptors in heart cells. *Front Biosci* 2002; 7: d1867–78.
- Bers DM. Cardiac excitation-contraction coupling. *Nature* 2002; 415: 198–205.
- Wang H, Oestreich EA, Maekawa N, Bullard TA, Vikstrom KL, Dirksen RT, *et al.* Phospholipase C epsilon modulates beta-adrenergic receptor-dependent cardiac contraction and inhibits cardiac hypertrophy. *Circ Res* 2005; 97: 1305–13.
- Ziolo MT, Maier LS, Piacentino V 3rd, Bossuyt J, Houser SR, Bers DM. Myocyte nitric oxide synthase 2 contributes to blunted beta-adrenergic response in failing human hearts by decreasing Ca<sup>2+</sup> transients. *Circulation* 2004; 109: 1886–91.
- Song LS, Wang SQ, Xiao RP, Spurgeon H, Lakatta EG, Cheng H.  $\beta$ -Adrenergic stimulation synchronizes intracellular Ca<sup>2+</sup> release during excitation-contraction coupling in cardiac myocytes. *Circ Res* 2001; 88: 794–801.
- Katra RP, Laurita KR. Cellular mechanism of calcium-mediated triggered activity in the heart. *Circ Res* 2005; 96: 535–42.
- Shen JX, Wang S, Song LS, Han T, Cheng H. Polymorphism of Ca<sup>2+</sup> sparks evoked from in-focus Ca<sup>2+</sup> release units in cardiac myocytes. *Biophys J* 2004; 86: 182–90.
- Wang SQ, Song LS, Lakatta EG, Cheng H. Ca<sup>2+</sup> signalling between single L-type Ca<sup>2+</sup> channels and ryanodine receptors in heart cells. *Nature* 2001; 410: 592–6.
- Cheng H, Song LS, Shirokova N, Gonzalez A, Lakatta EG, Rios E, *et al.* Amplitude distribution of calcium sparks in confocal images: theory and studies with an automatic detection method. *Biophys J* 1999; 76: 606–17.
- Bers DM, editors. Cardiac inotropy and Ca mismanagement, in excitation-contraction coupling and cardiac contractile force. Boston: Kluwer Academic Publishers; 2001. p273–331.
- Ziolo MT, Katoh H, Bers DM. Positive and negative effects of nitric oxide on Ca<sup>2+</sup> sparks: influence of beta -adrenergic stimulation. *Am J Physiol Heart Circ Physiol* 2001; 281: H2295–303.
- Sah R, Ramirez RJ, Backx PH. Modulation of Ca<sup>2+</sup> release in cardiac myocytes by changes in repolarization rate: role of phase-1 action potential repolarization in excitation-contraction coupling. *Circ Res* 2002; 90: 165–73.
- Brochet DXP, Yang D, Maio AD, Lederer WJ, Franzini-Armstrong C, Cheng H. Ca<sup>2+</sup> blinks: rapid nanoscopic store calcium signaling. *Proc Natl Acad Sci USA* 2005; 102: 3099–104.
- Valdivia HH, Kaplan JH, Ellis-Davies GC, Lederer WJ. Rapid adaptation of cardiac ryanodine receptors: modulation by Mg<sup>2+</sup> and phosphorylation. *Science* 1995; 267: 1997–2000.
- Marx SO, Gaburjakova J, Gaburjakova M, Henrikson C, Ondrias K, Marks AR. Coupled gating between cardiac calcium release channels (ryanodine receptors). *Circ Res* 2001; 88: 1151–8.
- Lu X, Xu L, Meissner G. Phosphorylation of dihydropyridine receptor II-III loop peptide regulates skeletal muscle calcium release channel function. *J Biol Chem* 1995; 270: 18459–64.
- Ginsburg KS, Bers DM. Modulation of excitation-contraction coupling by isoproterenol in cardiomyocytes with controlled SR Ca<sup>2+</sup> load and Ca<sup>2+</sup> current trigger. *J Physiol (Lond)* 2004; 556: 463–80.
- Shannon TR, Chu G, Kranias EG, Bers DM. Phospholamban decreases the energetic efficiency of the sarcoplasmic reticulum Ca pump. *J Biol Chem* 2001; 276: 7195–201.