

Full-length article

Metabolism and effect of para-toluene-sulfonamide on rat liver microsomal cytochrome P450 from *in vivo* and *in vitro* studiesJiang-quan ZHOU^{1,3}, Zhi-qiang TANG¹, Jin-nan ZHANG², Jing-cheng TANG²¹Department of Pharmacy, Cancer Hospital, Chinese Academy of Medical Science and Peking Union Medical College, Beijing 100021, China; ²School of Chemical Biology and Pharmaceutical Sciences, Capital University of Medical Sciences, Beijing 100069, China**Key words**

cytochrome P450; para-toluene-sulfonamide; liver microsomes; HPLC; inhibitor

³ Correspondence to Jiang-quan ZHOU.
Phn 86-10-8778-8428.
E-mail zhoujiangquan@gmail.comReceived 2005-10-08
Accepted 2005-12-04

doi: 10.1111/j.1745-7254.2006.00307.x

Abstract

Aim: To study the *in vivo* and *in vitro* metabolism and the effect of para-toluene-sulfonamide (PTS) on cytochrome P450 enzymes (CYP450). **Methods:** Total CYP450 and microsome protein content were determined after iv pretreatment of rats with PTS. CYP-specific substrates were incubated with rat liver microsomes. Specific CYP isoform activities were determined by using HPLC. CYP chemical inhibitors added to the incubation mixture were used to investigate the principal CYP isoforms involved in PTS metabolism. The effect of PTS on CYP isoforms was investigated by incubating PTS with specific substrates. **Results:** The groups treated with 33 and 99 mg/kg per d PTS, respectively, had a total CYP content of 0.66 ± 0.17 and 0.60 ± 0.12 nmol/mg. The K_m and V_{max} were $92.2 \mu\text{mol/L}$ and 0.0137 nmol/min per mg protein. CYP2C7, CYP2D1 and CYP3A2 might contribute to PTS metabolism in the rat liver. The inhibitory effects of sulfaphenazole and ketoconazole on PTS metabolism were shown to have a mixed mechanism, whereas PTS metabolism was inhibited noncompetitively by quinidine. PTS had little effect on the activities of the selected CYP isoforms. **Conclusion:** Generally speaking, it is relatively safe for PTS to be co-administered with other drugs. However, care should be taken when administering PTS with CYP inhibitors and the substrates of CYP2C, CYP2D and CYP3A.

Introduction

Para-toluene-sulfonamide (*p*-methylbenzenesulfonamide; PTS) is a novel anticancer reagent for which phase III clinical trials are being conducted in China. PTS is intended mainly for the treatment of breast cancer and head and neck squamous cancer. It is delivered by intravenous or intratumoral injection as an adjunct to chemotherapy and radiation therapy. The reagent has good lipophilic ability, and its clinical pharmacokinetics accord with the one-compartment model.

Because cytochrome P450 enzyme play a prominent role in the metabolism of many pharmaceutical agents and activation or deactivation of potential carcinogens, it would be useful to know as early in the development process as possible that which CYP enzymes are likely to process a new chemical entity (NCE), which CYP activities are likely to be

altered by an NCE and the magnitude of the alteration. Moreover, CYP genes have promise in enhancing the sensitivity of tumor cells to cancer chemotherapeutic drugs^[1]. Inhibition screening is very useful not only for investigating the potential influence of new drugs on liver CYP, but also for future applications of pharmacogenomics and pharmacogenetics to personalize treatment regimens^[2].

In our experiments, we chose five CYP isoforms and their specific substrates (phenacetin for CYP1A2, tolbutamide for CYP2C7, dextromethorphan for CYP2D1 and CYP3A2, and chlorzoxazone for CYP2E1). Dextromethorphan is reduced to dextrorphan by the rat CYP2D1 enzyme. In a parallel pathway, it is *N*-demethylated to 3-methoxymorphinan, which is catalyzed by CYP3A2^[3]. The known inhibitors selected for CYP1A2, CYP2C7, CYP2D1 and CYP3A2 were, respectively, α -naphthoflavone, sulfaphenazole, quinidine, and ketoconazole. The present study was designed to: (1)

investigate the effect of PTS pretreatment on rat liver CYP content; (2) examine the kinetic parameters of PTS incubation metabolism toward CYP; (3) to identify the principal CYP isoforms that metabolize PTS; (4) assess the inhibitory mechanism of known inhibitors on PTS metabolism *in vitro*; (5) investigate the modulation effect of PTS on the activities of selected CYP isoforms.

Materials and methods

Chemicals PTS standard was manufactured by Aldrich (Wyoming, IL, USA), phenacetin (Phe), acetaminophen (Ace), tolbutamide (Tol), 4-hydroxytolbutamide (4-OH-Tol), chlorzoxazone (Chl), 6-hydroxychlorzoxazone (6-OH-Chl), dextromethorphan hydrobromide monohydrate (Dex), dextrophan (Dor), 3-methoxymorphinan (3-MM), α -naphthoflavone (α -Naph), sulfaphenazole (Sulf), quinidine (Qui), ketoconazole (Ket), antipyrine, β -nicotinamide adenine dinucleotide phosphate (β -NADP), glucose-6-phosphate (G-6-P), and glucose-6-phosphate dehydrogenase (G-6-PDH) were obtained from Sigma-Aldrich (St Louis, MO, USA). Chromatographic grade acetonitrile was purchased from Fisher Scientific (Fair Lawn, NJ, USA). All other supplies were of the highest grades available from standard commercial sources.

Solutions Stock solutions of the analytes were prepared separately by dissolving each compound in water with acetonitrile at concentrations lower than 1% (v/v)^[4]. G-6-PDH was dissolved in 5 mmol/L sodium citrate and was kept at $-80\text{ }^{\circ}\text{C}$ until used.

Microsome preparation Male Wistar rats (3 months old, weighing 230–260 g) were purchased from the Department of Laboratory Animal Sciences, Capital University of Medical Sciences. Animals were divided into 4 groups, with 6 animals in every group. The groups received the following treatments: blank control, phenobarbital iv injection (40 mg/kg per d)^[5], or PTS intravenous injection via the tail vein at dosages of 33 and 99 mg/kg per d.

Rats were killed after the animals had been treated for 4 consecutive days. Microsomes were prepared by differential centrifugation and the preparation was stored at $-80\text{ }^{\circ}\text{C}$ until used. All procedures were performed at $0\text{--}4\text{ }^{\circ}\text{C}$ ^[6]. Total CYP content and microsome protein were measured by using the method of Omura^[7] and Lowry^[8]. SPSS pharmaceutical software was used for statistical comparisons. The significance level was set at $P < 0.05$.

Microsome incubation and sample preparation The incubation volume was 0.5 mL, containing 1.0 mg protein, 100 mmol/L potassium phosphate buffer (pH 7.4), and an

NADPH-generating system (0.5 mmol/L β -NADP, 2.0 mmol/L G-6-P, 5 mmol/L MgCl_2 , and 0.1 mmol/L ethylenediamine tetraacetic acid). The samples were preincubated for 5 min at $37\text{ }^{\circ}\text{C}$ prior to the addition of 2 U G-6-PDH. Reactions were carried out for 30 min and terminated by placement into icy water and the addition of 100 μL 7% perchloric acid. The samples were centrifuged at $16\ 100\times g$ for 10 min to pellet the protein precipitate. The supernatant was transferred for HPLC analysis. All incubations were carried out in triplicate, and the mean values were used for analysis^[9].

Apparatus and chromatographic conditions^[10] The samples were analyzed on an Agilent 1100 series liquid chromatograph (Agilent Technologies, Waldbronn, Germany) at room temperature. An aliquot (50 μL) from each sample was injected onto an Agilent XDB C_{18} column (4.6 mm \times 250 mm, 5 μm). Compounds were quantified using their peak areas.

For CYP1A2 phenacetin *O*-deethylase, the mobile phase consisted of a gradient of acetonitrile/water with 0.1% acetic acid (pH 3.5) monitored at 230 nm. The flow rate was 1.2 mL/min. The retention times of Ace and Phe were 4.7 min and 9.9 min, respectively.

For CYP2C7 tolbutamide methylhydroxylation, mobile phase A consisted of acetonitrile, and mobile phase B consisted of water with 0.1% acetic acid (pH 3.5; 40:60), monitored at 230 nm. The flow rate was 1.2 mL/min. The retention times of 4-OH-Tol and Tol were 3.8 min and 11.3 min, respectively.

For CYP2D1 dextro-*O*-demethylation and CYP3A2 dextro-*N*-demethylation, the mobile phase consisted of a gradient of acetonitrile/water with acetic acid and 0.1% triethylamine (pH 4.5) monitored at 277 nm. The flow rate was 1.0 mL/min. The retention times of Dor, 3MM and Dex were 4.7 min, 10.6 min and 11.1 min, respectively.

For CYP2E1 chlorzoxazone 6-hydroxylation, mobile phase A consisted of acetonitrile and mobile phase B consisted of water with 0.1% acetic acid (pH 3.5; 40:60) monitored at 278 nm. The flow rate was 1.2 mL/min. The retention times of 6-OH-Chl and Chl were 3.1 min and 6.2 min, respectively.

For PTS, mobile phase A consisted of acetonitrile and mobile phase B consisted of water with 0.1% acetic acid (pH 3.5; 20:80) monitored at 230 nm. The flow rate was 1.2 mL/min. The retention times of the metabolites, internal standard (IS) and PTS were 3.0 min, 5.8 min and 9.7 min, respectively.

PTS incubation and kinetics assays To determine the kinetics of PTS metabolism, an incubation mixture contained PTS at 0, 10, 20, 40, 80, 100, 120, 200, 300, 400, and 600 $\mu\text{mol/L}$. The apparent K_m and V_{max} values were estimated by nonlinear regression analysis of V (enzyme activity) and $[S]$ (substrate concentration) using the Michaelis-Menten model:

$$V=V_{\max}[S]/(K_m+[S])$$

Inhibition study in rat liver microsomes The incubation mixture contained microsome protein, the relevant inhibitor, PTS, and NADPH-generating system. Various concentrations of α -Naph (1.0–50.0 $\mu\text{mol/L}$), Sulf (1.0–200.0 $\mu\text{mol/L}$), Qui (1.0–100.0 $\mu\text{mol/L}$), or Ket (0.5–20.0 $\mu\text{mol/L}$) were co-incubated with 40 $\mu\text{mol/L}$ PTS. Because PTS metabolite standard was not available, the analyte to internal standard (antipyrine) peak area ratio was used instead of using absolute quantitation^[11]. Both positive (in the presence of known inhibitors and specific substrates) and negative (in the presence of PTS or substrate and in the absence of the inhibitor) control samples were included in each assay to ensure the integrity of the microsomal incubation system. The result, expressed as percentage of control activity, was calculated based on a comparison between the peak area ratio of the sample and that of the negative control samples^[2].

The inhibitory effects of Sulf, Qui, and Ket on PTS metabolism were derived from Lineweaver-Burk plots of PTS metabolite formation by varying the concentration of the PTS at several fixed concentrations of the inhibitor^[12]. Sulf at concentrations of 0, 50, and 100 $\mu\text{mol/L}$, Qui at concentrations of 0, 50, and 100 $\mu\text{mol/L}$, or Ket at concentrations of 0, 5, and 10 $\mu\text{mol/L}$ were co-incubated with PTS at concentrations ranging from 10.0 to 160.0 $\mu\text{mol/L}$. To investigate the effect of PTS on CYP isoform activity, PTS at concentrations of 0–120 $\mu\text{mol/L}$ were added to the incubation mixture with CYP substrates. A single fixed concentration at approximately the K_m for each substrate (20 $\mu\text{mol/L}$ Phe, 100 $\mu\text{mol/L}$ Tol, 20 $\mu\text{mol/L}$ Dex, or 40 $\mu\text{mol/L}$ Chl) was used in the incubation sample. The areas for Ace, 4-OH-Tol, Dor, 3-MM and

6-OH-Chl were used for analysis.

Results

Total CYP and microsome protein content There was no significant difference in total CYP and microsome protein content between the PTS pretreatment and blank control groups. PTS pretreatment at the investigated concentrations had little effect on rat liver CYP content (Table 1).

Validation of PTS assay Validation of the HPLC method developed for PTS assay was performed with 6 calibration standards ranging from 10.0–320.0 $\mu\text{mol/L}$, and 3 quality control (QC) samples (20, 40, and 80 $\mu\text{mol/L}$). Standard samples were added to the boiled microsome mixture and prepared as described earlier. The calibration curve was constructed by linear least-squares regression of standard concentrations against peak area ratio of PTS and internal standard. The calibration curve $Y=0.065X+0.2265$ had excellent linearity, with a correlation coefficient of 0.9996 ($n=6$). The average recovery for the method was $106.2\% \pm 4.5\%$ from different concentrations. The relative standard deviations of intra-day and inter-day variation in the concentrations determined were less than 9.4%.

Mass spectrometry for PTS and its metabolite We hoped to gather more information about the metabolites and metabolism of PTS by *in vitro* incubation. In HPLC chromatograms, the retention times of the PTS metabolite and PTS itself were, respectively, 3.0 min and 9.7 min. There was no interfering peak found at the same retention time for the metabolite and PTS in the chromatogram for microsomal incubation. We collected the material with a retention time




Figure 1. HPLC chromatograms. (A) Blank microsomes with standard IS and PTS; peaks: 1 antipyrine (IS), 2 PTS. (B) PTS incubation; peaks: 3 PTS metabolite. (C) Phe incubation; peaks: 1 Ace, 2 Phe. (D) Tol incubation; peaks: 1 4-OH-Tol, 2 Tol. (E) Chl incubation; peaks: 1 6-OH-Chl, 2 Chl. (F) Dex incubation; peaks: 1 Dor, 2 3-MM, 3 Dex.

of approximately 3.0 min during the HPLC separation of the PTS incubation sample. This material was analyzed by using a Finnigan LCQ Deca XP Max (Sunnyvale, CA, USA) mass spectrometer. The metabolite with $[M+1]^+$ 188>171 in the mass spectrometry could be a hydroxylated derivative (Figure 1).

Kinetics assays of PTS metabolism relative to CYP

The K_m and V_{max} values were estimated using the Michaelis-Menten model. The K_m and V_{max} of the CYP-catalyzed reaction of PTS metabolism were 92.2 $\mu\text{mol/L}$ and 0.0137 nmol/min per mg protein, respectively (Figure 2).

Inhibition screening of PTS metabolism PTS metabolism was inhibited by Sulf, Qui and Ket. The results indicated that CYP2C7, CYP2D1, and CYP3A2 might be respon-

Table 1. Microsome protein and total CYP450 activity. $n=6$. Data are mean \pm SD. ^b $P<0.05$ vs blank control.

	Blank control	PB	PTS (mg/kg per d)	
			33	99
Microsome protein (mg/mL)	9.85 \pm 0.14	9.97 \pm 0.10	8.91 \pm 0.22	9.23 \pm 0.14
CYP450 content (nmol/mg)	0.61 \pm 0.08	1.36 \pm 0.13 ^b	0.66 \pm 0.17	0.60 \pm 0.12




Figure 2. PTS incubation data. Each data point represents the mean of three determinations. (A) Plot of velocity versus substrate concentration for the formation of PTS metabolites in rat liver microsomes. (B) Values from Lineweaver-Burk plots. [V], enzyme activity (nmol/min per mg).




Figure 3. Plots of percentage of control activity versus concentration of inhibitor. PTS (40 $\mu\text{mol/L}$) or specific substrates (positive control) were incubated with varying concentrations of inhibitors. Experiments were carried out in triplicate. $n=3$. Data are mean \pm SD.

sible for the CYP-catalyzed metabolism of PTS in liver microsomes of male Wistar rats (Figure 3).

Mechanisms of inhibitors The three inhibitors (Sulf, Qui and Ket) were assessed with respect to their inhibitory mechanisms. Lineweaver-Burk plots of PTS metabolite formation in the presence of Sulf, Qui and Ket are shown. As seen in Figure 4, PTS metabolism was inhibited by both Sulf and Ket through a mixed inhibitory mechanism. Qui had a noncompetitive mechanism. Noncompetitive inhibition mainly implies that an intermediate substance hinders the release of the substrate (Figure 4).

Effect of PTS on rat CYP isoform activity No significant changes were found with respect to the areas of Ace, 4-OH-Tol, Dor, 3-MM and 6-OH-Chl when the concentration of PTS changed from 0 to 120 $\mu\text{mol/L}$. This study indicated

that PTS had neither an inhibitive nor an inductive effect on the CYP1A2, CYP2C7, CYP2D1, CYP3A2 and CYP2E1 reactions selected in our experiments (Figure 5).




Figure 5. Various concentrations of PTS were co-incubated with substrate. The areas of Ace, 4-OH-Tol, Dor, 3-MM and 6-OH-Chl as determined from HPLC detection were used for analysis. Each data point represents the mean of three determinations.




Figure 4. Lineweaver-Burk plots of PTS metabolite formation curves. Several fixed concentrations of the inhibitor were incubated with PTS in varying concentrations from 10–120 $\mu\text{mol/L}$. (A) PTS incubation with Sulf at concentrations of 0, 50, and 100 $\mu\text{mol/L}$. (B) PTS incubation with Qui at concentrations of 0, 50, and 100 $\mu\text{mol/L}$. (C) PTS incubation with Ket at concentrations of 0, 5 and 10 $\mu\text{mol/L}$.

Discussion

Acquiring metabolic information and determining the effect of an NCE on CYP are important in developing clinically safe and efficient medications^[13]. Drug co-administration and individual differences in therapeutic effectiveness are common in cancer therapy. In the present article, we provide a relatively complete description of PTS metabolism involving CYP. This was a prerequisite for further toxicological risk assessment using this animal model.

In preclinical animal pharmacokinetic research, it was found that PTS could be given to rats at dosages of 33–198 mg/kg per d, for which the $T_{1/2}$ was 2.90–3.48 h, and the C_{max} was 29.24–151.23 mg/mL. There was one metabolite detected in the aqueous and organic phases, respectively, in urine samples. PTS might be used at higher concentrations in the clinical setting to treat different severities and types of tumors. Further experiments in human microsomes with different PTS concentrations are expected.

Because of various anti-tumor mechanisms, the anticancer drug degradation process could be very different and could have no relation with liver CYP enzymes at all. We arrived at a preliminary conclusion that CYP was responsible for PTS metabolism because of two pieces of evidence. First, inhibition screening studies showed that inhibition of the activity of CYP2C7, CYP2D1, and CYP3A2 slowed down the metabolism of PTS. Secondly, the result that CYP was responsible for PTS metabolism had also been verified by PTS metabolism research in a rat *in situ* liver perfusion model. In the latter experiment, there was difference in the PTS metabolism curves between the groups receiving Ket and PB

pretreatment and the blank controls. The inhibitors Sulf, Qui and Ket gave us further evidence that CYP2C7, CYP2D1, and CYP3A2 might contribute to PTS rat liver metabolism. However, specific CYP isoforms and different effective inhibitors (especially for CYP3A) for screening research will be needed to verify our results.

Inhibitors of CYP2C7, CYP2D1, and CYP3A2 could slow down PTS metabolism and potentiate its activity or toxicity. Of Sulf (up to 100 $\mu\text{mol/L}$, 56.8% of control activity), Qui (up to 100 $\mu\text{mol/L}$, 65.1%), and Ket (up to 20 $\mu\text{mol/L}$, 40.1%), Ket was the most effective inhibitor of PTS metabolism. Agents that modulate CYP3A will always need to be administered with care, and the same is true for PTS administration.

PTS had no effect on phenacetin *O*-deethylase, tolbutamide methylhydroxylation, dextro *O*-demethylation, dextro *N*-demethylation, or chlorzoxazone 6-hydroxylation reactions, which respectively represent the activities of CYP1A2, CYP2C7, CYP2D1, CYP3A2 and CYP2E1 *in vitro*. Our data predict that PTS could be used relatively safely with many current drugs, and that PTS would be suitable for pharmaceutical development. PTS has a simple structure, and it could be manufactured easily and cheaply. In conclusion, studies of CYP correlations with NCE metabolism provide a firm scientific basis for the safe and effective use of drugs, personalized medicine, and drug discovery.

References

- 1 Zhou JQ, Tang ZQ. Cytochrome P450 directed prodrug activation therapy in the research of cancer enzymology. *J Chin Pharm Sci* 2005; 14: 1–9.
- 2 Peng SX, Barbone AG, Ritchie DM. High-throughput cytochrome P450 inhibition assays by ultrafast gradient liquid chromatography with tandem mass spectrometry using monolithic columns. *Rapid Commun Mass Spectrom* 2003; 17: 509–18.
- 3 Afshar M, Rouini MR, Amini M. Simple chromatography method for simultaneous determination of dextromethorphan and its main metabolites in human plasma with fluorimetric detection. *J Chromatogr* 2004; 802: 317–22.
- 4 Testino SA Jr, Patonay G. High-throughput inhibition screening of major human cytochrome P450 enzymes using an *in vitro* cocktail and liquid chromatography-tandem mass spectrometry. *J Pharm Biomed Anal* 2003; 30: 1459–67.
- 5 Mimura N, Kobayashi K, Nakamura Y, Shimada N, Hosokawa M, Chiba K. Metabolism of medroxyprogesterone acetate (MPA) via CYP enzymes *in vitro* and effect of MPA on bleeding time in female rats in dependence on CYP activity *in vivo*. *Life Sci* 2003; 73: 3201–12.
- 6 Paul LD, Springer D, Staack RF, Kraemer T, Maurer HH. Cytochrome P450 isoforms involved in rat liver microsomal metabolism of californine and protopine. *Eur J Pharmacol* 2004; 485: 69–79.
- 7 Omura T, Sato R. The carbon monoxide-binding pigment of liver microsomes. *J Biol Chem* 1964; 239: 2379–85.
- 8 Lowry OH, Rosebrough NJ, Farr AL, Randall J. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951; 193: 265–75.
- 9 Liu XQ, Zhao Y, Li D, Qian ZY, Wang GJ. Metabolism and metabolic inhibition of cilnidipine in human liver microsomes. *Acta Pharmacol Sin* 2003; 24: 263–7.
- 10 Zhou JQ, Tang ZQ. Effect of quercetin on CYP1A2, CYP2E1, CYP3A2 and its inhibition mechanism from *in vitro* rat liver microsomal studies. *J Chin Pharm Sci* 2005; 14: 231–6.
- 11 Yu C, Shin YG, Kosmeder JW, Pezzuto JM, van Breemen RB. Liquid chromatography/tandem mass spectrometric determination of inhibition of human cytochrome P450 isozymes by resveratrol and resveratrol-3-sulfate. *Rapid Commun Mass Spectrom* 2003; 17: 307–13.
- 12 Kim KA, Lee JS, Park HJ, Kim JW, Kim CJ, Shim IS, *et al*. Inhibition of cytochrome P450 activities by oleanolic acid and ursolic acid in human liver microsomes. *Life Sci* 2004; 74: 2769–79.
- 13 Hu YZ, Yao TW. *In vitro* metabolism and inductive or inhibitive effect of DL111 on rat cytochrome P4501A enzyme. *Chem Biol Inter* 2004; 147: 109–17.