

Invited review

Sympathetically evoked Ca²⁺ signaling in arterial smooth muscle¹Wei-jin ZANG^{2,4}, Joseph ZACHARIA³, Christine LAMONT³, Withrow Gil WIER^{2,3}

²Department of Pharmacology, Key Laboratory of Environment and Genes Related to Diseases of Ministry of Education, School of Medicine, Xi'an Jiaotong University, Xi'an 710061, China; ³Department of Physiology, University of Maryland School of Medicine, Baltimore, Maryland 21201, USA

Key words

sympathetic nerve; Ca²⁺ signaling; arterial smooth muscles; receptors; junctional Ca²⁺ transients; confocal microscope

¹Project supported by grants from the National Institutes of Health (USA) (Grant No HL64708) and the National Natural Science Foundation of China (No 30470633), the Research Fund for the Doctoral Program (No 20050698012) and the Cultivation Fund of the Key Scientific and Technical Innovation Project, Ministry of Education of China (No 705045).

⁴Correspondence to Prof Wei-jin ZANG.

Phn 86-29-8265-5150.

Fax 86-29-8265-5003.

E-mail zwj@mail.xjtu.edu.cn

Received 2006-06-20

Accepted 2006-08-23

doi: 10.1111/j.1745-7254.2006.00465.x

Abstract

The sympathetic nervous system plays an essential role in the control of total peripheral vascular resistance and blood flow, by controlling the contraction of small arteries. Perivascular sympathetic nerves release ATP, norepinephrine (NE) and neuropeptide Y. This review summarizes our knowledge of the intracellular Ca²⁺ signals that are activated by ATP and NE, acting respectively on P2X₁ and α₁-adrenoceptors in arterial smooth muscle. Each neurotransmitter produces a unique type of post-synaptic Ca²⁺ signal and associated contraction. The neural release of ATP and NE is thought to vary markedly with the pattern of nerve activity, probably reflecting both pre- and post-synaptic mechanisms. Finally, we show that Ca²⁺ signaling during neurogenic contractions activated by trains of sympathetic nerve fiber action potentials are in fact significantly different from that elicited by simple bath application of exogenous neurotransmitters to isolated arteries (a common experimental technique), and end by identifying important questions remaining in our understanding of sympathetic neurotransmission and the physiological regulation of contraction of small arteries.

Introduction

The autonomic nervous system controls the activity of smooth muscles in the cardiovascular-renal system, gastrointestinal system, and urogenital system. Afferent and efferent (sensory) nerves release a 'cocktail' of neurotransmitters that control the contractile and trophic state of the smooth muscle cells^[1]. These neurotransmitters activate several cell signaling systems, including the ubiquitous intracellular Ca²⁺ signaling systems^[2]. Yet, neurogenic Ca²⁺ signaling in smooth muscle (that elicited by neurally released transmitters, as opposed to exogenously applied receptor agonists) has been investigated so far only in a few types of arteries, the vas deferens, the bladder and the uterus. Until now, this remarkable state of affairs reflected the difficulty of observing intracellular (Ca²⁺) in the muscle cells of intact organs in which nerves can be stimulated. Our intent in this article is to review what is known about the changes in intracellular (Ca²⁺) that are elicited by sympathetic nerves, and to

identify the important remaining questions.

Many factors are involved in regulating arterial diameter including the central and peripheral nervous systems, endothelial cells, circulating hormones, locally released substances, blood pressure, blood flow, and intrinsic mechanisms of smooth muscle. In arteries that contribute significantly to total peripheral resistance, such as mesenteric small arteries^[3,4] (which control blood flow to the intestine), control of smooth muscle contraction by the sympathetic nervous system is of major importance. This control is complex, involving multiple neurotransmitters and receptors, as well as complex patterns of nerve fiber activity. It is known that at least 3 different sympathetic neurotransmitters [ATP, norepinephrine (NE), and neuropeptide Y (NPY)] are released from sympathetic varicosities, and that their effects vary with the pattern of nerve fiber activity. Actions of the transmitters to elicit Ca²⁺ signaling in smooth muscle are believed to be synergistic, although detailed information is lacking. Finally, the probability of transmitter release at a single

sympathetic nerve terminal is low (~ 0.05) and the pattern of sympathetic nerve activity *in vivo* is complex, consisting of single action potentials (AP) or bursts of AP superimposed on a background of tonic activity^[5,6].

All 3 sympathetic cotransmitters, NE, ATP, and NPY contribute to sympathetically mediated vasoconstriction. Stimulation of the nerves supplying the rat mesenteric arterial bed (the tissue with which we are mainly concerned in this review) elicits an increase in perfusion pressure that can be blocked completely only by the combined administration of prazosin, suramin and BIBP 3226 ((*R*)-*N*2-(diphenacetyl)-*N*-(4-hydroxyphenyl) methyl)-*D*-arginineamide) (a selective α_1 -adrenoceptor antagonist, a non-selective antagonist of all P2X receptor subtypes, and a selective antagonist of the NPY receptor subtype, Y_1), respectively^[7]. Neurogenic contractions of isolated mesenteric small arteries are typically bi-phasic (Figure 1), with the small, initial transient component being attributed to purinergic (P2X) receptors, and the later, slow, large component being attributed to neurally released NE. However, the relative importance of each type of receptor depends on the size and type of the artery, as well as on the pattern of sympathetic nerve activity. In mesenteric arteries, the purinergic component of the contraction is




Figure 1. A neurogenic contraction from a rat small mesenteric artery. The artery was mounted between two glass pipettes. One pipette was attached to a servo-controlled pressure-regulating device (Living Systems), while the other was attached to a closed stopcock. The inter-luminal pressure was set to 70 mmHg. Electrical field stimulation (EFS) (via two platinum electrodes placed parallel to the long axis of the artery) was applied for the period indicated by the bar. EFS characteristics were 0.2 ms and 40V. The small initial component is attributed predominantly to purinergic receptors (P2X₁), the later slower, larger component being attributed to neurally released nor-epinephrine (previously unpublished data of the authors).

relatively large in very small arteries (compared to the α_1 -receptor-mediated component), and the purinergic component predominates during brief bursts of sympathetic nerve fiber activity^[8]. Indeed, it has recently been concluded that, “all three cotransmitters contribute significantly to vascular responses and their contribution varies markedly with impulse numbers”^[9]. The varying contribution of the transmitters under different conditions is the result of both pre- and post-synaptic factors.

Our goal in this review is to summarize recent new information on sympathetic neuromuscular transmission and the resultant Ca^{2+} signaling in the smooth muscle cells of small arteries during neurogenic contractions. Ca^{2+} signaling during neurogenic contractions activated by trains of sympathetic nerve fiber action potentials is in fact significantly different from that elicited by the simple application of exogenous neurotransmitters (both ATP and NE) to isolated arteries (or single isolated smooth muscle cells). We end by identifying important questions remaining in our understanding of sympathetic neuromuscular transmission and the physiological regulation of arterial smooth muscle contraction by the sympathetic nervous system.

Sympathetic nerves in isolated small arteries

The perivascular nerve fibers present in rat mesenteric small arteries are of 2 major types: sympathetic and ‘sensory’. The latter, which will not be discussed further here, are the so-called non-adrenergic, non-cholinergic (NANC) or ‘sensory’ nerves, whose cell bodies are located in the dorsal root ganglia. The perivascular sympathetic nerves that remain with isolated arteries consist of the neuroeffector system, that is, the post-ganglionic fibers (cell bodies located in paravertebral ganglia) and their nodal areas of axoplasmic specialization, that is, varicosities and synaptic junctions with the smooth muscle. Experimentally, the perivascular nerves of an isolated artery can easily be electrically stimulated to release transmitters. The function of NANC nerves may be blocked through the use of capsaicin^[10], thus permitting selective activation of the sympathetic nerves.

We next review very briefly the salient features of current concepts on the pre-synaptic and post-synaptic mechanisms that are involved in sympathetic neuromuscular transmission at such junctions, particularly those involved in the differential release of sympathetic cotransmitters. A major physiological phenomenon to be explained is that the effects of ATP are predominant at low frequencies of sympathetic nerve fiber activity, while those of NE are predominant at high frequencies.

Synaptic vesicles in sympathetic varicosities

Varicosities on the sympathetic perivascular nerves in rat mesenteric small arteries contain several different types of synaptic vesicles, probably containing different proportions of NE, ATP and NPY. These vesicles may have different origins (cell body vs formation at synapse) and mechanisms of exocytosis (including regulation by varicosity Ca^{2+}). Rat mesenteric arteries adrenergic nerve terminals contain 3 types of vesicles: large dense-cored vesicles (LDCV; ~100 nm diameter), small dense core vesicles (SDCV; ~50 nm diameter) and small clear vesicles (SCV; ~50 nm diameter)^[11-13]. LDCV comprises 5% of the total vesicles in a varicosity, with the remainder being small vesicles (SV)^[10]. The majority of the SV have (88%) dense cores (SDCV); the remainder without, SCV^[10]. NE and ATP are thought to be stored in all vesicle types, but in different proportions. It is generally agreed that the ATP and NE are released from different nerve terminal stores^[12,14] although the type of vesicles involved is not generally agreed upon. The exact vesicular origin of the released transmitters remains to be elucidated and may be species and tissue specific.

Post junctional receptors

Post-synaptic receptors for ATP Receptors for purine (ATP, ADP) and pyrimidine (UTP, UDP) nucleotides are presently divided into 2 families: ionotropic P2X receptors (7 cloned subtypes) and metabotropic P2Y receptors (6 cloned subtypes)^[15]. In the cardiovascular system, P2X receptors are expressed predominantly on smooth muscle (but are present on endothelial cells), and P2Y receptors are predominantly expressed on endothelial cells (also present on smooth muscle cells). P1 receptors will not be considered here except as they may be activated by adenosine produced by endothelial nucleotidase activity^[16,17]. Neurally-released ATP activated P2X receptors (ligand-gated ion channels) on smooth muscle to produce the excitatory junction potentials, and activated P2Y receptors on endothelium (G protein-coupled receptors) to produce endothelium dependent hyperpolarizing factor. In rat mesenteric arteries, the predominant P2X receptor is P2X₁^[18].

Neurally-released ATP binds to both P2X and P2Y receptors. While 'overflow' of ATP may be quantified chemically, only the effect of ATP on P2X receptors is detectable with electrical and optical methods. There is no doubt that neurally-released ATP produces the excitatory junction currents (EJCs), junctional Ca^{2+} transients (jCaT) and in vas deferens, the neuroeffector Ca^{2+} transients (NCT). jCaT and NCT are post-junctional changes in Ca^{2+} of smooth

muscle cells in response to neurally-released ATP. In mouse vas deferens, EJC are of 2 types: (1) large and fast; and (2) small and slow. The 2 types are differentially affected by external (Ca^{2+}) and osmotic pressure^[12] and sensitivity to heptanol^[19]. Considering the work of these authors and the 'dual vesicle' hypothesis of Stjarne^[12], it may be speculated that the 'large and fast' EJC arise from the release of the 'big quanta' type of SV, and that the 'small and slow' EJC arise from the 'small quanta' type of SV. jCaT and NCT would arise from the 'big quanta' SV; the extent to which work in the smooth muscle of vas deferens may apply to that of arteries is not known.

Post-synaptic receptors for NE The second sympathetic cotransmitter, NE, binds to adrenoceptors, the classification and function of which have been reviewed recently^[20]. At least 9 subtypes of vascular adrenoceptors have been identified^[21]: α_{1A} , α_{1B} , α_{1D} , $\alpha_{2A/D}$, α_{2B} , α_{2C} , β_1 , β_2 , and β_3 . There is also believed to be a low affinity (to prazosin) state of α_{1A} -adrenoceptors, known as the α_{1L} -adrenoceptors, and they are found in various vascular beds^[22]. A useful heuristic generalization is that α_1 -adrenoceptors are located post-junctionally on vascular smooth muscle and have a primary role in controlling arterial tone, particularly in small resistance arteries. This accounts for the predominant use of the synthetic α_1 -adrenoceptor agonist, phenylephrine (PE), in the majority of experimental studies. Exogenous activation of adrenoceptors using PE and the use of an α_{1B} -knockout mouse has permitted the conclusion that α_{1D} is most important in conduit arteries, while α_{1A} is most important in small arteries and α_{1B} was reported to have a minor contribution^[23]. Recent studies using nerve-evoked contraction, also in small arteries isolated from α_{1D} -knockout mouse, have shown the importance of α_{1A} -adrenoceptors as the predominant subtype and α_{1D} having a small, but significant role^[24]. Pre-junctionally, transmitter release at sympathetic varicosities (neuromuscular junctions) is importantly regulated by $\alpha_{2A/D}$ and α_{2C} . Endothelial cells have at least 5 subtypes of adrenoceptors: $\alpha_{2A/D}$, α_{2C} , β_1 , β_2 , and β_3 . Adrenergic signaling mechanisms in arterial smooth muscle have been reviewed recently^[25], with the emphasis on Ca^{2+} activation of contraction and Ca^{2+} -sensitizing mechanisms activated by bath-applied phenylephrine.

Neurally-released NE has been detected classically by chemical methods in the effluent of neurally stimulated arteries ('overflow experiments') or by amperometric methods, in which the oxidation of NE on the surface of a carbon fiber electrode is detected as an electrical current. Most significantly, the release of single quanta of NE has been detected from the surface of rat mesenteric small arteries

through the use of carbon fiber microelectrodes (CFmE). These microelectrodes are 7 μm in diameter and are believed to detect the release of NE quanta from a distance of 8 μm ^[14]. Furthermore, spontaneous oxidation currents (SOC) were also recorded. These authors speculated that the SOC arose from NE released from large dense-cored vesicles (LDV). An important observation was that α -latrotoxin increased the frequency of SOC about 4-fold, but increased the frequency of spontaneous excitatory junction potentials (which monitor packeted or quantal ATP release) by 30-fold. This observation supports the suggestion that SEJP (spontaneous excitatory junction potentials activated by released ATP) and SOC (NE release) occur through different synaptic vesicles, under these conditions.

In most vascular beds, α_1 -adrenoceptors play an important role in vasoconstriction. The relative importance of the different α_1 -adrenoceptor subtypes in regulation of peripheral resistance and systemic arterial blood pressure is not clear, as the contribution of different subtypes to vasoconstriction differs with the mode of activation and species^[26]. For example, studies in rat mesenteric small arteries using exogenous agonists have revealed the role of α_{1A} ^[27,28] or α_{1B} ^[29] or α_{1L} -adrenoceptors^[30]. In contrast, nerve-evoked contractions in rat mesenteric arteries were predominantly mediated by α_{1A} -adrenoceptors^[27,28]. Further, studies in mesenteric small arteries using α_{1B} -adrenoceptor knockout mouse have revealed the predominance of α_{1A} -adrenoceptors in vasoconstriction to phenylephrine^[23] and nerve-evoked contractions showed the predominance of α_{1B} -adrenoceptors^[31]. Therefore, it is important to study *in vivo* preparations to understand the clinical relevance of the α_1 -adrenoceptor subtypes involved in vasoconstriction.

Post-synaptic receptors for NPY The third sympathetic cotransmitter, NPY, is believed to enhance the effects of both ATP and NE, by acting on post-junctional NPY- Y_1 receptors^[32]. Five distinct NPY receptors (Y_1 , Y_2 , Y_4 , Y_5 , and Y_6) have been cloned^[33]. Y_1 receptors are believed to be the major type present post-junctionally in the cardiovascular system and to mediate the response to NPY, although the possible involvement of pre- and post-junctional Y_2 receptors has been suggested. NPY receptors act via pertussis toxin-sensitive G proteins. A major effect of their activation is the inhibition of adenylyl cyclase. NPY shifts agonist dose-response curves to the right^[34]. It has been suggested that NPY activates mesenteric small arteries through 2 different mechanisms: activation of non-selective cation channels and consequent Ca^{2+} entry, and the inhibition of the hyperpolarization produced by cAMP^[35] (cyclic adenosine monophosphate).

Differential release of ATP and NE There is good evidence that ATP is the predominant sympathetic effector in response to 'single' action potentials, while at high frequencies of activation, NE is the predominant sympathetic effector. This may result from the pre- and post-synaptic mechanisms discussed earlier. Low frequency electrical field stimulation (EFS; eg 1 Hz) elicits relatively small, brief, phasic contraction of rat mesenteric arteries, whereas higher frequency EFS (eg 16 Hz) elicits larger, sustained contractions. The irreversible adrenergic antagonist, phenoxy-benzamine, abolished the response to bath-applied norepinephrine, but reduced the response to a single nerve stimulus by only 20%^[36]. Conversely, the stable ATP analogue, α,β -methylene ATP (which desensitizes P2X receptors), reduced the response to a single pulse by 70%, while reducing the contraction to high frequency stimulation only 10%. Similarly, in the rabbit ear artery, it has been concluded that 'short pulse bursts at low frequency favor the prazosin-resistant (purinergic) component of the response'^[37].

Post-synaptic Ca^{2+} signaling

Recent work has revealed in detail the Ca^{2+} signals in smooth muscle cells during sympathetically-mediated neurogenic contractions of small arteries^[38] and vas deferens^[39]. There are differences between these 2 tissues (which have very different functions), and we will focus here on what is known about sympathetic neuromuscular transmission and neurogenic contractions of vascular smooth muscle. The goals of the current research in this area are to attribute different types of post-synaptic Ca^{2+} signals to the activation of specific receptors, channels, and organelles, and to determine to what extent the different post-junctional Ca^{2+} signals actually activate contraction. Here, we summarize the recent studies in which Ca^{2+} signals, attributable to both neurally-released ATP and neurally released NE, have been observed.

jCaT: post-synaptic Ca^{2+} signals activated by neurally released ATP Neurally-released ATP activates a specific, localized, Ca^{2+} transient in arterial smooth muscle cells^[40] that has been termed a 'jCaT', for junctional Ca^{2+} transient. It was shown that these Ca transients arise in the vicinity of perivascular nerves^[38] or even directly beneath single visualized nerve fibers^[40]. The confocal images of jCaT in Figure 2 were obtained in pressurized (70 mmHg) rat mesenteric small arteries subjected to EFS. Low frequency, low voltage stimulation (0.67 Hz, 0.2 ms) excited nerve fibers without causing an appreciable contraction. This was referred to this as 'sub-threshold' EFS, as it is sub-threshold for muscle




Figure 2. Junctional Ca^{2+} transients in pressurized small artery. (A) Images of perivascular nerves and smooth muscle cells (SMC). The image is an average of 30 frames (1 s total), from a set of 600 images obtained with a real-time confocal microscope during EFS. (B) A virtual line-scan image was derived from this data, by taking a single line of pixels, extending through the image vertically, between the two black arrowheads in A, from each of the 600 frames. The chosen line crosses prominent nerve fibers at the top of the image, bisects a smooth muscle cell longitudinally, and crosses a region (white arrow in A) in which a small nerve fiber crosses the muscle cell. In the line-scan image, nerve fiber Ca^{2+} transients in the large fibers at the top of are seen as periodic (0.5 s^{-1}) increases in fluorescence occurring at the times of the stimulus pulses (black arrowheads). A jCaT is seen in the SMC at 4.8 s in the line-scan image. (C) jCaTs often arise at streaks in line-scan images. (D) A region where a jCaT occurred was scanned repeatedly. In (a) a jCaT occurred. In (b) no jCaT occurred, but a fluorescence transient occurred in the streak. We demonstrate that jCaTs can arise precisely in the region of an electrically excitable nerve fiber (from Lamont & Wier, 2002^[40]).

contraction. Thus, in these experiments, motion did not occur and the characteristics of the jCaT could be studied in detail. Figure 2 illustrates the basic appearance of jCaT in line-scan images and demonstrates that: (1) nerve fibers are being excited by each EFS pulse; (2) a jCaT occurs nearly simultaneously with an EFS pulse; (3) jCaT occur near nerve fibers; and (4) jCaT are events of very low probability.

Spatio-temporal characteristics of jCaT JCaT are larger in spatial spread and last longer than spontaneous Ca^{2+} sparks. JCaT always occur with brief latency to the EFS pulse. The spatio-temporal differences between the jCaT and the sparks are obvious in published records^[40]: the jCaT is larger in space, lasts longer, and occurs at the time of the stimulus pulse. These characteristics and their pharmacology are how jCaT are distinguished from sparks. The vast majority of jCaT occurred within 12 ms (4 scan lines) after the stimulus pulse. The evidence linking the occurrence of jCaT to the stimulus seems unequivocal. The spatial full-width-at-half-maximum (FWHM) for jCaT is $4.8 \mu\text{m}$, and the time taken to fall to half-amplitude, $t_{1/2}$, (from the peak) is 145 ms. The means of these distributions are quite different from those of sparks in smooth muscle (sparks: $t_{1/2}$, 48–56 ms;

FWHM, $2.4 \mu\text{m}$)^[41]. Some jCaT occurred before the stimulus, and some much after; we hypothesize that these are associated with spontaneous neurotransmitter release.

Pharmacology of jCaT Pharmacological studies of jCaT^[40] have provided strong evidence that they arise from the activation of purinergic receptors. JCaT persist, apparently unchanged, in the presence of capsaicin, and are thus not dependent on sensory nerves. They are completely absent in the presence of the purinergic receptor blocker, suramin ($300 \mu\text{mol/L}$). They persist in the presence of α_1 -adrenergic blocker prazosin ($10 \mu\text{mol/L}$), sufficient to block neurogenic adrenergic responses completely. They are also largely unaffected by ryanodine ($30 \mu\text{mol/L}$), while Ca^{2+} sparks are abolished. Although we favor the hypothesis that jCaT are due to ATP and P2X_1 receptors, further studies are required, particularly to be sure that it is the P2X_1 receptor subtype. NPY is also reported to activate non-specific cation channels^[36] and thus it could contribute to jCaT. This possibility could be tested using specific Y_1 receptor antagonists (BIBP 3226).

In our recent study^[42] using P2X_1 knockout animals^[43], we clearly showed that jCaT represent Ca^{2+} that enter vascu-

lar smooth muscle cells through P2X₁ receptors activated by neurally-released ATP. The P2X₁ knockout models also showed the importance of the involvement of ATP-mediated P2X₁ activation in the initial rapid component of the nerve-evoked contraction^[42].

Ca²⁺ waves: post-synaptic Ca²⁺ signals activated by α_1 -adrenergic agonists The Ca²⁺ signaling elicited by adrenergic agonists has been studied mainly through the use of exogenous, bath-applied synthetic catecholamines such as the α_1 -adrenoceptor specific agonist, PE. An early study was that of Zang and colleagues^[44], in which it was shown that PE elicited asynchronous propagating Ca²⁺ waves in rat mesenteric arteries. The basis of the dose-response to PE was increasing recruitment of individual smooth muscle cells to produce Ca²⁺ waves, and to produce them at higher and higher frequencies. As expected, asynchronous propagating Ca²⁺ waves seem also to be the response elicited by neurally released NE. This is discussed in more detail later in relation to the activation of contraction by neurally released NE, during neurogenic contractions of small arteries.

Neurogenic contractions of small arteries

The detailed studies on jCaTs described above were performed in pressurized small arteries that did not contract because the electrical stimulation was 'sub-threshold' for contraction. In this section, we review the studies in which jCaT and Ca²⁺ waves have been observed during isometric neurogenic contractions. For these studies, the arteries were mounted in a myograph that permitted simultaneous (i) high-speed confocal imaging of fluorescence from individual smooth muscle cells; (ii) electrical stimulation of perivascular nerves; and (iii) recording of isometric tension. Sympathetic neuro-muscular transmission was achieved by EFS (frequency, 10 Hz; pulse voltage, 40 V; pulse duration, 0.2 ms) in the presence of capsaicin and scopolamine (to inhibit 'sensory' and cholinergic nerves, respectively). As shown in Figure 3, during the first 20 s of EFS, force rose to a small peak, then declined, similar to that recorded previously^[45]. During this time, jCaT were present at a relatively high frequency. Propagating asynchronous Ca²⁺ waves, previously associated with bath-applied α_1 -adrenoceptor agonists, were not initially present. During the next 2.5 min of EFS, force rose slowly, and asynchronous propagating Ca²⁺ waves appeared. The selective α_1 -adrenoceptor antagonist, prazosin, abolished both the slowly developing contraction and the Ca²⁺ waves, but reduced the initial transient contraction by only ~25%.

Purinergic component of neurogenic contraction In

order to study selectively the arterial contractions generated by neurally-released ATP, arteries were exposed to prazosin (1–10 μ mol/L) to block α_1 -adrenergic receptors (Figure 4). Others^[8] have shown that purinergic receptor antagonists, such as suramin, abolish the small contractions that remain in prazosin. We found that after prazosin treatment 73.7% \pm 14.0% ($n=7$) of the initial transient contraction remained and 5.00% \pm 0.98% ($n=5$) of the maintained contraction. We then sought to characterize the changes in frequency and amplitude of jCaT that might occur during the EFS, and which activate contraction. Because the jCaT is a local Ca²⁺ signal, it was not necessarily clear that jCaT would activate contraction effectively. Confocal imaging of Fluo-4 fluorescence at 30 images·s⁻¹ was performed for 3 periods of 20 s in the beginning (0–20 s), middle (80–100 s), and end (160–180 s) of 3 min EFS (periods indicated by bars in Figure 4A). The frequency of jCaT declined markedly during the 3 min of EFS (Figure 4B, iv–vi). In contrast to the frequency, the peak amplitude of the jCaT changed little during 3 min EFS. jCaT occurred in sufficient numbers during the first 20 s of EFS to produce a detectable elevation of average (Ca²⁺; fluorescence ratio), which paralleled the transient contraction that occurred during this time. On the other hand, jCaT occurred at a very low frequency later in the EFS, when contractile force fell to very low levels. Thus, it seems reasonable to attribute the contractile activation to jCaT, despite their limited spatial extent and frequency.

Adrenergic component of neurogenic contraction Later during neurogenic contractions, asynchronous Ca²⁺ waves propagated within individual smooth muscle cells of the arterial wall. A representative Ca²⁺ signal, obtained as the average fluorescence with an area of interest (AOI; 1.35 mm square) within a single smooth muscle cell is shown in Figure 5A (lower trace, right hand scale). For this data, images were obtained at 2 s⁻¹; a rate which is too slow to resolve the jCaT^[40] generated during the initial purinergic component. For the analysis of the adrenergic component of the experiment illustrated in Figure 5, 74 individual smooth muscle cells were identified and an AOI placed on each. In these 74 cells, 809 Ca²⁺ waves were detected and the time of onset and peak amplitude of each was determined, and the data presented as a histogram.

Different roles of ryanodine (RyR) and inositol(1,4,5)-trisphosphate receptors (InsP₃R) in neurogenic contractions The role of the sarcoplasmic reticulum (SR) and its Ca²⁺-release channels (RyR, (Ins(1,4,5)P₃R) of individual smooth muscle cells of the arterial wall and in these Ca²⁺ signals is not completely known. Zang *et al*^[44] used confocal laser scanning microscopy and Fluo-4 to visualize Ca²⁺




Figure 3. Occurrence of Junctional Ca^{2+} transients (jCaTs) during the predominantly purinergic component (A) and Ca^{2+} waves during the predominantly adrenergic component (B) of neurogenic contraction. Records at left of both (A) and (B) are the force produced by EFS for 3 minutes. The dotted vertical bars indicate the times during which images in (a)–(c) and traces in (d) were obtained. (Aa and Ba) Image obtained by averaging 30 frames (1 sec) during the period indicated. Ab and Bb are virtual line-scan images derived from a selected single vertical line of pixels in images Aa and Bb, respectively (solid white line). Ac and Bc are virtual line-scan images derived from a single horizontal line of pixels in images Aa and Ba, respectively (dotted white line). Traces in (Ad) are fluorescence pseudo-ratios (F/F_0) derived from the average fluorescence within selected areas-of-interest (AOIs) in the images in Aa (white boxes). Traces in (Bd) are derived from the line-scan image in Bb, at the lines indicated by the black arrows. All the data illustrated here were obtained at 30 images·s⁻¹ (from Lamont, Vainorius & Wier 2003^[38]).

transients within individual smooth muscle cells of rat resistance arteries during α_1 -adrenoceptor activation. They noticed that in the presence of PE, caffeine also elicited a massive release of Ca^{2+} , at a time when Ca^{2+} waves had died away completely, or when further responses to PE would have been much diminished. This result shows that caffeine-sensitive Ca^{2+} stores are not depleted of Ca^{2+} in the presence of PE and further indicates that the α_1 -agonist-releasable Ca^{2+} store and the caffeine releasable Ca^{2+} store are different^[44]. More recent experiments by our group (Lamont and Wier, 2004)^[46] showed that Ins(1,4,5)P₃R are essential for adrenergically-induced asynchronous Ca^{2+} waves and the associ-

ated steady vasoconstriction, but RyR are not appreciably opened during adrenergic activation (because PE did not facilitate the development of the effects of ryanodine). Also Ins(1,4,5)P₃R are not essential for Ca^{2+} sparks. This provides an explanation of the fact that adrenergic stimulation decreases the frequency of Ca^{2+} sparks (previously reported) while simultaneously increasing the frequency of asynchronous propagating Ca^{2+} waves; different SR Ca^{2+} -release channels are involved^[46].

Both the contraction and the underlying Ca^{2+} signals during the adrenergic component of the neurogenic isometric contraction are also distinctly different from those occur-




Figure 4. Characteristics of junctional Ca^{2+} transients (jCaTs) during 3 minutes EFS in the presence of prazosin. A, Upper trace, force under control conditions. Lower trace, force in the presence of prazosin ($10 \mu\text{mol/L}$). Bars below the force records indicate the times during which imaging was performed and during which the data illustrated in (B) was obtained. B, (a–c) Scatter plots of the peak fluorescence ratios of jCaTs, as a function of their time-of-occurrence, obtained during the three periods of time indicated by the bars in (A); 0–20 s, 80–100 s, and 160–180 s. Different symbols represent data from different arteries. B, (d–f) Frequency histograms of jCaTs during the same periods. Bin size; 1s. Data in (B) obtained from 5 arteries, consisting of 549 jCaTs. Mean values of peak fluorescence ratio (F/F_0) are: 2.033 ± 0.0261 ($n=408$), 1.751 ± 0.040 ($n=90$), and 1.782 ± 0.0648 ($n=51$), in (a), (b), and (c), respectively (from Lamont, Vainorius & Wier 2003^[38]).




Figure 5. Characteristics of Ca^{2+} waves during the predominantly adrenergic component of neurogenic contraction. A, Recordings of force (upper trace) and fluorescence pseudo-ratio (lower trace) from a representative single cell during 3 minutes of EFS. B, Frequency histogram of the times of occurrence of Ca^{2+} waves in 809 waves from 74 cells in one artery. C, Scatter plot of peak pseudo-fluorescence ratio as a function of time of occurrence of the Ca^{2+} waves (from Lamont, Vainorius & Wier 2003^[38]).

ring during externally applied α_1 -adrenoceptor agonist (typically PE). After the initial purinergic component, the adrenergic component of the neurogenic contractions, even at maximally effective EFS, rises much more slowly than does the contraction in response to bath-applied α_1 -adrenoceptor agonist. Maximally effective concentrations of exogenous PE elicit an initial synchronous release of Ca^{2+} , followed by asynchronous propagating Ca^{2+} waves, both in veins^[45] and

in the arteries^[34,44]. Thus, the initial rapid rise in force in response to externally applied PE appears to be generated by a synchronous release of Ca^{2+} from intracellular stores. This does not occur during neurogenic contractions, possibly because neuronally released NA does not initially reach the uniformly high levels that are achieved rapidly after external application.

Influence of myogenic tone

Another significant complication that arises from arteries pressurized at 70 mmHg above room temperature is the development of tone. Earlier studies (eg Mauban *et al*^[47]), were at room temperature, which is not conducive for the development of myogenic tone. The only previous study in which spatially resolved imaging was performed in mesenteric small arteries at 37 °C^[48] was complicated by the development of adrenoceptor-mediated vasomotion, preventing clear observation of Ca²⁺ signaling in individual smooth muscle. However, mouse mesenteric small arteries at 32 °C showed the development of myogenic tone and also rarely resulted in vasomotion on stimulation by PE. In such arteries, PE elicited only a spatially uniform increase in Ca²⁺ with little or no Ca²⁺ waves, suggesting that the rise in Ca²⁺_i leads to the development of myogenic tone and also inactivation of IP₃ receptors^[49].

Conclusion

The confocal microscope, along with other new technologies has provided much new information on the physiology and pharmacology of sympathetically evoked Ca²⁺ signaling in arterial smooth muscle. We have previously advanced a scheme to explain the Ca²⁺ signals and isometric contraction elicited by electrical field stimulation of perivascular sympathetic nerves of a rat mesenteric small artery (Figure 6). Early during a train of nerve fiber action potentials, smooth muscle contraction is activated mainly by jCaT induced by neurally released ATP. jCaTs are localized to the post-junctional region, and arise from Ca²⁺ that has entered via P2X₁ receptors. At this time, sympathetic varicosities may release mainly synaptic vesicles that contain a relatively high concentration of ATP (the relatively few ‘big’ quanta proposed by Stjarne^[12]). Later during a train of nerve fiber action potentials, jCaTs are rare, and contraction is activated by Ca²⁺ waves that arise from SR. Ca²⁺ release from SR is




Figure 6. Real and hypothetical events of sympathetic neuromuscular transmission in a small artery. (A) Early during a train of nerve fiber action potentials, smooth muscle contraction is activated mainly by post-junctional Ca²⁺ transients (‘jCaTs’) induced by neurally released ATP. jCaTs are localized to the post-junctional region, and arise from Ca²⁺ that has entered via P2X receptors. At this time, sympathetic varicosities may release mainly small vesicles that contain a relatively high concentration of ATP (*viz* the relatively few ‘big’ quanta proposed by Stjarne, 2001). (B) Later during a train of nerve fibre action potentials, jCaTs are rare, and contraction is activated by Ca²⁺ waves that arise from sarcoplasmic reticulum (SR). Ca²⁺ release from SR is activated by InsP₃, produced after binding of NA (norepinephrine) to α₁-adrenoceptors. At this time, sympathetic varicosities may release small synaptic vesicles (the more numerous ‘small’ quanta, green) that contain a relatively high concentration of NA. (Figure reproduced from Wier GW and Lamont C. Confocal Ca imaging in intact small arteries reveals smooth muscle Ca²⁺ transients attributable to neurally released jATP and NA. *J Physiol* 2004; 557P: SA 18, Research Symposium).

activated by InsP_3 , produced after the binding of NE to α_1 -adrenoceptors. At this time, sympathetic varicosities may release small synaptic vesicles (the more numerous 'small' quanta, green; Figure 6) that contain a relatively high concentration of NE. We stress that the mechanisms accounting for differential release of ATP and NE are quite speculative at this time.

JCaT are distinct from Ca^{2+} transients activated in isolated venous myocytes by exogenously applied ATP^[50]. Previous studies of the effects of ATP on small arteries utilized spatially averaged measurements of Ca^{2+} ^[51] and we can not determine therefore whether jCaT might be produced by bath-applied ATP or not. In rat mesenteric small arteries similar to those used here, low (0.01–1 mmol/L) concentrations of exogenous ATP caused 'global' Ca^{2+} transients that seemed to involve Ca^{2+} influx through channels sensitive to nifedipine and the putative blocker of receptor operated channels, SKF 96365^[52], whereas higher concentrations (1–3 mmol/L) caused a release of Ca^{2+} from intracellular stores (Ca^{2+} transients were elicited by high ATP in the absence of external Ca^{2+}). When applied to isolated venous myocytes, low concentrations of ATP (0.1 $\mu\text{mol/L}$) induced rather uniform increases in Ca^{2+} (which started from the edges of the cell), and at higher concentrations (1 $\mu\text{mol/L}$), propagating Ca^{2+} waves^[50].

In the arteries studied here in the presence of prazosin, no propagating Ca^{2+} waves were ever observed during EFS. We interpret this to mean that neurally-released ATP does not evoke significant release of Ca^{2+} from intracellular stores, a result in agreement with previous pharmacological studies on rat mesenteric small arteries^[7]. We speculate that the differences between the effects of bath-applied ATP and neurally-released ATP are due to a markedly different spatio-temporal pattern of ATP on the smooth muscle cell in the 2 cases.

Both the contraction and the underlying Ca^{2+} signals during the adrenergic component of the neurogenic isometric contraction are also distinctly different from those occurring during externally applied α_1 -adrenoceptor agonist (typically phenylephrine, PE). After the initial purinergic component, the adrenergic component of the neurogenic contractions, even at maximally effective EFS, rises much more slowly than the contraction in response to bath-applied α_1 -adrenoceptor agonist. Maximally effective concentrations of exogenous PE elicit an initial synchronous release of Ca^{2+} , followed by asynchronous propagating Ca^{2+} waves, both in veins^[53] and arteries^[44,47]. Thus, the initial rapid rise in force in response to externally applied PE appears to be generated by a synchronous release of Ca^{2+} from intracellular stores. This does not occur during neurogenic contractions, possibly because neurally-released NE does

not initially reach the uniformly high levels that are achieved rapidly after external application.

Acknowledgement

We gratefully acknowledge the assistance of Becky SAUNDERS in the preparation of this manuscript.

References

- 1 Todorov LD, Mihaylova-Todorova ST, Bjur RA, Westfall DP. Differential cotransmission in sympathetic nerves: role of frequency of stimulation and prejunctional autoreceptors. *J Pharmacol Exp Ther* 1999; 290: 241–66
- 2 Cheng H, Wei S, Verkhatsky A. Calcium signaling in physiology and pathophysiology. *Acta Pharmacol Sin* 2006; 27: 767–72.
- 3 Christensen KL, Mulvany MLJ. Mesenteric arcade arteries contribute substantially to vascular resistance in conscious rats. *J Vas Res* 1993; 30: 73–9.
- 4 Fenger-Gron J, Mulvany MJ, Christensen KL. Intestinal blood flow is controlled by both feed arteries and microcirculatory resistance vessels in freely moving rats. *J Physiol* 1997; 498: 215–24.
- 5 Johnson CD, Gilbey MP. Focally recorded single sympathetic postganglionic neuronal activity supplying rat lateral tail vein. *J Physiol* 1998; 508 (Pt 2): 575–85.
- 6 Johnson CD, Gilbey MP. Effects of aortic nerve stimulation on discharges of sympathetic neurons innervating rat tail artery and vein. *Am J Physiol* 1998; 275: R942–9.
- 7 Donoso MV, Steiner M, Huidobro-Toro JB. BIBP 3226, suramin and prazosin identify neuropeptide Y, adenosine 5'-triphosphate and noradrenaline as sympathetic cotransmitters in the rat mesenteric bed. *J Pharmacol Exp Ther* 1997; 282: 691–8.
- 8 Gitterman DP, Evans RJ. Nerve evoked P2X receptor contractions of rat mesenteric arteries; dependence on vessel size and lack of role of L-type calcium channels and calcium induced calcium release. *Br J Pharmacol* 2001; 132: 1201–8.
- 9 Bradley E, Law A, Bell D, Johnson CD. Effects of varying impulse number on cotransmitter contributions to sympathetic vasoconstriction in rat tail artery. *Am J Physiol* 2003; 284: H2007–14.
- 10 Holzer P. Capsaicin: cellular targets, mechanisms of action, and selectivity for thin sensory neurons. *Pharmacol Rev* 1991; 43: 143–201.
- 11 Tranzer JP. New aspects of the localisation of catecholamine in adrenergic neurons. *Frontiers Catecholamine Res* 1973; 453–8.
- 12 Stjarne L. Novel dual 'small' vesicle model of ATP- and noradrenaline-mediated sympathetic neuromuscular transmission. *Auton Neurosci* 2001; 87: 16–36.
- 13 Blair DH, Lin YQ, Bennett MR. Differential sensitivity to calcium and osmotic pressure of fast and slow ATP currents at sympathetic varicosities in mouse vas deferens. *Auton Neurosci* 2003; 105: 45–52.
- 14 Brock JA, Dunn WR, Boyd NSF, Wong DKY. Spontaneous release of large packets of noradrenaline from sympathetic nerve terminals in rat mesenteric arteries *in vivo*. *Br J Pharmacol* 2000; 131: 1507–11.
- 15 Ralevic V. The involvement of smooth muscle P2X receptors in the prolonged vasorelaxation response to purine nucleotides in the rat mesenteric arterial bed. *Br J Pharmacol* 2002; 35: 1988–94.
- 16 North RA, Surprenant A. Pharmacology of cloned P2X receptors.

- Ann Rev Pharmacol Toxicol 2000; 40: 563–80.
- 17 Fredholm BB, Abbracchio MP, Burnstock G, Dubyak GR, Harden TK, Jacobson KA, *et al*. Towards a revised nomenclature for P₁ and P₂ receptors. Trends Pharmacol Sci 1997; 18: 79–82.
 - 18 Lewis CJ, Evans RJ. Comparison of P2X receptors in rat mesenteric, basilar and septal (coronary) arteries. J Auton Nerv Sys 2000; 81: 69–74.
 - 19 Manchanda R, Venkateswarlu K. Quantal evoked depolarizations underlying the excitatory junction potential of the guinea-pig isolated vas deferens. J Physiol 1999; 520.2: 527–37.
 - 20 Guimaraes S, Moura D. Vascular adrenoceptors: An update. Pharmacol Rev 2001; 53: 319–56.
 - 21 Alexander SPH, Peters JA. Receptor and ion channel nomenclatures. Trends Pharmacol Sci 1998; Suppl: 1–98.
 - 22 Ford AP, Daniels DV, Chang DJ, Gever JR, Jasper JR, Lesnick JD. Pharmacological pleiotropism of the human recombinant α_{1A} -adrenoceptor: implications for α_1 -adrenoceptor classification. Br J Pharmacol 1997; 121: 1127–35.
 - 23 Daly CJ, Deighan C, McGee A, Mennie D, Ali Z, McBride M, *et al*. A knockout approach indicates a minor vasoconstrictor role for vascular alpha 1B-adrenoceptors in mouse. Physiol Genomics 2002; 9: 85–91.
 - 24 Zacharia J, Hillier C, Tanoue A, Tsujimoto G, Daly CJ, McGrath JC, *et al*. Br J Pharmacol 2005; 146: 679–91.
 - 25 Wier WG, Morgan KG. Adrenergic signaling mechanisms in mammalian resistance arteries. Rev Physiol Biochem Pharmacol 2003; 150: 91–139.
 - 26 Vargas HM, Gorman AJ. Vascular alpha-1 adrenergic receptor subtypes in the regulation of arterial pressure. Life Sci 1995; 57: 2291–308.
 - 27 Kong JQ, Taylor DA, Fleming WW. Functional distribution and role of α_1 -adrenoceptor subtypes in the mesenteric vasculature of the rat. J Pharmacol Exp Ther 1994; 268: 1153–9.
 - 28 Williams T, Clarke D. Characterization of α_1 -adrenoceptors mediating vasoconstriction to noradrenaline and nerve stimulation in the isolated perfused mesentery of rat. Br J Pharmacol 1995; 114: 531–6.
 - 29 Piascik MT, Hrometz SL, Edelmann SE, Guarino RD, Hadley RW, Brown RD. Immunocytochemical localization of the α_{1B} adrenergic receptor and the contribution of this and the other subtypes to vascular smooth muscle contraction: analysis with selective ligands and antisense oligonucleotides. J Pharmacol Exp Ther 1997; 283: 854–68.
 - 30 Stam WB, Van Der Graaf PH, Saxena PR. Analysis of α_{1L} -adrenoceptor pharmacology in rat small mesenteric artery. Br J Pharmacol 1999; 127: 661–70.
 - 31 Townsend SA, Jung AS, Gillian Hoe YS, Lefkowitz RY, Khan SA, Lemmon CA. Critical role of α_{1B} -adrenergic receptor at sympathetic neuroeffector junction. Hypertension 2004; 44: 776–82.
 - 32 Westfall TC, Yang CL, Curfman-Falvey M. Neuropeptide -Y-ATP interactions at the vascular sympathetic neuro-effector junction. J Cardiovasc Pharmacol 1995; 26: 682–7.
 - 33 Michel MC, Beck-Sickinger A, Cox H, Doods HN, Herzog H, Larhammar D, *et al*. International Union of Pharmacology recommendations for the nomenclature of neuropeptide Y, peptide YY, and pancreatic polypeptide receptors. Pharmacol Rev 1998; 50: 143–50.
 - 34 Chen H, Fetscher C, Schäfers RF, Wambach G, Philipp T, Michel MC. Effects of noradrenaline and neuropeptide Y on rat mesenteric microvessel contraction. Naunyn-Schmiedeberg's Arch Pharmacol 1996; 353: 314–23.
 - 35 Prieto D, Buus CL, Mulvany MJ, Nilsson H. Interactions between neuropeptide Y and the adenylyl cyclase pathway in rat mesenteric small arteries: role of membrane potential. J Physiol 1997; 502: 281–92.
 - 36 Sjoblom-Widfeldt N, Gustafsson H, Nilsson H. Transmitter characteristics of small mesenteric arteries from the rat. Acta Physiol Scand 1990; 138: 203–12.
 - 37 Kennedy C, Saville VL, Burnstock G. The contributions of noradrenaline and ATP to the responses of the rabbit central ear artery to sympathetic nerve stimulation depend on the parameters of stimulation. Eur J Pharmacol 1986; 122: 291–300.
 - 38 Lamont C, Vainorius E, Wier WG. Purinergic and adrenergic Ca²⁺ transients during neurogenic contractions of rat mesenteric small arteries. J Physiol 2003; 549.3: 801–8.
 - 39 Brain KL, Cuprian AM, Williams DJ, Cunnane TC. The sources and sequestration of Ca²⁺ contributing to neuroeffector Ca²⁺ transients in the mouse vas deferens. J Physiol 2003; 553 (Pt 2): 627–35.
 - 40 Lamont C, Wier WG. Evoked and spontaneous purinergic junctional Ca²⁺ transients (jCaTs) in rat small arteries. Circ Res 2002; 91: 454–6.
 - 41 Jaggar JH, Porter VA, Lederer MR, Nelson MT. Calcium sparks in smooth muscle. Am J Physiol 2000; 278: C235–56.
 - 42 Lamont C, Vial C, Evans RJ, Wier GW. P2X₁ receptors mediate sympathetic post-junctional Ca²⁺ transients (jCaTs) in mesenteric small arteries. Am J Physiol Heart Circ Physiol 2006; 291 (6): H3106–13.
 - 43 Mulryan K, Gitterman DP, Lewis CJ, Vial C, Leckie BJ, Cobb AL, *et al*. Reduced vas deferens contraction and male fertility in mice lacking P2X₁ receptors. Nature 2000; 403: 86–9.
 - 44 Zang WJ, Balke CW, Wier WG. Graded α_1 -adrenoceptor activation of arteries involves recruitment of smooth muscle cells to produce 'all or none' Ca²⁺ signals. Cell Calcium 2001; 29: 327–34.
 - 45 Nilsson H, Goldstein M, Nilsson O. Adrenergic innervation and neurogenic response in large and small arteries and veins from the rat. Acta Physiol Scand 1986; 126: 121–33.
 - 46 Lamont C, Wier WG. Different roles of ryanodine receptors and inositol (1,4,5)-trisphosphate receptors in adrenergically stimulated contractions of small arteries. Am J Physiol Heart Circ Physiol 2004; 287: H617–25.
 - 47 Mauban JRH, Lamont C, Balke CW, Wier WG. Adrenergic stimulation of rat resistance arteries affects Ca²⁺ sparks, Ca²⁺ waves, and Ca²⁺ oscillations. Am J Physiol 2001; 280: H2399–405.
 - 48 Miriel VA, Mauban JR, Blaustein MP, Wier WG. Local and cellular Ca²⁺ transients in smooth muscle of pressurized rat resistance arteries during myogenic and agonist stimulation. J Physiol 1999; 518 (Pt 3): 815–24.
 - 49 Zacharia J, Zhang J, Wier GW. Calcium signaling in mouse mesenteric small arteries: myogenic tone and adrenergic vasoconstriction. FASEB J 2006; 20: A1174.
 - 50 Mironneau J, Coussin F, Morel JL, Barbot C, Jeyakumar LH, Fleischer S, *et al*. Calcium signaling through nucleotide receptor P2X₁ in rat portal vein myocytes. J Physiol 2001; 536: 339–50.
 - 51 Lagaud GJ, Stoclet JC, Andriantsitohaina R. Calcium handling and purinoceptor subtypes involved in ATP-induced contraction in rat small mesenteric arteries. J Physiol 1996; 492 (Pt 3): 689–703.
 - 52 Putney JW. The pharmacology of capacitative calcium entry. Mol Interv 2001; 1: 84–94.
 - 53 Ruehlmann DO, Lee CH, Poburko D, van Breemen C. Asynchronous Ca²⁺ waves in intact venous smooth muscle. Circ Res 2000; 86: E72–9.