

Full-length article

Astragalus polysaccharide reduces hepatic endoplasmic reticulum stress and restores glucose homeostasis in a diabetic KKAY mouse model¹Xian-qing MAO², Yong WU³, Ke WU², Ming LIU², Jing-fang ZHANG⁴, Feng ZOU², Jing-ping OU-YANG^{2,5}

²Department of Pathophysiology, Medical College of Wuhan University, Hubei Provincial Key Laboratory of Allergy and Immune-Related Diseases, Wuhan 430071, China; ³Division of Endocrinology and Diabetes, Department of Medicine, University of Oklahoma Health Science Center, Oklahoma city, Oklahoma 73104, USA; ⁴Department of Pathology and Physiology, Jingmen Vocational College, Jingmen 448000, China

Key words

Astragalus membranaceus; polysaccharide; type 2 diabetes mellitus; insulin resistance; endoplasmic reticulum; stress

¹Project supported by grants from the National Natural Science Foundation of China (No 30370673).

⁵Correspondence to Dr Jing-ping OU-YANG.
Phn 86-27-8733-1146.
Fax 86-27-8733-1077.
E-mail jpyou@163.com

Received 2007-04-26

Accepted 2007-06-14

doi: 10.1111/j.1745-7254.2007.00674.x

Abstract

Aim: To examine the potential effects of *Astragalus* polysaccharide (APS) on hepatic endoplasmic reticulum (ER) stress *in vivo* and *in vitro* and its link with hypoglycemia activity, thus establishing the mechanism underlying the hypoglycemic action of APS. **Methods:** The obese and type 2 diabetic KKAY mouse model, which is the yellow offspring of the KK mice expressed A^y gene (700 mg·kg⁻¹·d⁻¹, 8 weeks) and a high glucose-induced HepG2 cell model (200 µg/mL, 24 h) were treated with APS. The oral glucose tolerance test was measured to reflex insulin sensitivity with the calculated homeostasis model assessment (HOMA-IR) index. XBP1 (*Xho1* site-binding protein 1) transcription and splicing, an indicator of ER stress, was analyzed by RT-PCR and real-time PCR. The expression and activation of glycogen synthase kinase 3 beta (GSK3β), an insulin signaling protein, was measured by Western blotting. **Results:** APS can alleviate ER stress in cultured cells *in vivo*. The hyperglycemia status, systemic insulin sensitivity, fatty liver disease, and insulin action in the liver of diabetic mice were partly normalized or improved in response to APS administration. **Conclusion:** Our results indicate that APS enables insulin-sensitizing and hypoglycemic activity at least in part by enhancing the adaptive capacity of the ER, which can further promote insulin signal transduction. Thus, APS has promising application in the treatment of type 2 diabetes.

Introduction

Type 2 diabetes mellitus (T2DM) is one of the toughest cases of human diseases worldwide. The number of T2DM patients is expected to affect 300 million people by the year 2025^[1] due to an increased number of elderly people, a greater prevalence of obesity, and sedentary lifestyles. Therefore, it is very important to reinforce the effective prevention and cure of this disease. Traditional Chinese medicines and their extractions demonstrate the characteristics of economy and effectiveness to cure diabetes and its complications. In traditional Chinese medicine, *Astragalus* is commonly found in mixtures with other herbs, and is used in the treatment of

numerous ailments, including heart, liver, and kidney diseases, as well as cancer, viral infections, and immune system disorders. Western herbalists began using *Astragalus* in the 1800s as an ingredient in various tonics. The use of *Astragalus* became popular in the 1980s based on theories about anticancer properties, although these proposed effects have not been clearly demonstrated in reliable human studies. *Astragalus* polysaccharides (APS), the polysaccharide component of the ethanol extract of *Astragalus* roots is an important active component of *Astragalus*. Apart from the actions of the antioxidant, antihypertensive, and immunomodulatory activities^[2-4] of APS, we found that it also shows insulin-sensitizing and

hypoglycemic activity by decreasing the elevated expression and activity of protein-tyrosine phosphatase 1B (PTP1B) in the skeletal muscles of T2DM rats in our previous research^[5].

Among the 3 major insulin-responsive tissues (fat, muscle, and liver), liver plays a central role in the control of glucose homeostasis and is subject to complex regulation by insulin and other hormones^[6]. Many studies have proved that the impaired regulation of hepatic glucose production is a characteristic feature of the metabolic syndrome, which is also known as “insulin resistance syndrome”^[7]. It has recently been discovered that endoplasmic reticulum (ER) stress maybe a key link between obesity, insulin resistance, and T2DM^[8]. Hepatocytes have a well-developed ER structure and ER stress is involved in the development of hepatic insulin resistance^[9]. In light of the important role of the liver in glucose homeostasis and the pathogenesis of diabetes, we sought to examine the potential effect of APS on the insulin signal and ER stress response signal in hepatocytes of diabetic animals and a high glucose-treated cell model.

In this study, we provide evidence that the mechanistic link mentioned earlier can be exploited for therapeutic purposes with the traditional oral Chinese herb APS, which alleviates ER stress in high glucose-treated HepG2 cells and a diabetic animal model. The treatment of obese and diabetic KKAY mice, which is the yellow offspring of the KK mice, expressed *A^y* gene, with APS resulted in the normalization of hyperglycemia, restoration of systemic insulin sensitivity, resolution of fatty liver disease, and the enhancement of insulin action in the liver. Our results demonstrated that APS can improve insulin sensitivity coupled with the enhanced adaptive capacity of the ER and acts as potent antidiabetic modalities with potential application in the treatment of type 2 diabetes.

Materials and methods

Plant materials and preparation of APS *Astragalus membranaceus* (Fisch) and *Bunge var mongholicus* (Bunge) Hsiao were purchased from Shanghai Medicinal Materials (Shanghai, China) and identified by the Department of Authentication of Chinese Medicine, Hubei College of Chinese Traditional Medicine (Wuhan, China). We used the representative specimen which had been kept in our laboratory by anterior researchers. In brief, APS was extracted with optimized techniques using direct water decoction, as described previously^[10]. Three subtypes of APS are defined by phytochemical screening: APSI, II, and III (1.47:1.21:1). APSI consists of *d*-glucose, *d*-galactose, and *l*-arabinose

in molar ratios of 1.75:1.63:1 and has an average molecular weight of 36 300 kDa. Both APSII and APSIII are dextrans, the linkage mode of which is mainly α -(1 \rightarrow 4) linkages, and in which α -(1 \rightarrow 6) linkages are exiguous. APS is a hazel-colored and water-soluble powder. It was diluted to 12% in normal saline before use.

Biochemical reagents GAPDH (ab9845), glycogen synthase kinase 3 beta (GSK3 β , 9332), p (ser9)-GSK3 β (9336), and p (ser641)-GS (3891) were purchased from Abcam (Abcam, Cambridge, UK) and Cell Signaling Technology (Danvers, MA, USA). DMSO (D098-100) and tunicamycin (Tun, T-7765) were from Sigma (St Louis, MO, USA). Dulbecco's modified Eagle's medium (1 \times), liquid (no glucose) (11966025) without glucose and sodium pyruvate containing *L*-glutamine were from Invitrogen-Gibco (Frederick, MD, USA). Enhanced chemiluminescence (ECL) was performed by using the protein detector Lumi-GLO western blot kit from Kierkegaard and Perry Laboratories (Gaithersburg, MA, USA). The BCA (bicincho-ninic acid) protein assay kit was from Pierce Biotechnology (Rockford, IL, USA). The RevertAid first strand cDNA synthesis kit (K1622) and reagents for PCR were from Fermentas (Glen Burnie, MD, USA). The EZNA gel extraction kit was from Omega (Peqlab, Erlangen, Germany). The DyNAmo SYBR green qPCR kit was from Finnzymes (Ipswich, MA, USA). All other chemicals and reagents were of analytical grade.

Mouse models and administration of APS Female KKAY and C57BL/6J mice from the age of 8 weeks were obtained from the Chinese Academy of Medical Sciences (Beijing, China) and housed individually in plastic cages at 20 °C, with lighting on from 6:00–18:00. The C57BL/6J mice were fed a normal chow diet consisting (as a percentage of total kcal) of 12% fat, 60% carbohydrates, and 28% protein. The KKAY mice were fed a high-fat diet consisting of 41% fat, 41% carbohydrates, and 18% protein. The KKAY mice were a cross between glucose-intolerant black KK female mice and male, yellow, obese Ay mice and are known to serve as excellent models of T2DM, while C57BL/6J mice with normal diets are generally used as non-diabetic controls^[11–14]. All experimental procedures were approved and carried out in compliance with the guidelines of the Wuhan University School of Medicine Committee on Animals. From 12 weeks of age the animals were given APS (700 mg·kg⁻¹·d⁻¹) orally or vehicle treatment (PBS, phosphate-buffered saline) for 2 months at the same time. Plasma glucose, insulin, glycogen^[15], free fatty acid (FFA), and triglyceride^[16] were measured with commercial kits. Before necropsy, saline- and APS-treated animals were intraperitoneally administered insulin (10 U/kg body wt) in saline

or vehicle (saline) after overnight fasting. After 10 min, the mice were killed and the liver was rapidly excised and snap frozen in liquid nitrogen.

Insulin sensitivity (oral glucose tolerance test and the homeostasis model assessment, HOMA-IR) Insulin sensitivity was identified by the comprehensive analysis of the oral glucose tolerance test (OGTT) and calculated HOMA-IR (HOMA-IR index = FPG (Fasting plasma glucose) [mmol/L] × FINS (fasting insulin) [μU/mL] / 22.5)^[17]. The OGTT was performed after a 16 h overnight fast. Glucose (2 g/kg) was administered orally and blood was collected from the orbital sinus at 0, 30, 60, and 120 min, respectively^[18]. The area under the curve (AUC) was calculated for glucose during the OGTT (AUC = 0.5 × [Bg0 + Bg30] / 2 + 0.5 × [Bg30 + Bg60] / 2 + 1 × [Bg60 + Bg120] / 2).

Liver histology The liver samples were embedded in paraffin, and sections were cut into 6 mm slices. Neutral lipid was stained with Sudan III on frozen sections. The hepatocyte ultrastructure was presented by transmission electron microscope (TEM). The total operative procedures complied with the standard protocols.

Cell culture and pretreatment with APS The human hepatocarcinoma cell line HepG2 cells were cultured in minimal essential medium containing 10% fetal bovine serum and 0.5 mg/mL geneticin, supplemented with 100 U/mL penicillin, 100 μg/mL streptomycin, and 2 mmol/L *L*-glutamine in a humidified atmosphere with 5% CO₂ at 37 °C. The cells were plated at a density of 3 × 10⁴ cells/cm² and maintained in culture medium for 24 h before treatment. The HepG2 cells were grown on coverslips and treated with tunicamycin (Tm, 10 μg/mL)^[19] for 16 h as positive ER stress cell control and treated with high glucose culture medium (30 and 45 mmol/L, respectively) for 5 h to induce ER stress response *in vitro*. The HepG2 cells were pretreated with APS (200 μg/mL) for 24 h.

Analysis of XBP1 (XhoI site-binding protein 1) mRNA transcription and splicing by real-time RT-PCR RNA was extracted from the liver samples and HepG2 cells using Trizol reagent. Gene transcription was analyzed by semiquantitative RT-PCR and real-time PCR. The total RNA concentration and purity were determined by measuring the OD₂₆₀ and OD₂₆₀/OD₂₈₀ ratio. The specific primers for the mouse and human samples are as follows: mXBP1 mRNA, sense: 5'-AAACAGAGTAGCAGCGCAGACTGC-3' and antisense: 5'-TCCTTCTGGGTAGACCTCTGGGAG-3'; mβ-actin mRNA, sense: 5'-TCATCACTATTGGCAACGAGC-3' and antisense: 5'-AACAGTCCGCTAGAAGCAC-3'; hXBP1 mRNA, sense: 5'-AAACAGAGTAGCAGCTCAGACTGC-3' and antisense: 5'-TCCTTCTGGGTAGACCTCTGGGAG-3'; and hβ-actin

mRNA, sense: 5'-CAGGGCGTGATGGTGGGCA-3' and antisense: 5'-CAAACATCATCTGGGTCATCTTCTC-3'. In brief, 1 μg total RNA was used to prepare cDNA. Real-time PCR reaction was performed by the following thermal cycling conditions: 94 °C for 4 min, 94 °C for 10 s, 65 °C for 30 s, repeated for 40 cycles, and 72 °C for 30 s. Digestion with *Pst*I (which cuts only in the unspliced cDNA) was used to distinguish the unspliced from the spliced bands and then we ran a carefully prepared 2% gel that would present the 2 PCR products clearly. This protocol works for the human and mouse genes. The quantity of specific mRNA was normalized as a ratio to the amount of β-actin mRNA.

Analysis of protein expression and phosphorylation by Western blotting The cell lysates from tissues or cells were prepared in 1 mL lysis buffer (20 mmol/l Tris, pH 7.5, 5 mmol/L EDTA, 10 mmol/L Na₄P₂O₇, 100 mmol/L NaF, 2 mmol/L Na₃VO₄, 1% Nonidet P-40, 1 mmol/L phenyl-methylsulfonyl fluoride, and 10 μg/mL aprotinin) on ice in 1.5 mL microtubes. The lysates were solubilized by continuous stirring for 1 h at 4 °C and centrifuged for 10–15 min at 14000 × g. The supernatants were collected and protein concentrations were measured with BCA protein assay reagent and then stored at -80 °C until further analysis. The cell lysates were subjected to SDS-PAGE and blotted onto a polyvinylidene difluoride membrane followed by incubation with the primary antibodies anti-GSK3β, anti-ser9GSK3β, and anti-ser641GS. The proteins were detected with an ECL system.

Statistical analysis All values are expressed as mean ± SEM. Statistical significance was determined using ANOVA followed by Turkey's test. *P* < 0.05 was considered statistically significant.

Results

Effect of APS treatment on systematic glucose metabolism and insulin sensitivity in KKAY mice

Characteristics of experimental animals To investigate the *in vivo* effects of APS, we employed obese and diabetic (KKAY) mice, a model of severe obesity and insulin resistance. The glucose levels of both fasting and fed mice were significantly upregulated in KKAY mice (T2DM) than those in normal C57BL/6J mice (control) (1.6-fold, *P* < 0.05; 3.6-fold, *P* < 0.05, respectively; Figure 1), which were significantly reduced after treatment with APS (700 mg·kg⁻¹·d⁻¹, *po*) for 8 weeks. Notably, the insulin concentrations in KKAY mice were higher than the values of the control mice (6-fold, *P* < 0.05), but treatment with APS did not affect the insulin levels in both the control and type 2 diabetic mice, suggesting that the action of APS may not be




Figure 1. Characteristics of the experimental animals. Data are expressed as mean±SEM. ^b*P*<0.05 vs C57BL/6J mice (the same age); ^c*P*<0.05 vs C group mice; ^e*P*<0.05 vs KA group mice. *n* indicates the number of animals studied, which is presented in the parenthesis of the figure.

mediated by changing the insulin levels. The KKAY diabetic mice weighed 20 g more than the normal chow-fed C57BL/6J mice (20.4±0.4 vs 40.3±0.6, *P*<0.05). APS treatment could significantly inhibit body weight gain in diabetic mice (KA: 41.0±0.9 vs KK: 45.6±1.4, *P*<0.05) and had no effect on that of the control mice (CA: 20.9±0.1; C: 21.1±0.1, *P*>0.05; Figures 2, 3). All data are expressed as mean±SEM calculated from the results of 8–10 mice

Insulin sensitivity The KKAY mice did not show significant reductions in insulin levels after APS treatment (KA mouse group; Figure 1). To confirm whether APS improved glucose intolerance in KKAY mice, we performed a OGTT and observed that the mice treated with APS showed a lower peak of plasma glucose concentration at 30 min after the




Figure 2. Comparison of appearance in KKAY mice with *Astragalus* polysaccharide therapy.




Figure 3. Growth curve of body weight in experimental animals. Data are expressed as mean±SEM. ^b*P*<0.05 vs C group; ^e*P*<0.05 vs KA group.

glucose load. The plasma glucose level declined more rapidly compared with that in the vehicle-treated KKAY mice (Figure 4). Accordingly, after APS therapy, the HOMA-IR index in the KA mouse group was significantly lower than that in the KK mouse group (10.0±0.51 vs 15.5±0.52, *P*<0.05). These results suggest that the blood glucose-lowering effect of APS is due to increased systemic insulin sensitivity. In addition, neither of these parameters, blood glucose and insulin levels, were different the between APS-treated (C group mice) and vehicle-treated lean control C57BL/6J mice (CA group mice). Taken together, the impaired insulin sensitivity in the T2DM rats was improved following APS treatment.

Liver pathology and biochemistry Obesity and diabetes in mice and humans is associated with alterations in liver lipid metabolism and fatty liver disease^[19–21]. APS treatment resulted in the resolution of the obesity-induced lipid accumulation and glycogen synthesis in the liver from KKAY mice (Figure 5). There was a significant reduction in liver triglyceride and FFA content in the APS-treated KKAY mice compared to the control animals (*P*<0.05; Figure




Figure 4. Effect of APS on OGTT in experimental animals. Data are expressed as mean±SEM. ^b*P*<0.05 vs C57BL/6J mice; ^c*P*<0.05 vs C group mice; ⁱ*P*<0.05 vs KA group mice. *n* indicates the number of animals studied, which is presented in the parenthesis of the figure. APS, *Astragalus polysaccharide*.




Figure 5. Biochemical parameters of experimental animals. FFA indicates free fatty acid. Data are expressed as mean±SEM. ^b*P*<0.05 vs C group; ^c*P*<0.05 vs KA group. *n* indicates the number of animals studied, which is presented in the parenthesis of the figure.

5). Consistent with this, liver Sudan III staining also showed a significant alleviation of fatty degeneration in APS-treated KKAY mice (Figure 6). Additionally, the dilated ER of hepatocytes was observed by TEM, which can be ameliorated with APS therapy (Figure 7).

Effect of APS treatment on hepatic insulin signal transduction in KKAY mice In an attempt to understand the ameliorating effect of APS on insulin signal transduction in the liver tissue, we next examined whether APS affected the expression and activity of hepatic GSK3β in the obese and diabetic KKAY mouse model. APS treatment significantly reduced GSK3β protein levels in KKAY mice (*P*<0.05, Figure 8). Importantly, treatment with APS significantly induced

GSK3β phosphorylation at serine 9, which is the inactivated form of this kinase (*P*<0.05, Figure 8). Hepatic glycogen synthase (GS), which is a key enzyme in the regulation of glycogen synthesis, is regulated by the phosphorylation of the sites between Ser₆₄₁ and Ser₆₅₃ targeted by GSK3^[22]. To support this, GSK3 inhibitors stimulate hepatic glycogen synthase. Thus, we further examined the effect of APS on the insulin-induced Ser₆₄₁ phosphorylation of GS. APS treatment significantly reduced GS phosphorylation at the Ser₆₄₁ site (*P*<0.05, Figure 9). These results indicate that APS has a positive effect on hepatic insulin transduction.

Effect of APS treatment on markers of ER stress in KKAY mice As metabolic demands increase, for example, in the




Figure 6. Sudan III staining of the liver in experimental animals. Effect of APS on fatty liver in KKAY mice was presented by Sudan III staining. A shows C group, B shows KK group, C shows CA group, and D shows KA group. Original magnifications of histological sections are $\times 400$. Nuclei are in blue, and neutral fat in salmon pink. Figure shows typical hepatic steatosis in KKAY mice at 20 weeks of age.




Figure 7. Effect of APS on hepatocyte ultrastructure alterations. Figure shows dilated ER of hepatocytes (white arrow) by TEM. Original magnifications of histological sections are $\times 17\,000$. KK group is shown with white arrow on the left and KA group is presented on the right.

state of hyperglycemia, ER becomes over loaded, which can perturb the protein folding in this protein factory. The distressed ER may further contribute to impaired insulin action in obesity and ER stress leads to the development of insulin resistance and eventually type 2 diabetes^[8]. To confirm whether APS can act as an agent to enhance the adaptive capacity of the ER, XBP1 transcription and splicing in the liver of the KK group mice was detected with a significant increase compared with the controls ($P < 0.05$), indicating that high ER stress exists in this diabetic animal model (Figure 10). Notably, APS administration could reduce the level of XBP1 ($P < 0.05$) in KKAY mice. Spliced XBP1 levels in the liver from the KK group were also significantly increased compared with those of the KA group ($P < 0.05$). XBP1 transcription and splicing in the liver of normal control mice was not affected by APS treatment (Figure 10A, 10D).

Effect of APS treatment on the high glucose-treated HepG2 cell model The transcription factor XBP1 is a basic motif-leucine zipper protein. The spliced or processed form of XBP1 (XBP1s) is a key factor in ER stress through the transcriptional regulation of an array of genes, including molecular chaperones^[23–26]. The modulation of XBP1s in cells could alter insulin action via its potential impact on the magnitude of the ER stress responses^[27]. Hyperglycemia is a direct cause of insulin resistance, which is the hallmark of T2DM. We next examined whether ER stress is increased in high glucose conditions and whether APS can perform this effect on ER stress *in vitro* by measuring XBP1 transcription and splicing in the high glucose-treated cell model. High glucose (30 mmol/L) is suitable to induce significant ER stress. We observed that APS inhibited high glucose-induced ER stress responses in cultured HepG2 cells (Figure




Figure 8. Effect of APS on the expression and phosphorylation of hepatic GSK3β in KKAy and C57BL/6J mice. Data are expressed as mean±SEM of four (for both of KK and KA group) or five (for both of C and CA group) independent experiments from different animals. ^b*P*<0.05 vs C group; ^e*P*<0.05 vs KA group.




Figure 9. Effect of APS on insulin-induced Ser₆₄₁ phosphorylation of hepatic GS in experimental animals. Data are expressed as mean±SEM of 4 for both of KK and KA group or 5 for both of C and CA group in independent experiments from different animals. ^b*P*<0.05 vs C group; ^e*P*<0.05 vs KA group.

10), while it had no significant effect on Tm-induced ER stress. These may indicate that APS is not an antagonist of tunicamycin, and APS inhibition of ER stress is likely by modification of metabolic disturbance and maintaining glucose homeostasis.

Discussion

T2DM is one of the most prevalent and serious metabolic diseases in the world. Hyperglycemia can directly cause insulin resistance, which is associated with an imbalance

between endocrine pancreatic function and hepatic and extrahepatic insulin sensitivity^[28]. Among the 3 major insulin-responsive tissues (fat, muscle, and liver), the liver plays a central role in the control of glucose homeostasis; insulin signaling in liver is critical in maintaining normal hepatic function^[6,7]. Many studies have proved that the impaired regulation of hepatic glucose production is a characteristic feature of the metabolic syndrome^[29], which is also known as “insulin resistance syndrome”, including obesity, insulin resistance, type 2 diabetes, and other metabolic disorders^[20].

The ER is a membranous network that provides a specialized environment for processing and folding newly synthesized proteins. As metabolic demands increase, which can perturb the protein folding in the ER, so does the workload of this protein factory, collectively called ER stress^[19]. Since hepatocytes have a well-developed ER structure, ER stress is involved in liver-related diseases^[9]. Sustained ER stress, which appears to occur as a result of obesity and diabetes, modulates insulin action in the liver^[8]. The development of hepatocellular ER stress as a result of diabetes/obesity appears to be a major contributor to insulin resistance^[9]. Inhibiting ER stress in the liver or increasing hepatic sensitivity to insulin might break the vicious circle linking hyperinsulinemia and insulin resistance that leads to elevated triglyceride and FFA concentrations, progressive steatosis, ultrastructural mitochondrial lesions in the hepatocytes, and ultimately hepatocyte death. Therefore, we wonder whether the beneficial function of APS on the insulin signal pathway is associated with suppressing ER stress in the liver.




Figure 10. Effect of APS on indicators of ER stress in the liver of KKAY mice and HepG2 cells. Analysis of spliced XBP1 mRNA by RT-PCR and subsequent PstI treatment in the liver of KKAY mice (A); high glucose-treated HepG2 cells (B); high glucose-treated HepG2 cells with APS (200 $\mu\text{g/mL}$) pre-treated for 24 h (C). Quantitative analysis for transcription of XBP1 mRNA by real-time PCR (D). Data are expressed as mean \pm SEM of three independent experiments. ^b $P < 0.05$ vs C group; ^e $P < 0.05$ vs KA group; ^h $P < 0.05$ vs G45 group; ^k $P < 0.05$ vs GA group.

In this study, we adopted KKAY mice as a model of T2DM. KKAY mice show hyperglycemia, hyperlipidemia, and hyperinsulinemia compared with C57BL/6J mice. Consistent with our previous study^[5], we prove that APS has significant hypoglycemic activity and insulin-sensitizing effects in the present study. The diabetic mouse model was significantly obesity-resistant and showed alleviated hepatic fatty degeneration in response to APS therapy. As the liver is the hinge of nutritive material metabolism, hepatic insulin resis-

tance has been suggested to be a later factor in the development of hyperglycemia. Increased hepatic glucose production is tightly correlated with fasting hyperglycemia in type 2 diabetics^[30]. The presence of fatty liver in T2DM and obese patients, which is also known as non-alcohol fatty liver, has long been reported. Obesity-related insulin resistance might be partly responsible for liver fat deposition^[21], so it is important for us to detect whether APS can ameliorate hepatic insulin sensitivity and its mechanism.

To further analyze the action of APS on hepatic glucose metabolism and insulin action, we observed the expression and activity of hepatic GSK3 β , one of the important negative regulators of insulin signal transduction in the liver. Previous studies suggested that the inhibition of GSK3 (ser9 for β subunit) in animal models of diabetes leads to the normalization of blood glucose levels and improved hepatic and peripheral insulin resistance, while high GSK3 activity has been reported in T2DM, which is also involved in diminished levels of the IPF1/PDX1 (islet transcription factor 1, also known as IPF-1, IDX-1, and STF-1) protein and β cell dysfunction during the progression of diabetes^[31,32]. Our findings indicate that the hypoglycemic activity of APS is mediated by insulin sensitivity improvement at least partly related to GSK3 inhibition. Further-more, it has been recently reported that GSK3 may play a central role in signaling the downstream effects of ER stress^[33,34]. Since ER stress-induced lipid accumulation and cell death play a role in the pathogenesis of disorders, diabetes mellitus, and hepatic steatosis^[34,35], one can speculate that the protection against ER stress-induced cellular dysfunction occurs while GSK3 activity is inhibited. Thus, GSK3 inhibition undoubtedly leads to insulin sensitivity improvement which will promote a beneficial cycle coupled with the enhancement of ER function to cope with metabolic alterations.

To determine how APS performs its function on the management of metabolic abnormalities associated with obesity and diabetes, in the present study we hypothesize that the reversal of hyperglycemia, increases glucose tolerance and insulin sensitivity induced by APS is related to a decrease in ER stress so APS-treated KKAY mice should display a reduction in ER stress. Spliced *XBP1* mRNA induced by activated IRE1 (Inositol-Requiring Enzyme 1) is translated to the protein, a potent transcription factor that induces BiP expression^[21,25]. *XBP1* is also induced by activated ATF6 (Activating Transcription Factor 6)^[23]. It is thus thought to be an important marker reflecting both IRE1 and ATF6 signaling in response to ER stress^[24,36]. So it is important to distinguish between the spliced and non-spliced form of *XBP1* mRNA for the quantitative measurement of *XBP1* gene expression^[37,38]. Our data indicated that in APS-treated KKAY mice, the transcription and splicing of *XBP1* in the liver was markedly reduced in comparison with the controls. Similar to these results, high glucose (30 mmol/L) was used to induce ER stress in a cultured cell model, and increased *XBP1* gene expression and splicing was significantly suppressed in HepG2 cells pretreated with APS. Therefore, APS has a role in inhibiting hepatic ER stress in the state of hyperglycemia. We concluded that APS promotes insulin

signal transduction, thus it enables insulin-sensitizing and hypoglycemic activity, which is related to the enhanced adaptive capacity of the ER. The alleviation of ER stress also contributes to insulin signaling. This positive interaction shows that APS has a promising application in the treatment of type 2 diabetes.

Although our results in this study show a remarkable pharmacological effect on hepatic insulin resistance, we cannot directly extrapolate our results to humans. Hence, it is necessary to reveal a more detailed mechanism of APS in the improvement of insulin resistance in humans in our future research. In conclusion, ER stress is a key link between obesity, insulin resistance, and type 2 diabetes. Our study provides new evidence that APS renders its hypoglycemic action through decreasing liver insulin resistance coupled with alleviating ER stress. In addition, the treatment of obese and diabetic mice with APS resulted in a significant alleviation of hyperglycemia, restoration of systemic insulin sensitivity, resolution of fatty liver disease, and enhancement of insulin action in liver tissue. Our research demonstrates that APS can enhance the adaptive capacity of the ER and act as potent antidiabetic modalities with promising application in the treatment of type 2 diabetes.

Acknowledgment

We appreciate the help given by the Experimental Animal Center of Wuhan University (No SCXK 2003-0003) and other cooperative partners.

References

- 1 Kiberstis PA. A surfeit of suspects (special section—Type 2 diabetes). *Science* 2005; 307: 3692
- 2 Grover JK, Yadav S, Vats V. Medicinal plants of India with anti-diabetic potential. *J Ethnopharmacol* 2002; 81: 81–100.
- 3 Mao CP, Xie ML, Gu ZL. Effects of konjac extract on insulin sensitivity in high fat diet rats. *Acta Pharmacol Sin* 2002; 23: 855–9.
- 4 Wu F, Chen X. A review of pharmacological study on *Astragalus membranaceus* (Fisch) Bge. *Zhong Yao Cai* 2004; 27: 232–4.
- 5 Wu Y, Ou-Yang JP, Wu K, Wang Y, Zhou YF, Wen CY. Hypoglycemic effect of *Astragalus polysaccharide* and its effect on PTP1B. *Acta Pharmacol Sin* 2005; 26: 345–52.
- 6 Michael MD, Kulkarni RN, Postic C, Previs SF, Shulman GI, Magnuson MA, *et al*. Loss of insulin signaling in hepatocytes leads to severe insulin resistance and progressive hepatic dysfunction. *Mol Cell* 2000; 6: 87–97.
- 7 Klover PJ, Mooney RA. Hepatocytes: critical for glucose homeostasis. *Int J Biochem Cell Biol* 2004; 36: 753–8.
- 8 Ozcan U, Cao Q, Yilmaz E, Lee AH, Iwakoshi NN, Ozdelen E, *et al*. Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes. *Science* 2004; 306: 457–61.

- 9 Ji C, Kaplowitz N. ER stress: can the liver cope? *J Hepatol* 2006; 45: 321–33.
- 10 Ni Y, Su Q, Liu X, Li XR. Experimental study of optimized techniques of water decoction extraction of *Astragalus* polysaccharide. *Zhongguo Zhong Yao Za Zhi* 1998; 23: 284–6.
- 11 Suto J, Matsuura S, Imamura K, Yamanaka H, Sekikawa K. Genetic analysis of non-insulin-dependent diabetes mellitus in KK and KK-Ay mice. *Eur J Endocrinol* 1998; 139: 654–61.
- 12 Iwatsuka H, Shino A, Suzuoki Z. General survey of diabetic features of yellow KK mice. *Endocrinol Jpn* 1970; 17: 23–35.
- 13 Butler L. The inheritance of glucosuria in the KK and AY mouse. *Can J Genet Cytol* 1972; 14: 265–9.
- 14 JAX mice data sheet: strain details of C57BL/6J. Available from URL: <http://jaxmice.jax.org/strain/000664.html>
- 15 Yang R, Cao L, Gasa R, Brady MJ, Sherry AD, Newgard CB. Glycogen-targeting subunits and glucokinase differentially affect pathways of glycogen metabolism and their regulation in hepatocytes. *J Biol Chem* 2002; 277: 1514–23.
- 16 Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 1957; 226: 497–509.
- 17 Bonora E, Targher G, Alberiche M, Bonadonna RC, Saggiani F, Zenere MB, *et al*. Homeostasis model assessment closely mirrors the glucose clamp technique in the assessment of insulin sensitivity studies in subjects with various degrees of glucose tolerance and insulin sensitivity. *Diabetes Care* 2000; 23: 57–63.
- 18 Shiuchi T, Cui TX, Wu L, Nakagami H, Takeda-Matsubara Y, Iwai M, *et al*. ACE inhibitor improves insulin resistance in diabetic mouse via bradykinin and NO. *Hypertension* 2002; 40: 329–34.
- 19 Gu F, Nguyen DT, Stuibler M, Dube N, Tremblay ML, Chevet E. Protein-tyrosine phosphatase 1B potentiates IRE1 signaling during endoplasmic reticulum stress. *J Biol Chem* 2004; 279: 49 689–93.
- 20 Berge RK, Tronstad KJ, Berge K, Rost TH, Wergedahl H, Gudbrandsen OA, *et al*. The metabolic syndrome and the hepatic fatty acid drainage hypothesis. *Biochimie* 2005; 87: 15–20.
- 21 Eriksson S, Eriksson KF, Bondesson L. Nonalcoholic steatohepatitis in obesity: a reversible condition. *Acta Med Scand* 1986; 220: 83–8.
- 22 Roach PJ. Glycogen and its metabolism. *Curr Mol Med* 2002; 2: 101–20.
- 23 Calton M, Zeng H, Urano F, Till JH, Hubbard SR, Harding HP, *et al*. IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. *Nature* 2002; 415: 92–6.
- 24 Yoshida H, Matsui T, Yamamoto A, Okada T, Mori K. XBP1 mRNA is induced by ATF6 and spliced by IRE1 in response to ER stress to produce a highly active transcription factor. *Cell* 2001; 107: 881–91.
- 25 Lee AH, Iwakoshi NN, Glimcher LH. XBP-1 regulates a subset of endoplasmic reticulum resident chaperone genes in the unfolded protein response. *Mol Cell Biol* 2003; 23: 7448–59.
- 26 Shen X, Ellis RE, Lee K, Liu CY, Yang K, Solomon A, *et al*. Complementary signaling pathways regulate the unfolded protein response and are required for *C elegans* development. *Cell* 2001; 107: 893–903.
- 27 Ji C, Kaplowitz N. ER stress: can the liver cope? *J Hepatol* 2006; 45: 321–33.
- 28 Bavenholm PN, Pigon J, Ostenson CG, Efendic S. Insulin sensitivity of suppression of endogenous glucose production is the single most important determinant of glucose tolerance. *Diabetes* 2001; 50: 1449–54.
- 29 Marchesini G, Brizi M, Bianchi G, Tomassetti S, Bugianesi E, Lenzi M, *et al*. Nonalcoholic fatty liver disease: a feature of the metabolic syndrome. *Diabetes* 2001; 50: 1844–50.
- 30 DeFronzo RA. Pathogenesis of type 2 diabetes: metabolic and molecular implications for identifying diabetes genes. *Diabetes Rev* 1997; 5: 177–269.
- 31 Jope RS, Johnson GV. The glamour and gloom of glycogen synthase kinase-3. *Trends Biochem Sci* 2004; 29: 95–102.
- 32 Boucher MJ, Selander L, Carlsson L, Edlund H. Phosphorylation marks IPF1/PDX1 protein for degradation by glycogen synthase kinase 3-dependent mechanisms. *J Biol Chem* 2006; 281: 6395–403.
- 33 Song L, De Sarno P, Jope RS. Central role of glycogen synthase kinase-3 beta in endoplasmic reticulum stress-induced caspase-3 activation. *J Biol Chem* 2002; 277: 44 701–8.
- 34 Kim AJ, Shi Y, Austin RC, Werstuck GH. Valproate protects cells from ER stress-induced lipid accumulation and apoptosis by inhibiting glycogen synthase kinase-3. *J Cell Sci* 2005; 118: 89–99.
- 35 Werstuck GH, Lentz SR, Dayal S, Hossain GS, Sood SK, Shi YY, *et al*. Homocysteine-induced endoplasmic reticulum stress causes dysregulation of the cholesterol and triglyceride biosynthetic pathways. *J Clin Invest* 2001; 107: 1263–73.
- 36 Yoshida H, Nadanaka S, Sato R, Mori K. XBP1 is critical to protect cells from endoplasmic reticulum stress: evidence from Site-2 protease-deficient Chinese hamster ovary cells. *Cell Struct Funct* 2006; 31: 117–25.
- 37 Hirota M, Kitagaki M, Itagaki H, Aiba S. Quantitative measurement of spliced *XBP1* mRNA as an indicator of endoplasmic reticulum stress. *J Toxicol Sci* 2006; 31: 149–56.
- 38 Iwakaki T, Akai R. Analysis of the XBP1 splicing mechanism using endoplasmic reticulum stress-indicators. *Biochem Biophys Res Commun* 2006; 350: 709–15.