

Letter to the Editor

Comparable rate of EGFR kinase domain mutation in lung adenocarcinomas from Chinese male and female never-smokers

Yi-hua SUN^{1,2}, Rong FANG¹, Bin GAO¹, Xiang-kun HAN¹, Jun-hua ZHANG², William PAO³, Hai-quan CHEN^{2,*}, Hong-bin JI^{1,*}

¹Laboratory of Molecular Cell Biology, Institute of Biochemistry and Cell Biology, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences, Shanghai 200031, China; ²Thoracic Surgery, Shanghai Cancer Hospital, Fudan University, Shanghai 200032, China; ³Vanderbilt-Ingram Cancer Center, 2220 Pierce Avenue, 777 Preston Research Building, Nashville, TN 37232–6307, USA

Acta Pharmacologica Sinica (2010) 31: 647–648; doi: 10.1038/aps.2010.43; published online 26 April 2010

Lung cancer patients with the epidermal growth factor receptor (EGFR) kinase domain mutations frequently show good responses to small molecule tyrosine kinase inhibitors, including Iressa and Tarceva, in clinical studies^[1–3]. Previous studies have demonstrated that EGFR kinase domain mutations are commonly observed in lung adenocarcinomas, never-smokers, East Asian, and females^[4–8]. In contrast to caucasian females, most East Asian females do not smoke, while tobacco smoking is very common in the male population. This gender-associated tobacco usage may bias the EGFR mutation incidence in lung adenocarcinoma patients. Here we analyzed the EGFR mutation rate together with gender and smoking status in Chinese lung adenocarcinoma patients. Interestingly, our results showed a comparable rate of EGFR kinase domain mutations in lung adenocarcinomas from Chinese male and female never-smokers.

We have sequenced the EGFR kinase domain using cDNA extracted from 224 lung adenocarcinomas continuously collected from September 2007 to October 2009 at Shanghai Cancer Hospital, Fudan University, with patient consent. Clinical information including age, smoking history, stage and lung tumor differentiation status is listed in Table 1. All patients were Chinese; they came from Shanghai and 10 other provinces. About 62% (139/224) of the lung adenocarcinomas were from never-smokers, and about 38% (85/224) were from ever-smokers (Table 1). Only 2 of 110 Chinese females smoked, while 83 of 114 males were ever-smokers. About 63% (142/224) of lung adenocarcinomas harbored EGFR kinase domain mutations. Without considering gender-associated

tobacco usage, our data were similar to those of previous reports^[4,5]. EGFR kinase domain mutations tended to occur in never-smokers (75.54%) more frequently than in ever-smokers (43.53%), and female patients had a higher mutation rate of the EGFR kinase domain (75.45%) than did male patients (51.75%) ($P < 0.01$).

We then analyzed the association of EGFR kinase domain mutations with gender in never-smokers and ever-smokers separately. About 55% (46/83) of male ever-smokers had EGFR mutations (Table 1). The only two female smokers had no EGFR mutations. Interestingly, a comparable EGFR kinase domain mutation rate was found in both male never-smokers (70.97%; 22/31) and female never-smokers (76.85%; 83/108) (Table 1). These data clearly showed that gender did not influence EGFR kinase domain mutation rates in never-smoker patients with lung adenocarcinomas. The low rate of EGFR mutations in males could possibly be ascribed to the low number/percentage of male never-smoker patients with lung adenocarcinomas.

Similar conclusion has been drawn using Japanese clinical samples^[9,10]. Although Chinese and Japanese patients are both from East Asia, it has not been conclusively demonstrated that Japanese lung cancer patients are biologically the same as Chinese lung cancer patients. Our study using 224 lung adenocarcinoma specimens mainly from eastern China extends previous studies^[6,7] and corroborates the notion that gender does not play a significant role in the distribution of EGFR mutations in lung adenocarcinoma. Further studies to include other areas in China and that significantly increase the patient number will be very helpful to solidify and extend our results. The status of EGFR kinase domain mutation has been proposed as one of most important factors used for Iressa or Tarceva treatment decision-making in clinic^[11]. Based on our findings in this study, not only Chinese female never-smokers

* To whom correspondence should be addressed.

E-mail hbjj@sibs.ac.cn (Hong-bin Ji)

hqchen1@yahoo.com (Hai-quan CHEN)

Received 2010-03-03 Accepted 2010-03-16

Table 1. No gender influence on EGFR kinase domain mutation rates in Chinese never-smokers with lung adenocarcinomas.

Lung adenocarcinoma	Total	Male	Female	P value
Patients, No	224	114	110	
Median age, year \pm sd	58.09 \pm 9.90	58.35 \pm 9.78	57.82 \pm 10.06	0.691
Smoking history, No (%)				
Never-smoker	139(62.05%)	31 (27.19%)	108 (98.18%)	<0.001
Ever-smoker	85(37.95%)	83 (72.81%)	2 (1.82%)	
Differentiation, No (%)				
Well	37	13	24	0.001
Moderate	118	53	65	
Poor	69	48	21	
Stage, No (%)				
I	100 (44.64%)	42 (36.84%)	58 (52.73%)	0.102
II	21 (9.38%)	13 (11.40%)	7 (6.36%)	
III	94 (41.96%)	53 (46.49%)	40 (36.36%)	
IV	9 (4.02%)	6 (5.26%)	5 (4.55%)	
EGFR, No (%)				
Wild type	82 (36.61%)	55 (48.25%)	27 (24.55%)	<0.001
Mutated	142 (63.39)	59 (51.75%)	83 (75.45%)	
Never-smoker, No (%)				
EGFR wild type	34 (24.46%)	9 (29.03%)	25 (23.15%)	0.502
EGFR mutation	105 (75.54%)	22 (70.97%)	83 (76.85%)	
Ever-smoker, No (%)				
EGFR wild type	48 (56.47%)	46 (55.42%)	2 (100%)	0.505
EGFR mutation	37 (43.53%)	37 (44.58%)	0 (0%)	

Never-smoker: less than 100 cigarettes in life time.

but also male never-smokers are likely to benefit from EGFR-targeted therapy.

Acknowledgements

This work was supported by the National Basic Research Program of China (No 2010CB912102), the National Natural Science Foundation of China (No 30623003, 30740084, and 30871284), the Chinese Academy of Sciences (No 2008KIP101, KSCX1-YW-22), the Science and Technology Commission of Shanghai Municipality (No 08PJ14105, 09JC1416300) and the Hundred Talents Program of the Chinese Academy of Sciences to Prof Hong-bin Ji.

References

- Lynch TJ, Bell DW, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, Harris PL, *et al.* Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004; 350: 2129–39.
- Paez JG, Janne PA, Lee JC, Tracy S, Greulich H, Gabriel S, Herman P, *et al.* EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 2004; 304: 1497–500.
- Pao W, Miller V, Zakowski M, Doherty J, Politi K, Sarkaria I, Singh B, *et al.* EGF receptor gene mutations are common in lung cancers from "never smokers" and are associated with sensitivity of tumors to gefitinib and erlotinib. *Proc Natl Acad Sci USA* 2004; 101: 13306–11.
- Yang CH. EGFR tyrosine kinase inhibitors for the treatment of NSCLC in East Asia: present and future. *Lung Cancer* 2008; 60 Suppl 2: S23-S30.
- Yatabe Y, Mitsudomi T. Epidermal growth factor receptor mutations in lung cancers. *Pathol Int* 2007; 57: 233–44.
- Shigematsu H, Lin L, Takahashi T, Nomura M, Suzuki M, Wistuba II, Fong KM, *et al.* Clinical and biological features associated with epidermal growth factor receptor gene mutations in lung cancers. *J Natl Cancer Inst* 2005; 97: 339–46.
- Bai H, Mao L, Wang HS, Zhao J, Yang L, An TT, *et al.* Epidermal growth factor receptor mutations in plasma DNA samples predict tumor response in Chinese patients with stages IIIB to IV non-small-cell lung cancer. *J Clin Oncol* 2009; 27: 2653–9.
- Li Q, Zhao YL, Hao HJ, Li XH. EGFR gene mutation status among lung cancer patients in China. *Zhonghua Zhong Liu Za Zhi* 2007; 29: 270–3.
- Kosaka T, Yatabe Y, Endoh H, Kuwano H, Takahashi T, Mitsudomi T. Mutations of the epidermal growth factor receptor gene in lung cancer: biological and clinical implications. *Cancer Res* 2004; 64: 8919–23.
- Tanaka T, Matsuoka M, Sutani A, Gemma A, Maemondo M, Inoue A, Okinaga S, *et al.* Frequency of and variables associated with the EGFR mutation and its subtypes. *Int J Cancer* 126: 651–5.
- Eberhard DA, Giaccone G, Johnson BE. Biomarkers of response to epidermal growth factor receptor inhibitors in Non-Small-Cell Lung Cancer Working Group: standardization for use in the clinical trial setting. *J Clin Oncol* 2008; 26: 983–94.