

## Original Article

# No association between *LRP5* gene polymorphisms and bone and obesity phenotypes in Chinese male-offspring nuclear families

Jin-bo YU, Yao-hua KE, Jin-wei HE, Hao ZHANG, Wei-wei HU, Yun-qiu HU, Miao LI, Yu-juan LIU, Jie-mei GU, Wen-zhen FU, Gao GAO, Hua YUE, Wen-jin XIAO, Zhen-lin ZHANG\*

Department of Osteoporosis and Bone Diseases, Metabolic Bone Disease and Genetics Research Unit, The Shanghai Sixth People's Hospital, Shanghai Jiaotong University, Shanghai 200233, China

**Aim:** To investigate the effect of low-density lipoprotein receptor-related protein 5 (*LRP5*) gene polymorphisms on bone and obesity phenotypes in young Chinese men.

**Methods:** A total of 1244 subjects from 411 Chinese nuclear families were genotyped by using the polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) technique at the Q89R, N740N, and A1330V sites in the *LRP5* gene. Bone mineral density (BMD) in the lumbar spine and the hip, total fat mass and total lean mass were measured using dual-energy X-ray absorptiometry. The association between *LRP5* gene polymorphisms and peak BMD, body mass index (BMI), total fat mass, total lean mass and percentage of fat mass was assessed using a quantitative transmission disequilibrium test (QTDT).

**Results:** No significant within-family associations were found between genotypes or haplotypes of the *LRP5* gene and peak BMD, BMI, total fat mass, total lean mass and percentage of fat mass. The 1000 permutations that were subsequently simulated were in agreement with these within-family association results.

**Conclusion:** Our results suggest that common polymorphic variations of the *LRP5* gene do not influence peak bone mass acquisition and obesity phenotypes in young Chinese men.

**Keywords:** *LRP5*; BMD; fat mass; lean mass; transmission disequilibrium test

Acta Pharmacologica Sinica (2010) 31: 1464–1469; doi: 10.1038/aps.2010.92; published online 18 Oct 2010

## Introduction

Osteoporosis and obesity are two common and complex diseases. Interestingly, the correlation between osteoporosis and obesity has been established both genetically and phenotypically<sup>[1–4]</sup>. In addition, genes associated with osteoporosis may also be candidates for obesity. The low-density lipoprotein receptor-related protein 5 (*LRP5*) functions as a cell membrane co-receptor for *Wnt* and plays an important role in the *Wnt* signaling pathway<sup>[5]</sup>. It is expressed in many tissues, including bone and liver, and it regulates bone and cholesterol metabolism<sup>[5–8]</sup>. From a molecular perspective, the *LRP5*/*Wnt* signaling pathway plays a key role in the switch between osteogenesis and adipogenesis<sup>[9,10]</sup>.

Because *LRP5* is a key regulator of osteoblast proliferation and bone formation, loss-of-function mutations of the *LRP5*

gene have been associated with osteoporosis-pseudoglioma syndrome (OPPG), an autosomal recessive disorder characterized by low bone mass, spontaneous fractures and blindness<sup>[5]</sup>. In contrast, gain-of-function mutations of the *LRP5* gene, such as dominant G171V, cause high bone mass and increased bone biomechanical properties<sup>[11]</sup>. Because of its crucial role in bone development, common variations such as single nucleotide polymorphisms in the *LRP5* gene and their relation to bone phenotypes have been extensively studied. *LRP5* polymorphisms have been shown to determine bone mass variation in the general population<sup>[12–15]</sup>.

*LRP5*, a member of the low-density lipoprotein receptor family, is also important for glucose and cholesterol metabolism<sup>[7,8]</sup>. Several studies have found that *LRP5* polymorphisms are associated with complex diseases or traits that are related to obesity<sup>[16,17]</sup>. However, only one study has reported the relationship between *LRP5* polymorphisms and obesity in Caucasian nuclear families<sup>[18]</sup>.

Therefore, the *LRP5* gene could be a pleiotropic genetic fac-

\* To whom correspondence should be addressed.

E-mail ZYL2002@medmail.com.cn

Received 2010-03-24 Accepted 2010-06-17

tor influencing both osteoporosis and obesity phenotypes. However, a clear relationship between *LRP5*'s single nucleotide polymorphisms (SNPs) and peak BMD and obesity phenotypes has not been elucidated, and studies on men are especially lacking. Here, focusing on the well-characterized polymorphisms of the *LRP5* gene related to osteoporosis (Q89R, N740N, and A1330V), we investigated the relationship of these polymorphisms with peak BMD, BMI, total fat mass, total lean mass and the percentage of fat mass in Chinese male offspring of nuclear families using a quantitative transmission disequilibrium test (QTDT).

## Materials and methods

### Subjects

The study was approved by the Ethics Committee of the Shanghai Jiaotong University affiliated Sixth People's Hospital (Shanghai, China). All the subjects involved in the study were selected from the local population of Shanghai City by the Department of Osteoporosis, and they signed informed consent forms before entering the project. We recruited 427 nuclear families from 2004 to 2007. However, only 411 families were analyzed because the genotypes of 15 individuals could not be determined due to poor DNA quality and one family deviated from Mendelian inheritance. These 411 nuclear families were composed of both parents and at least one healthy male child whose age was largely between 20 and 40 years old. The great majority of the families (400) had one child, and 11 families had two children. All the recruited sons were healthy. The following criteria were used to exclude individuals from the study: (1) serious consequences from a cerebral vascular disease; (2) diabetes mellitus; (3) chronic renal disease; (4) serious chronic liver disease or alcoholism; (5) significant chronic lung disease; (6) corticosteroid therapy at pharmacologic levels for more than 6 months; (7) treatment with anticonvulsant therapy for more than 6 months; (8) evidence of other metabolic or inherited bone disease such as hyperparathyroidism or hypoparathyroidism, Paget's disease of bone, osteomalacia and osteogenesis imperfecta; (9) rheumatoid arthritis or collagen disease; (10) a significant disease of any endocrine organ that would affect bone mass; (11) hyperthyroidism; and (12) any neurological or musculoskeletal condition that would cause a nongenetic low bone mass.

### Phenotype measurements

The BMD ( $\text{g}/\text{cm}^2$ ) of the anteroposterior lumbar spine (L1-4) and the left proximal femur (including the femoral neck, the trochanter and the total hip), the total fat mass and the total lean mass were measured with a dual-energy X-ray absorptiometry densitometer. The coefficient of variation (CV) was obtained from three repeated measurements on 15 individuals. The CV values of the BMD in L1-4, the femoral neck, the trochanter and the total hip were 1.39%, 2.22%, 1.41%, 0.70%, respectively<sup>[19]</sup>. For body composition, the CVs were 3.72% and 1.18% for total fat mass and total lean mass, respectively. Height and body weight were measured using standardized equipment. The BMI was defined as the weight/height<sup>2</sup> in  $\text{kg}/\text{m}^2$ . The percentage of fat mass (PFM) was calculated as the total fat mass divided by weight.

### Genotyping

The DNA was isolated from peripheral blood leukocytes using conventional methods. Polymerase chain reaction (PCR) was performed in the following steps: 95 °C for 5 min and then 32 cycles of 95 °C for 30 s, 65 °C for 30 s, 72 °C for 45 s, and finally 72 °C for 5 min (Okubo et al<sup>[20]</sup>). The PCR primers and the restriction endonucleases used for genotyping are summarized in Table 1. The PCR products were digested with *Ava* II, *Ase* I and *Dra* III restriction endonucleases, respectively. The Q89R genotypes (c.314A>G) were separated by electrophoresis in a 2% agarose gel. The AA genotype produces a 436-bp fragment, the GG genotype produces two fragments of 274 bp and 162 bp, and the heterozygous AG genotype produces three fragments of 436 bp, 274 bp, and 162 bp. The N740N (c.2268T>C) and A1330V (c.4037C>T) genotypes were separated by electrophoresis in a 12% polyacrylamide gel. The N740N CC genotype produces a 237-bp fragment, the TT genotype produces two fragments of 216 bp and 21 bp, and the heterozygous CT genotype produces three fragments of 237 bp, 216 bp, and 21 bp. The A1330V CC genotype produces a 143-bp fragment, the TT genotype produces two fragments of 119 bp and 24 bp, and the heterozygous CT genotype produces three fragments of 143 bp, 119 bp, and 24 bp<sup>[15]</sup>.

### Statistical analysis

A Chi-square test assessed if the genotype frequencies of every

**Table 1.** Information of the analyzed *LRP5* SNPs in this study.

SNP	Location	Amino acid change	PCR primer	Restriction enzyme
Q89R (c.314A>G)	Exon 2	Gln [Q]>Arg [R]	Forward: 5'-TCTGGGCATAGTGCTCCATC-3' Reverse: 5'-TTCGGGATGTGCCATTGAG-3'	<i>Ava</i> II
N740N (c.2268T>C)	Exon 10	Asn [N]>Asn [N]	Forward: 5'-CTACTGGGCCGACACTGGGATTAA-3' Reverse: 5'-ACAGCTCTAATCACTGAGGG-3'	<i>Ase</i> I
A1330V (c.4037C>T)	Exon 18	Ala [A]>Val [V]	Forward: 5'-GACTGTCCAGGACCCGCTCACACG-3' Reverse: 5'-AAGGTTTTTCAGAGCCCCTAC-3'	<i>Dra</i> III

SNP were in Hardy-Weinberg equilibrium. The Shapiro-Wilks test and the Bartlett test were used to assess if the variables of BMD, BMI, fat mass and lean mass had a normal distribution and to test the homogeneity of variances within each of the three SNPs genotypes. Haplotypes were constructed from the three common SNPs in *LRP5* using Phase software version 2.0.2<sup>[21]</sup>. We examined the Lewontin D and linkage disequilibrium (LD) coefficient  $r^2$  between all pairs of biallelic loci using Haploview version 3.2<sup>[22]</sup>. The QTDT was used to measure the population stratification, total association and within-family association between SNPs and BMD and obesity phenotypes. The QTDT software package is available on the Internet (<http://www.sph.umich.edu/csg/abecasis/QTDT/>). As indicated by Abecasis *et al*<sup>[23]</sup>, total association effects are partitioned into between- and within-family components. The between-family component of association is specific to each nuclear family and could be influenced by population stratification. The within-family association tests are immune to the stratification of the population and, therefore, they are significant only in the presence of linkage disequilibrium. Because all of the children in our nuclear families were sons and the effects of the parents' phenotypes were excluded in the QTDT, sex was not used as a covariate to adjust the sons' bone and obesity phenotypic values. Raw BMD values were adjusted by age, height and weight as covariates. BMI, total fat mass, total lean mass and percentage of fat mass values were adjusted by age as a covariate. Because false-positive results might be generated in multiple tests, 1000 simulations were performed to generate empirical  $P$  values<sup>[24]</sup> to assess the reliability of the results. The significance level was set at  $P < 0.05$  for all the analyses. The statistical analyses were conducted using SPSS package version 11.5.

## Results

### The distribution of genotype and haplotype

A total of 1244 individuals from 411 nuclear families comprising 822 parents and 422 offspring were recruited. The basic characteristics of the study subjects are summarized in Table 2. Parental total fat mass and total lean mass were not obtained. All the study subjects were men with an average age of 30.6±6.3 (mean±SD) years. The three SNP genotypes and allele frequencies in the son group are presented in Table 3. All SNPs had a minor allele frequency (MAF) greater than 10% in our population, and the genotype frequencies of these SNPs did not deviate from the Hardy-Weinberg equilibrium. Each pair of biallelic SNPs had intermediate values of LD ( $0.432 < D' < 0.729$ ,  $0.109 < r^2 < 0.510$ ). The most common haplotype was ACC, which had a frequency of 72.3%. The characteristics of the 411 unrelated sons were classified according to the three SNP genotypes and are summarized in Table 4. All the raw phenotypic values of the son group followed a normal distribution. The genotype data of each family were verified for Mendelian inheritance.

### Association between SNPs in the *LRP5* gene and peak BMD

There were 127, 212, and 223 informative nuclear families

**Table 2.** Basic characteristics of parents and son groups (mean±SD).

	Father (n=411)	Mother (n=411)	Son (n=422)
Age (years)	61.1±7.0	58.2±6.3	30.6±6.3
Height (cm)	167.9±5.9	155.8±5.4	173.0±6.0
Weight (kg)	69.9±9.4	58.3±8.1	71.1±11.1
BMI (kg/m <sup>2</sup> )	24.8±3.1	24.0±3.1	23.7±3.4
L1-4 (g/cm <sup>2</sup> )	1.142±0.173	0.995±0.172	1.139±0.137
Femoral neck BMD (g/cm <sup>2</sup> )	0.893±0.131	0.801±0.144	0.998±0.144
Trochanter BMD (g/cm <sup>2</sup> )	0.819±0.141	0.692±0.149	0.825±0.147
Total hip BMD (g/cm <sup>2</sup> )	0.969±0.129	0.872±0.149	1.016±0.139
Total fat mass (kg)	-	-	16.0±7.0
Total lean mass (kg)	-	-	51.4±5.8
PFM (%)	-	-	21.9±7

**Table 3.** The three SNPs genotypes and allele frequencies in unrelated son group (n=411).

Polymorphism	Genotype frequencies			Allele frequencies	
Q89R	AA	AG	GG	A	G
	325 (79.1%)	81 (19.7%)	5 (1.2%)	0.889	0.111
N740N	CC	CT	TT	C	T
	279 (67.9%)	121 (29.4%)	11 (2.7%)	0.826	0.174
A1330V	CC	CT	TT	C	T
	271 (65.9%)	126 (30.7%)	14 (3.4%)	0.813	0.187

for the QTDT analyses of the Q89R, N740N and A1330V sites, respectively, and each of those families had at least one heterozygous parent. The results of the QTDT analysis are presented in Table 5. No significant population stratification was found for BMD at any bone site. Regarding the total association, N740N was significantly associated with BMD at the total hip, and A1330V was significantly associated with BMD at L1-4, the femoral neck and the total hip. However, we did not find significant associations between SNPs in *LRP5* and peak BMD at any bone site for the within-family data. The 1000 permutations were in agreement with these within-family association results for all the tested parameters. Furthermore, we observed an association between the most common haplotype (ACC) and peak BMD using QTDT. There were 272 informative nuclear families for the QTDT analysis at haplotype ACC. None of the tests for population stratification, total association and within-family association between haplotype ACC and peak BMD were significant.

### Association between SNPs in *LRP5* and obesity

The results of the QTDT analysis are presented in Table 6. No population stratification was identified for BMI, total fat mass, total lean mass and percentage of fat mass. Regarding

**Table 4.** Characteristics of the sons (n=422) classified according to the three SNPs genotypes.

	AA	Q89R AG	GG	CC	N740N CT	TT	CC	A1330V CT	TT
n	334	83	5	287	124	11	279	129	14
Age (years)	30.5±5.8	30.9±6.4	32.4±8.8	30.3±5.9	31.4±6.0	28.9±5.3	30.2±5.9	31.4±5.8	30.1±7.0
Height (cm)	173.0±6.1	172.6±5.1	172.4±6.0	172.7±6.0	173.6±5.6	171.5±5.6	172.7±6.1	173.5±5.6	172.9±4.8
Weight (kg)	71.1±11.0	69.5±10.7	70.1±10.3	70.9±11.0	71.0±11.0	66.2±8.5	70.5±10.9	71.7±11.3	68.5±8.7
BMI (kg/m <sup>2</sup> )	23.8±3.4	23.3±3.4	23.8±5.0	23.8±3.4	23.5±3.4	22.5±2.9	23.7±3.4	23.8±3.5	23.0±3.0
L1-4 BMD (g/cm <sup>2</sup> )	1.140±0.138	1.127±0.138	1.173±0.039	1.142±0.137	1.132±0.142	1.090±0.095	1.140±0.136	1.136±0.144	1.112±0.114
Neck BMD (g/cm <sup>2</sup> )	1.002±0.139	0.980±0.161	1.052±0.115	1.004±0.140	0.991±0.154	0.940±0.090	1.006±0.143	0.984±0.149	0.971±0.085
Trochanter BMD (g/cm <sup>2</sup> )	0.823±0.142	0.814±0.152	0.816±0.102	0.826±0.137	0.816±0.162	0.760±0.072	0.824±0.139	0.819±0.160	0.788±0.086
Total hip BMD (g/cm <sup>2</sup> )	1.017±0.134	1.002±0.154	1.066±0.113	1.024±0.137	1.000±0.142	0.960±0.088	1.022±0.138	1.001±0.140	0.984±0.092
Total fat mass (kg)	16.1±7.1	15.3±6.0	13.3±7.4	15.8±6.9	16.4±6.9	13.0±5.9	15.6±6.9	16.6±7.1	15.3±5.7
Total lean mass (kg)	51.5±5.5	50.6±5.9	51.8±8.1	51.4±5.4	51.6±6.0	47.9±3.2	51.2±5.6	51.8±5.8	49.8±4.0
PFM(%)	22.0±7	21.5±7	18.4±8	21.8±7	22.3±7	19.7±7	21.6±7	22.4±7	21.8±6

**Table 5.** P value of tests for population stratification, total association, and within-family association between *LRP5* SNPs and peak BMD. Bold indicates significant P value. BMD values were adjusted by age, height and weight as covariates.

	Q89R	N740N	A1330V
Tests of population stratification			
L1-4 BMD	0.0711	0.4160	0.4438
Femoral neck BMD	0.8255	0.2695	0.7829
Trochanter BMD	0.0590	0.2141	0.4719
Total hip BMD	0.1585	0.2111	0.4345
Tests of total association			
L1-4 BMD	0.2093	0.2629	<b>0.0170</b>
Femoral neck BMD	0.1443	0.2668	<b>0.0227</b>
Trochanter BMD	0.8298	0.1669	0.0566
Total hip BMD	0.3663	<b>0.0471</b>	<b>0.0162</b>
Tests of within-family association			
L1-4 BMD	0.2782	0.8628	0.5468
Femoral neck BMD	0.4148	0.6528	0.3853
Trochanter BMD	0.0656	0.7239	0.6944
Total hip BMD	0.3803	0.9189	0.5943
P 1000 permutation of within-family association			
L1-4 BMD	0.1950	0.8440	0.5270
Femoral neck BMD	0.4350	0.6330	0.3430
Trochanter BMD	0.0750	0.6890	0.6880
Total hip BMD	0.3900	0.9160	0.5630

**Table 6.** P value of tests for population stratification, total association, and within-family association between *LRP5* SNPs and obesity phenotypes. BMI, total fat mass, total lean mass and PFM values were adjusted by age as a covariate.

	Q89R	N740N	A1330V
Tests of population stratification			
BMI	0.4588	0.8774	0.3393
Fat mass	0.6145	0.5621	0.5837
Lean mass	0.2699	0.7653	0.8126
PFM	0.5204	0.7361	0.5028
Tests of total association			
BMI	0.2141	0.3195	0.8180
Fat mass	0.1876	0.8340	0.2456
Lean mass	0.6004	0.6035	0.7569
PFM	0.3762	0.7813	0.5047
Tests of within-family association			
BMI	0.8871	0.5397	0.4746
Fat mass	0.6754	0.7736	0.2276
Lean mass	0.5868	0.5691	0.9509
PFM	0.9643	0.9511	0.3444
P 1000 permutation of within-family association			
BMI	0.8730	0.5470	0.5010
Fat mass	0.7590	0.8350	0.3630
Lean mass	0.6640	0.6420	0.9730
PFM	0.9750	0.9570	0.3770

the obesity phenotypes, no significant total association and within-family association was detected. The subsequent 1000

permutation tests confirmed these negative within-family association results. In addition, we also investigated the



association between the most common haplotype (ACC) and obesity phenotypes using QTDT. No significant population stratification, total association and within-family association were found between haplotype ACC and BMI, total fat mass, total lean mass and percentage of fat mass.

## Discussion

A number of studies have investigated the association between *LRP5* polymorphisms and BMD, revealing inconsistent results. Most of the association studies used the traditional association approach in a random population. The regular association approach may yield spurious results due to population stratification. In addition, the linkage approach is often short on statistical power with the currently used sample sizes. The TDT, a family-based association approach, is immune to population stratification and is more powerful when compared with the traditional linkage approach. The TDT has been proposed by Spielman *et al*<sup>[25]</sup> and, when extended to quantitative traits<sup>[26]</sup>, can be used in nuclear families with or without parental phenotypes.

The *LRP5* cooperates with members of the frizzled family of seven-pass transmembrane receptors to bind *Wnt* proteins and forms a functional ligand-receptor complex that activates the canonical *Wnt*/ $\beta$ -catenin pathway. The *LRP5*/*Wnt* signaling pathway is important for osteoblast differentiation, and mutations in the YWTD/EGF domains of the *LRP5* gene are associated with low bone mass syndromes. Numerous studies have revealed the association of exonic and intronic SNPs of *LRP5* with BMD variation. Most of the studies that test these associations have been performed in old women; few studies have investigated the relationship between *LRP5* gene polymorphisms and peak BMD variation in young men. Koh *et al*<sup>[27]</sup> reported that the Q89R polymorphism was significantly associated with BMD in the femoral neck in 219 young Korean men. After adjusting for age, weight and height, a marginal association was observed in the femoral neck ( $P=0.098$ ). No statistically significant relationship was found between the A1330V polymorphism and BMD in the lumbar spine or the hip. In another study, the polymorphic valine variant at position A1330V of the *LRP5* gene was significantly associated with reduced bone mineral content (BMC) and BMD values in healthy young Finnish men<sup>[28]</sup>. The Gothenburg Osteoporosis and Obesity Determinants (GOOD) Study ( $n=1068$  aged 18–20 years) showed no association between the A1330V genotype and bone phenotypes in men<sup>[14]</sup>. In the present study, after adjusting for the covariates of age, height and weight, we failed to find significant associations between Q89R, N740N, and A1330V in the *LRP5* gene and peak BMD variation in young Chinese men. Our results did not provide evidence to suggest that Q89R, N740N, and A1330V were quantitative trait loci (QTLs) or were in strong disequilibrium with a QTL underlying spine and hip BMD variation in young Chinese men. Our previous study found that Q89R and N740N were significantly associated with BMD in the femoral neck in postmenopausal Chinese women<sup>[15]</sup>. Sex-specific factors might be related to the action of *LRP5* on BMD.

Although the importance of the *LRP5* gene to bone biology is widely acknowledged, its importance to obesity has seldom been reported. Fujino *et al*<sup>[7]</sup> found that *LRP5* was required for cholesterol and glucose metabolism. Guo *et al*<sup>[18]</sup> reported a significant association between SNPs and haplotypes in the *LRP5* gene and human obesity, but the associations observed were found mainly in women. We measured whole-body fat mass, lean mass and BMI as indices of the degree of obesity. Our study did not find a significant association between *LRP5* polymorphisms and obesity in young Chinese men. These results seem to agree with Guo *et al*<sup>[18]</sup>.

Our study has several limitations. We investigated only three common polymorphisms in the *LRP5* gene. We cannot rule out the possibility that an association may exist between other polymorphisms in the *LRP5* gene and bone and obesity phenotypes. Despite the relatively large number of nuclear families (411), we tested only the most common haplotype (ACC) in the haplotype analyses because a limited number of subjects had the other haplotypes.

Our study also has some strengths. First, we selected young men aged between 20 to 40 years because they were expected to have reached their peak BMD. Second, we measured total fat mass and total lean mass as indices of obesity. Third, we investigated the relationship between *LRP5* gene polymorphisms and peak BMD and obesity phenotypes in 411 male offspring of nuclear families using QTDT. Due to our large sample size, the power to test a candidate gene as a QTL was greater than 80%, which can explain about 10% of the variation in BMD and obesity phenotypes.

In summary, we failed to find a significant association between three common polymorphisms in the *LRP5* gene and peak BMD and obesity phenotypes in young Chinese men. Further studies with denser markers and larger sample sizes might shed light on the role of the *LRP5* gene in osteoporosis and obesity.

## Acknowledgements

This study was supported by the National Natural Science Foundation of China (No. 30570891, 30771019, and 30800387) and the Program of Shanghai Chief Scientist (No. 08XD1403000).

## Author contribution

Zhen-lin ZHANG designed research; Jin-bo YU, Hao ZHANG, Wei-wei HU, Yu-juan LIU, Jie-mei GU recruited research subjects; Yun-qiu HU, Miao LI measured BMD, total fat mass and total lean mass; Jin-bo YU, Jin-wei HE, Wen-zhen FU, Yao-hua KE, Gao GAO, Hua YUE, Wen-jin XIAO performed research; Zhen-lin ZHANG contributed new analytic tools; Jin-bo YU, Zhen-lin ZHANG analyzed data; Jin-bo YU wrote the paper.

## References

- 1 Coin A, Sergi G, Benincà P, Lupoli L, Cinti G, Ferrara L, *et al*. Bone mineral density and body composition in underweight and normal elderly subjects. *Osteoporos Int* 2000; 11: 1043–50.
- 2 Tóth E, Ferenc V, Mészáros S, Csupor E, Horváth C. Effects of body

- mass index on bone mineral density in men. *Orv Hetil* 2005; 146: 1489–93.
- 3 Bustamante M, Nogués X, Mellibovsky L, Agueda L, Jurado S, Cáceres E, *et al*. Polymorphisms in the interleukin-6 receptor gene are associated with bone mineral density and body mass index in Spanish postmenopausal women. *Eur J Endocrinol* 2007; 157: 677–84.
  - 4 Li H, Xie H, Liu W, Hu R, Huang B, Tan YF, *et al*. A novel microRNA targeting HDAC5 regulates osteoblast differentiation in mice and contributes to primary osteoporosis in humans. *J Clin Invest* 2009; 119: 3666–77.
  - 5 Gong Y, Slee RB, Fukai N, Rawadi G, Roman-Roman S, Reginato AM, *et al*. LDL receptor-related protein 5 (LRP5) affects bone accrual and eye development. *Cell* 2001; 107: 513–23.
  - 6 Little RD, Carulli JP, Del Mastro RG, Dupuis J, Osborne M, Folz C, *et al*. A mutation in the LDL receptor-related protein 5 gene results in the autosomal dominant high-bone-mass trait. *Am J Hum Genet* 2002; 70: 11–9.
  - 7 Fujino T, Asaba H, Kang MJ, Ikeda Y, Sone H, Takada S, *et al*. Low-density lipoprotein receptor-related protein 5 (LRP5) is essential for normal cholesterol metabolism and glucose-induced insulin secretion. *Proc Natl Acad Sci USA* 2003; 100: 229–34.
  - 8 Jin T. The WNT signaling pathway and diabetes mellitus. *Diabetologia* 2008; 51: 1771–80.
  - 9 Bennett CN, Ross SE, Longo KA, Bajnok L, Hemati N, Johnson KW, *et al*. Regulation of Wnt signaling during adipogenesis. *J Biol Chem* 2002; 277: 30998–1004.
  - 10 Ross SE, Hemati N, Longo KA, Bennett CN, Lucas PC, Erickson RL, *et al*. Inhibition of adipogenesis by Wnt signaling. *Science* 2000; 289: 950–3.
  - 11 Boyden LM, Mao J, Belsky J, Mitzner L, Farhi A, Mitnick MA, *et al*. High bone density due to a mutation in LDL-receptor-related protein 5. *N Engl J Med* 2002; 346: 1513–21.
  - 12 Lee YH, Woo JH, Choi SJ, Ji JD, Song GG. Association between the A1330V polymorphism of the low-density lipoprotein receptor-related protein 5 gene and bone mineral density: a meta-analysis. *Rheumatol Int* 2009; 29: 539–44.
  - 13 Tran BN, Nguyen ND, Eisman JA, Nguyen TV. Association between LRP5 polymorphism and bone mineral density: a Bayesian meta-analysis. *BMC Med Genet* 2008; 9: 55.
  - 14 Grundberg E, Lau EM, Lorentzon M, Karlsson M, Holmberg A, Groop L, *et al*. Large-scale association study between two coding LRP5 gene polymorphisms and bone phenotypes and fractures in men. *Osteoporos Int* 2008; 19: 829–37.
  - 15 Zhang ZL, Qin YJ, He JW, Huang QR, Li M, Hu YQ, *et al*. Association of polymorphisms in low-density lipoprotein receptor-related protein 5 gene with bone mineral density in postmenopausal Chinese women. *Acta Pharmacol Sin* 2005; 26: 1111–6.
  - 16 Figueroa DJ, Hess JF, Ky B, Brown SD, Sandig V, Hermanowski-Vosatka A, *et al*. Expression of the type I diabetes-associated gene LRP5 in macrophages, vitamin A system cells, and the Islets of Langerhans suggests multiple potential roles in diabetes. *J Histochem Cytochem* 2000; 48: 1357–68.
  - 17 Bruford EA, Riise R, Teague PW, Porter K, Thomson KL, Moore AT, *et al*. Linkage mapping in 29 Bardet-Biedl syndrome families confirms loci in chromosomal regions 11q13, 15q22.3-q23, and 16q21. *Genomics* 1997; 41: 93–9.
  - 18 Guo YF, Xiong DH, Shen H, Zhao LJ, Xiao P, Guo Y, *et al*. Polymorphisms of the low-density lipoprotein receptor-related protein 5 (LRP5) gene are associated with obesity phenotypes in a large family-based association study. *J Med Genet* 2006; 43: 798–803.
  - 19 Gao G, Zhang ZL, Zhang H, Hu WW, Huang QR, Lu JH, *et al*. Hip axis length changes in 10554 males and females and the association with femoral neck fracture. *J Clin Densitom* 2008; 11: 360–6.
  - 20 Okubo M, Horinishi A, Kim DH, Yamamoto TT, Murase T. Seven novel sequence variants in the human low density lipoprotein receptor related protein 5 (LRP5) gene. *Hum Mutat* 2002; 19: 186.
  - 21 Stephens M, Smith NJ, Donnelly P. A new statistical method for haplotype reconstruction from population data. *Am J Hum Genet* 2001; 68: 978–89.
  - 22 Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 2005; 21: 263–5.
  - 23 Abecasis GR, Cardon LR, Cookson WO. A general test of association for quantitative traits in nuclear families. *Am J Hum Genet* 2000; 66: 279–92.
  - 24 Zhang ZL, He JW, Qin YJ, Hu YQ, Li M, Zhang H, *et al*. Association between myostatin gene polymorphisms and peak BMD variation in Chinese nuclear families. *Osteoporos Int* 2008; 19: 39–47.
  - 25 Spielman RS, McGinnis RE, Ewens WJ. Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am J Hum Genet* 1993; 52: 506–16.
  - 26 Allison DB. Transmission-disequilibrium tests for quantitative traits. *Am J Hum Genet* 1997; 60: 676–90.
  - 27 Koh JM, Jung MH, Hong JS, Park HJ, Chang JS, Shin HD, *et al*. Association between bone mineral density and LDL receptor-related protein 5 gene polymorphisms in young Korean men. *J Korean Med Sci* 2004; 19: 407–12.
  - 28 Saarinen A, Välimäki VV, Välimäki MJ, Löytyniemi E, Auro K, Uusen P, *et al*. The A1330V polymorphism of the low-density lipoprotein receptor-related protein 5 gene (LRP5) associates with low peak bone mass in young healthy men. *Bone* 2007; 40: 1006–12.