

Full-length article

Role of arachidonic acid in hyposmotic membrane stretch-induced increase in calcium-activated potassium currents in gastric myocytes¹

Meng YANG², Xing-lan LI³, Hui-ying XU³, Jia-bin SUN³, Bin MEI⁴, Hai-feng ZHENG³, Lian-hua PIAO³, De-gang XING³, Zhai-liu LI³, Wen-xie XU^{2,5}

²Department of Physiology, Shanghai Jiaotong University School of Medicine, Shanghai 200030; ³Department of Physiology, Yanbian University College of Medicine, Yanji 133000; ⁴East China Normal University School of Life Science, Shanghai 200062, China

Key words

gastric myocyte; arachidonic acid; hyposmotic membrane stretch; calcium-activated potassium current.

¹Project supported by the National Natural Science Foundation of China (No. 30160028).

⁵Correspondence to Prof Wen-xie XU.

Phn 86-21-5474-7639.

Fax 86-21-6283-2528.

E-mail wenxiexu@sjtu.edu.cn

Received 2005-05-22

Accepted 2005-07-06

doi: 10.1111/j.1745-7254.2005.00201.x

Abstract

Aim: To study effects of arachidonic acid (AA) and its metabolites on the hyposmotic membrane stretch-induced increase in calcium-activated potassium currents ($I_{K(Ca)}$) in gastric myocytes. **Methods:** Membrane currents were recorded by using a conventional whole cell patch-clamp technique in gastric myocytes isolated with collagenase. **Results:** Hyposmotic membrane stretch and AA increased both $I_{K(Ca)}$ and spontaneous transient outward currents significantly. Exogenous AA could potentiate the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$. The hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ was significantly suppressed by dimethyleicosadienoic acid (100 $\mu\text{mol/L}$ in pipette solution), an inhibitor of phospholipase A_2 . Nordihydroguaiaretic acid, a lipoxygenase inhibitor, significantly suppressed AA and hyposmotic membrane stretch-induced increases in $I_{K(Ca)}$. External calcium-free or gadolinium chloride, a blocker of stretch-activated channels, blocked the AA-induced increase in $I_{K(Ca)}$ significantly, but it was not blocked by nifedipine, an L-type calcium channel blocker. Ryanodine, a calcium-induced calcium release agonist, completely blocked the AA-induced increase in $I_{K(Ca)}$; however, heparin, a potent inhibitor of inositol triphosphate receptor, did not block the AA-induced increase in $I_{K(Ca)}$. **Conclusion:** Hyposmotic membrane stretch may activate phospholipase A_2 , which hydrolyzes membrane phospholipids to ultimately produce AA; AA as a second messenger mediates Ca^{2+} influx, which triggers Ca^{2+} -induced Ca^{2+} release and elicits activation of $I_{K(Ca)}$ in gastric antral circular myocytes of the guinea pig.

Introduction

Mechanical stretch is an important physiological stimulus in gut smooth muscles. It is well known that mechanical stretch induces myogenous contraction of gut smooth muscle, but the mechanism underlying this ionic channel process remains unknown. Mechanical stretch regulates the activities of ionic channels, which exist widely in the membranes of various cells and activate many signal transduction pathways. A hypothesis was proposed that membrane stretch induces alterations in the lipid bilayer, which transmits membrane tension to channel proteins or generates lipid-soluble second messengers, such as arachidonic acid (AA)

and other endogenous fatty acids, by membrane-bound phospholipases^[1,2]. Unsaturated fatty acids are major components of membrane lipids and they are mainly released by phospholipase A_2 (PLA₂) activation. AA in the cell membrane is esterified in phospholipids and can be released by PLA₂ in response to various extracellular stimuli^[3,4]. AA and other unsaturated fatty acids modulate the activities of various ion channels and enzymes through direct or indirect pathways. For example: AA potentiates hKir2.3 in part by decreasing inward rectification of the channel^[5]; AA induces membrane depolarization by inhibiting K_{ATP} currents in murine colonic smooth muscle cells^[6]; and AA increases choline acetyltransferase activity in spinal cord neurons, and

this effect is mediated by protein kinase C (PKC)^[7]. In addition, it was shown that AA induces endothelium-dependent hyperpolarization and relaxation of rabbit aorta through activation of apamin-sensitive K⁺ currents^[8]. Abundant evidence has revealed that AA is an important mediator in hyposmotic stress. It was observed that swelling induces AA release via the 85 kDa cell phospholipase A₂ (cPLA₂) in human neuroblastoma cells^[9], and that cell swelling activates PLA₂ in ehrlich ascites tumor cells^[10]. Tinel *et al*^[11] also reported that AA acts as a second messenger for hypotonic-induced calcium transients in rat inner medullary collecting duct (IMCD) cells.

The calcium-activated potassium current ($I_{K(Ca)}$) has been considered to play an important role in excitability and functional regulation in excitable cells. In our previous study, AA and other unsaturated fatty acids directly inhibited calcium currents^[12] and muscarinic currents^[13]. For both AA and hyposmotic membrane stretch-activated $I_{K(Ca)}$ ^[14,15], activation by hyposmotic membrane stretch is associated with calcium-induced calcium release (CICR)^[16], which is triggered by extracellular calcium influx through the stretch-activated channels (SAC) in gastric antral circular myocytes of the guinea pig. However, the roles of AA and other unsaturated fatty acids in the process of $I_{K(Ca)}$ activation by membrane stretch in gastric myocytes remains unclear. In the present study, we therefore investigated the effects of AA and its metabolites on hyposmotic membrane stretch-induced increases in $I_{K(Ca)}$ in gastric antral circular myocytes of guinea pig.

Materials and methods

Preparation of cells EWG/B guinea pigs (obtained from the Experimental Animal Department of Norman Bethune University, Changchun, China; Certificate No 10-6004) of either sex, weighing 250–350 g, were killed by lethal dose of pentobarbital sodium (50 mg/kg, iv). The antral part of the stomach was cut rapidly, and the mucosal layer was separated from the muscle layer. Circular muscle was dissected from the longitudinal layer using fine scissors, and cut into small segments (1 mm×4 mm). These segments were kept in a modified Kraft-Bruhe (K-B) medium at 4 °C for 15 min. The muscle segments were incubated at 36 °C in 4 mL of digestion medium [calcium-free physiological salt solution (Ca²⁺-free PSS)] containing 0.1% collagenase (II), 0.1% dithioerythritol, 0.15% trypsin inhibitor and 0.2% bovine serum albumin for 25–35 min. The softened muscle segments were then transferred into the modified K-B medium, and cells were dispersed individually by gentle agitation with a wide-pore fire-polished glass pipette. Isolated gastric myocytes

were kept in modified K-B medium at 4 °C until use.

Electrophysiological recording Isolated cells were transferred to a small chamber (0.1 mL) on the stage of an inverted microscope (IX-70 Olympus, Tokyo, Japan) for 10–15 min to settle down. The cells were superfused continuously with isosmotic PSS by gravity (0.9–1.0 mL/min). An 8-channel perfusion system (L/M-sps-8, List Electronics, Darmstadt, Germany) was used to change solution. Experiments were carried out at 20–25 °C and the whole-cell configuration of the patch-clamp technique was used. Patch-clamp pipettes were manufactured from borosilicate glass capillaries (GC 150T-7.5, Clark Electromedical Instruments, Kent, UK) using a 2-stage puller (PP-83, Narishige, Japan). The resistance of the patch pipette was 3–5 MΩ when filled with pipette solution. Liquid junction potentials were canceled before seal formation. Whole-cell currents were recorded with an EPC-10 patch-clamp amplifier (HEKA Instrument, Darmstadt, Germany) and command pulses were applied by using the Pentium IV-grade computer and pCLAMP software (Version 6.02; Axon Instruments, USA).

Drugs and solutions All drugs were purchased from Sigma (Sigma-Aldrich Corp St Louis, MO, USA). Tyrode's solution contained 147 mmol/L NaCl, 4 mmol/L KCl, 1.05 mmol/L MgCl₂·6H₂O, 2 mmol/L CaCl₂·2H₂O, 0.42 mmol/L NaH₂PO₄, 1.81 mmol/L Na₂HPO₄·2H₂O and 5.5 mmol/L glucose, and the pH was adjusted to 7.35 with NaOH. Ca²⁺-free solution contained 134.8 mmol/L NaCl, 4.5 mmol/L KCl, 10 mmol/L HEPES, 1 mmol/L MgCl₂ and 5 mmol/L glucose, and the pH was adjusted to 7.4 with TRIZMA BASE (Tris). The isosmotic solution (290 Osmmol/kg) contained 80 mmol/L NaCl, 4.5 mmol/L KCl, 1 mmol/L MgCl₂·6H₂O, 2 mmol/L CaCl₂·2H₂O, 5 mmol/L glucose, 10 mmol/L HEPES and 110 mmol/L sucrose, and the pH was adjusted to 7.4 with Tris. Hypoosmotic solution (200 Osmmol/kg) contained 30 mmol/L sucrose, with the other ingredients at the same concentrations as in the isosmotic solution. Modified K-B solution contained 50 mmol/L L-glutamate, 50 mmol/L KCl, 20 mmol/L taurine, 20 mmol/L KH₂PO₄, 3 mmol/L MgCl₂·6H₂O, 10 mmol/L glucose, 10 mmol/L HEPES, and 0.5 mmol/L egtazic acid, and the pH was adjusted to 7.4 with KOH. The pipette solution contained 110 mmol/L K-aspartic acid, 5 mmol/L Mg-ATP, 1 mmol/L MgCl₂·6H₂O, 20 mmol/L KCl, 0.1 mmol/L or 10 mmol/L egtazic acid, 2.5 mmol/L di-tris-creatine phosphate and 2.5 mmol/L disodium-creatine phosphate, and the pH was adjusted to 7.3 with KOH. AA, nordihydroguaiaretic acid (NDGA) and dimethyleicosadienoic acid (DEDA) were all prepared as aqueous stock solutions (1 mmol/L).

Data analysis Data were expressed as mean±SD. Statistical significance was evaluated using the Student's *t*-test.

Differences were considered to be statistically significant when $P < 0.05$.

Results

Effects of hyposmotic membrane stretch and AA on $I_{K(Ca)}$ and STOC Under whole-cell configuration, membrane potential was clamped at -60 mV, and $I_{K(Ca)}$ was elicited by a single-step command pulse from -60 mV to $+60$ mV for 400 ms at 15 s intervals. $I_{K(Ca)}$ started increasing at 139.3 ± 11.3 s after cells were exposed to hyposmotic solution (200 mOsm),

and at 165.0 ± 25.1 s after cells were exposed to $10 \mu\text{mol/L}$ AA (Figure 1A, 1B). There was no significant difference between the 2 latent periods. Hyposmotic membrane stretch and $10 \mu\text{mol/L}$ AA increased markedly in peak current of $I_{K(Ca)}$ to $168.3\% \pm 16.1\%$ and $158.5\% \pm 20.5\%$, respectively ($n=6$, Figure 1B.).

The calcium-activated potassium current is activated by intracellular Ca^{2+} and can be monitored by spontaneous transient outward currents (STOC)^[17]. We therefore observed STOC to investigate effects of hyposmotic membrane stretch and AA on $I_{K(Ca)}$. In whole cell configurations, the holding




Figure 1. Effects of hyposmotic membrane stretch and arachidonic acid (AA) on calcium-activated potassium currents ($I_{K(Ca)}$). (A) Raw traces of hyposmotic membrane stretch and AA-induced increase in $I_{K(Ca)}$. (B) The time courses of $I_{K(Ca)}$ increased by hyposmotic membrane stretch and arachidonic acid. Membrane potential was depolarized from -60 mV to $+60$ mV at 15 s intervals to elicit $I_{K(Ca)}$. (C) Effects of hyposmotic membrane stretch and AA on spontaneous transient outward currents.

potential was clamped at -20 mV, and STOC were elicited and enhanced by hyposmotic membrane stretch and 10 μmol/L AA, respectively (Figure 1C).

Effects of exogenous AA on hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ Under whole-cell configuration, membrane potential was clamped at -60 mV, and $I_{K(Ca)}$ was elicited by a step voltage command pulse from -40 mV to +100 mV for 400 ms with a 20-mV increment at 10-s

intervals. Exogenous AA significantly increased $I_{K(Ca)}$ elicited by the command step pulse when membrane potential was depolarized from -40 mV to +100 mV, and hyposmotic membrane stretch potentiated the AA-induced increase in $I_{K(Ca)}$ when the membrane potential was depolarized from -40 mV to +100 mV (Figure 2A, 2B). Hyposmotic membrane stretch also increased $I_{K(Ca)}$ elicited by the command step pulse when the membrane potential was depolarized from




Figure 2. The effect of exogenous arachidonic acid (AA) on the hyposmotic membrane stretch-induced increase in calcium-activated potassium currents ($I_{K(Ca)}$). (A) Raw traces of exogenous AA on the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$. (B) I/V relation curves. (C) Summary of the effects of exogenous AA on the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ and those of hyposmotic membrane stretch on the exogenous AA-induced increase in $I_{K(Ca)}$ at +60 mV. (D) The effect of exogenous AA on the hyposmotic membrane stretch-induced increase in spontaneous transient outward currents. ^b $P < 0.05$, ^c $P < 0.01$ vs control. ^e $P < 0.05$, ^f $P < 0.01$ vs 10 μmol/L AA. ^h $P < 0.05$, ⁱ $P < 0.01$ vs 202 mOsm.

-40 mV to +100 mV, and AA potentiated hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ when membrane potential was depolarized from -40 mV to +100 mV (Figure 2A, 2B). The peak current of $I_{K(Ca)}$ was increased to 184.2%±17.7% by hyposmotic membrane stretch and then the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ was potentiated by 10 μmol/L AA, and the peak current increased to 281.3%±28.3% at +60 mV ($n=8$, Figure 2C). In another way, the peak current of $I_{K(Ca)}$ was increased to 171.8%±20.3% by 10 μmol/L AA, and the AA-induced increase in $I_{K(Ca)}$ was potentiated by hyposmotic membrane stretch, with the peak current increasing to 311.5%±44.4% at +60 mV ($n=8$; Figure 2C). However, there was no significant difference between the potentiated effects of AA on the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ and hyposmotic membrane stretch on the AA-induced increase in $I_{K(Ca)}$ ($P>0.05$, Figure 2C). As with the protocol above, the effects of hyposmotic membrane stretch and AA on STOC were investigated.

Hyposmotic membrane stretch markedly increased STOC, and 10 μmol/L AA potentiated this effect ($n=2$, Figure 2D).

Effects of endogenous AA and its metabolites on hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ Intracellular free AA are metabolized by 3 enzymes: cyclooxygenase, lipoxygenase and epoxygenase. To determine the effects of endogenous AA and its metabolites on the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$, DEDA, a non-selective inhibitor of PLA₂, and NDGA, a lipoxygenase inhibitor, were used to inhibit the hydrolyzation of AA from membranes and to decrease the production of AA metabolites. DEDA (100 μmol/L in pipette) significantly suppressed the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ (Figure 3A,3B), and the increased percentage was reduced from 164.3%±9.8% of the control to 113.4%±3.6% at +60 mV ($n=15$, Figure 3C). When cells were pretreated for 15 min with 10 μmol/L NDGA, AA (Figure 4A,4B) and hyposmotic stretch-induced increases in $I_{K(Ca)}$ could be significantly suppressed (Figure 5A,




Figure 3. The inhibitory effect of dimethyleicosadienoic acid (DEDA), an inhibitor of phospholipase A₂ (PLA₂), on calcium-activated potassium currents ($I_{K(Ca)}$) increased by hyposmotic membrane stretch. (A) Raw traces of the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ in the presence of DEDA. (B) I/V relation curves of the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ in the presence of DEDA and control. Peak values of the currents were normalized to the values at +100 mV under control conditions. (C) The inhibitory effect of DEDA on $I_{K(Ca)}$ increased by hyposmotic membrane stretch at +60 mV. ^b $P<0.05$, ^c $P<0.01$ vs control. ^e $P<0.05$ vs 200 mOsm.




Figure 4. The inhibitory effect of nordihydroguaiaretic acid (NDGA), a lipoxygenase inhibitor, on calcium-activated potassium currents ($I_{K(Ca)}$) increased by arachidonic acid (AA). (A) Raw traces of the AA-induced increase in $I_{K(Ca)}$ in the presence of NDGA. (B) I/V relation curves of the AA-induced increase in $I_{K(Ca)}$ in the presence of NDGA and control. The peak values of currents were normalized to the values at +100 mV under control conditions. (C) The inhibitory effect of NDGA on $I_{K(Ca)}$ increased by AA at +60 mV. ^b $P < 0.05$, ^c $P < 0.01$ vs control. ^ε $P < 0.05$ vs NDGA.

5B). The AA-induced increase in $I_{K(Ca)}$ was decreased from $145\% \pm 10\%$ to $110\% \pm 4\%$ by NDGA ($n=8$, Figure 4C). NDGA also suppressed the hypotonic membrane stretch-induced increase in $I_{K(Ca)}$ (Figure 5A,5B) and the increased percentage was reduced from $170\% \pm 10\%$ to $142\% \pm 3\%$ at +60 mV (Figure 5C). These results showed that endogenous AA and its metabolites were involved in the hypotonic membrane stretch-induced increase in $I_{K(Ca)}$.

Effect of calcium mobilization on AA-induced increase in $I_{K(Ca)}$ It is well known that $I_{K(Ca)}$ is activated by intracellular free calcium, while extracellular calcium is necessary for efficiently controlling calcium homeostasis. To determine whether the AA-induced increase in $I_{K(Ca)}$ was mediated by calcium influx, the effect of AA on $I_{K(Ca)}$ was observed following the removal of extracellular calcium and the addition of 10 μmol/L EGTA in bath solution. The AA-induced increase in $I_{K(Ca)}$ was completely blocked by the removal of extracellular calcium, and the changes in the percentage of $I_{K(Ca)}$ were $146.30\% \pm 10.4\%$ and $95.64\% \pm 11.7\%$ in the presence or absence of extracellular calcium, respectively ($n=6$,

Figure 6A). Our previous study indicated that hypotonic membrane stretch activated L-type calcium currents^[12] and calcium-activated potassium currents via extracellular calcium influx through SAC in gastric myocytes of guinea pig^[15,16]. We therefore examined the relationship between the AA-induced increase in $I_{K(Ca)}$ and L-type calcium channels or SAC. However, 5 mmol/L nifedipine, an L-type calcium channel blocker, did not block the AA-induced increase in $I_{K(Ca)}$, but gadolinium (Gd^{3+}), a blocker of SAC, completely suppressed the AA-induced increase in $I_{K(Ca)}$. The changes in the percentage of $I_{K(Ca)}$ were $146.3\% \pm 10.4\%$, $151.1\% \pm 14.4\%$ and $102.5\% \pm 2.2\%$ in the control, nifedipine and Gd^{3+} groups, respectively ($n=6$, Figure 6A).

Intracellular free calcium has 2 sources: extracellular calcium influx and intracellular calcium release from calcium stores. Intracellular calcium is released through 2 pathways, one is CICR and the other is inositol-triphosphate-induced calcium release (IICR). We therefore investigated the role of intracellular calcium release in the AA-induced increase in $I_{K(Ca)}$. Heparin (3 mg/mL), a potent inhibitor of IICR, could




Figure 5. The inhibitory effect of nordihydroguaiaretic acid (NDGA), a lipoxygenase inhibitor, on calcium-activated potassium currents ($I_{K(Ca)}$) increased by hyposmotic membrane stretch. (A) Raw traces of the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ in the presence of NDGA. (B) I/V relation curves of the hyposmotic membrane stretch-induced increase in $I_{K(Ca)}$ in the presence of NDGA and control. The peak values of currents were normalized to the values at +100 mV under control conditions. (C) The inhibitory effect of NDGA on $I_{K(Ca)}$ increased by hyposmotic membrane stretch at +60 mV. ^b $P < 0.05$ vs control; ^e $P < 0.05$ vs NDGA; ^h $P < 0.05$ vs 200 mOsm.




Figure 6. The effect of calcium mobilization on $I_{K(Ca)}$ increased by arachidonic acid (AA) 10 μmol/L. (A) Summary of $I_{K(Ca)}$ increased by AA at +60 mV under conditions of no external Ca^{2+} , administration of nifedipine (Ni) and gadolinium (Gd^{3+}). (B) Effect of inositol triphosphate-induced calcium release on the AA-induced increase in spontaneous transient outward currents (STOC) and effect of calcium-induced calcium release on the AA-induced increase in STOC. ^b $P < 0.05$ vs control. ^e $P < 0.05$ vs Ni.

inhibit STOC significantly, but did not block the AA-induced increase in STOC (Figure 6B). Ryanodine (10 $\mu\text{mol/L}$), a specific CICR inhibitor, binds to CICR channels and locks them in a subconductance state, thereby functionally depleting calcium stores^[18]. In the present study, ryanodine increased STOC, and STOC were then almost abolished by ryanodine after approximately 8 min with caffeine, a CICR activator; AA could not then enhance them again (Figure 6B). These results indicated that CICR, but not IICR, was involved in the AA-induced increase in $I_{K(\text{Ca})}$.

Discussion

Our previous study demonstrated that $I_{K(\text{Ca})}$ was activated by hyposmotic membrane stretch and Ca^{2+} signaling played an important role in the process in gastric antral circular myocytes of guinea pig^[15,16]. Under hyposmotic conditions, extracellular calcium influx through SAC triggered CICR, and intracellular free calcium then activated $I_{K(\text{Ca})}$. However, it remains obscure how the membrane stretch is turned into the signal for Ca^{2+} entry from the extracellular space. We therefore investigated whether AA is involved in the hyposmotic membrane stretch-induced increase in $I_{K(\text{Ca})}$.

In the present study, both hyposmotic membrane stretch and AA significantly increased $I_{K(\text{Ca})}$ with a similar latent period. Moreover, exogenous AA potentiated the hyposmotic membrane stretch-induced increase in $I_{K(\text{Ca})}$ (Figure 1). The results indicated that there may be a similar mechanism for the $I_{K(\text{Ca})}$ activated by hyposmotic membrane stretch and AA. Activation of various signaling pathways may induce an increase in the production of AA, for example, phospholipase C, phospholipase D and PLA_2 . In mammalian tissues AA is mainly liberated directly from phospholipids by PLA_2 , which is a ubiquitous enzyme^[4]. We have observed that when cells are exposed to DEDA, a non-selective inhibitor of PLA_2 , the hyposmotic membrane stretch-induced increase in $I_{K(\text{Ca})}$ is significantly blocked by DEDA (Figure 3). The results suggest that hyposmotic membrane stretch activates PLA_2 , which hydrolyzes membrane phospholipids to produce AA, and AA as a second messenger mediates the hyposmotic membrane stretch-induced increase in $I_{K(\text{Ca})}$ in gastric myocytes of guinea pig. Many experiments also support our results. It was observed that hyposmotic cell swelling induced AA release from cell membranes in human neuroblastoma cells^[9] and Ehrlich ascites tumor cells^[10]. In rat inner medullary collecting duct cells, AA acted as a second messenger in hypotonicity-induced calcium transients^[11]. AA was also a second messenger in cultured rabbit principal cells^[19] and ciliary epithelial cells under hyposmotic condi-

tions^[20]. Meanwhile, many reports have described that AA is able to affect cell functions, via its metabolites, under hyposmotic conditions in several cell systems. Leucotrienes, for example, appeared to mediate the inositol efflux in glial cells^[21], and to activate chloride and potassium conductances as well as taurine transport in Ehrlich ascites cells^[22]. In the present study, we also examined the possibility that AA metabolites could be involved in activating potassium currents under hyposmotic conditions by using NDGA, a lipoxygenase inhibitor. NDGA significantly inhibited AA and the hyposmotic membrane stretch-induced increase in $I_{K(\text{Ca})}$ (Figure 4). These results suggest that AA metabolites generated by lipoxygenase mediate the hyposmotic membrane stretch-induced increase in $I_{K(\text{Ca})}$ in gastric myocytes.

In various cell types, AA was found to induce Ca^{2+} flux and to mobilize intracellular calcium to trigger different Ca^{2+} -dependent physiological functions in cells. For example, AA or its metabolites mobilized Ca^{2+} from intracellular stores, and intracellular Ca^{2+} then activated ion transport^[19]. In several cell types, for example, in rat IMCD cells^[11], human embryonic kidney (HEK293) cells^[23] and *Dictyostelium discoideum*^[24], AA released Ca^{2+} from the stores to trigger extracellular Ca^{2+} entry, and Ca^{2+} released from calcium stores was a prerequisite for extracellular Ca^{2+} entry. However, Murthy *et al*^[20] found that AA induces Ca^{2+} influx, which triggers CICR in longitudinal smooth muscle of the intestine. Our previous study also demonstrated that hyposmotic membrane stretch activates $I_{K(\text{Ca})}$ via CICR in gastric myocytes^[15]. In the present study, the roles of AA and its metabolites in the relationships among hyposmotic membrane stretch-induced increase in $I_{K(\text{Ca})}$, extracellular Ca^{2+} and intracellular calcium mobilization were investigated. Under extracellular calcium-free conditions, $I_{K(\text{Ca})}$ was not increased by AA or hyposmotic membrane stretch (Figure 6). It was elucidated that extracellular calcium is necessary for AA and the hyposmotic membrane stretch-induced increase in $I_{K(\text{Ca})}$, and some ionic channels participate in extracellular calcium influx. McCarty and O'Neil indicated that there are 2 alternative kinds of channel activated by hyposmotic swelling: voltage-activated Ca^{2+} channels and stretch-activated channels^[25]. We observed previously that hyposmotic membrane stretch increases L-type current in gastric myocytes of guinea pig^[12], and Yamamoto and Suzuki^[26] also observed that there are 2 kinds of SAC in gastric myocytes of guinea pig. In the present study we examined whether these 2 channels are associated with extracellular calcium influx. Nicardipine, an L-type calcium channel blocker, could not block the AA-induced increase in $I_{K(\text{Ca})}$. However, it was completely blocked by Gd^{3+} , which blocks not only SAC but also store-operated

Ca²⁺ channels (Figure 6). A similar effect of Gd³⁺ in blocking AA-induced Ca²⁺ entry has also been observed in other cell types, such as IMCD^[11] and HEK293 cells^[23].

Intracellular Ca²⁺ release from Ca²⁺ stores is the primary source of the increase in intracellular calcium. It was found that the entry of extracellular calcium via activating stretch-sensitive channels is amplified by calcium release from internal stores in toad gastric myocytes^[26]. Sutko and Airey^[27] suggested that ryanodine-sensitive calcium stores were positioned near the surface membrane in some smooth muscle cells, Ca²⁺ release from which was found to influence the activity of *I*_{K(Ca)}. Our previous study also indicated that hyposmotic membrane stretch activates *I*_{K(Ca)}^[15,16] and carbachol currents^[13], and the activations are associated with CICR, which is triggered by extracellular calcium influx^[15,28]. In the present study, heparin, a potent inhibitor of inositol triphosphate receptor, did not block the AA-induced increase in *I*_{K(Ca)}; however, ryanodine, a CICR agonist, completely blocked the AA-induced increase in *I*_{K(Ca)} (Figure 6B). The results suggest that AA mobilizes intracellular calcium via triggering CICR and activates *I*_{K(Ca)} in gastric antral circular myocytes of guinea pig. The potassium efflux through *I*_{K(Ca)} hyperpolarized the membrane potential of smooth muscle cells, thereby limiting depolarization-dependent calcium and promoting relaxation. CICR can thus participate in both the contraction and relaxation of smooth muscle cells. Therefore, AA may be involved in both the contraction via activating extracellular Ca²⁺ influx and relaxation via activating *I*_{K(Ca)} in gastric antral circular myocytes of the guinea pig.

In summary, hyposmotic membrane stretch may act on cell membranes to activate PLA₂ and then generate AA. AA may then act as a second messenger to mediate extracellular calcium entry and trigger CICR to activate *I*_{K(Ca)} in gastric myocytes of the guinea pig. AA and its metabolites may play an important role in regulating many cell functions under the hyposmotic conditions in gastric antral circular myocytes of the guinea pig.

References

- 1 Morris CE. Are stretch-sensitive channels in molluscan cells and elsewhere physiological mechanotransducers? *Experientia* 1992; 48: 852–8.
- 2 Ordway RW, Petrou S, Kirber MT, Walsh JV Jr, Singer JJ. Stretch activation of a toad smooth muscle K⁺ channel may be mediated by fatty acid. *J Physiol* 1995; 484: 331–7.
- 3 Dennis EA, Rhee SG, Billah MM, Hannun YA. Role of phospholipase in generating lipid second messengers in signal transduction. *FASEB J* 1991; 5: 2068–77.
- 4 Graber R, Sumida C, Nunez EA. Fatty acids and cell signal transduction. *J Lipid Mediat* 1994; 9: 91–116.

- 5 Liu Y, Liu D, Krafe DS. Decrease of inward rectification as a mechanism for arachidonic acid-induced potentiation of hKir2.3. *Eur Biophys J* 2002; 31: 497–503.
- 6 Jun JY, Yeum CH, Park YW, Jang IY, Kong ID, Sim JH, *et al*. Effects of arachidonic acid on ATP-sensitive K⁺ current in murine colonic smooth muscle cells. *Jpn J Pharmacol* 2002; 90: 81–7.
- 7 Chalimoniuk M, King-Pospisil K, Pedersen WA, Malecki A, Wylegala E, Mattson MP, *et al*. Arachidonic acid increases choline acetyltransferase activity in spinal cord neurons through a protein kinase C-mediated mechanism. *J Neurochem* 2004; 90: 629–36.
- 8 Gauthier KM, Spitzbarth N, Edwards EM, Campbell WB. Apamin-sensitive K⁺ currents mediate arachidonic acid-induced relaxations of rabbit aorta. *Hypertension* 2004; 43: 413–9.
- 9 Basavappa S, Pedersen SF, Jorgensen NK, Ellory JC, Hoffmann EK. Swelling-induced AA release via the 85-kDa cPLA₂ in human neuroblastoma cells. *J Neurophysiol* 1998; 79: 1441–9.
- 10 Thoroe SM, Lauritzen L, Lambert IH, Hansen HS, Hoffmann EK. Cell swelling activates phospholipase A₂ in Ehrlich ascites tumor cells. *J Membr Biol* 1997; 160: 47–58.
- 11 Tinel H, Wehner F, Kinne RKH. Arachidonic acid as a second messenger for hypotonicity-induced calcium transients in rat IMCD cells. *Pflügers Arch* 1997; 433: 245–53.
- 12 Xu WX, Kim SJ, Kim SJ, So I, Kang TM, Rhee JC, *et al*. Effect of stretch on calcium channel currents recorded from the antral circular myocytes of guinea-pig stomach. *Pfluegers Arch* 1996; 432: 159–64.
- 13 Cui YF, Li L, Yu YC, Jin ZY, Li ZL, Xu WX. Role of unsaturated fatty acids in the enhancement of muscarinic current by hyposmotic membrane stretch in guinea pig smooth muscle cells. *Acta Physiol Sin* 2003; 55: 96–100.
- 14 Zheng HF, Li XL, Jin ZY, Sun JB, Li ZL, Xu WX. Effect of unsaturated fatty acid on calcium-activated potassium current in guinea pig gastric myocytes. *World J Gastroenterol* 2005; 11: 672–5.
- 15 Piao L, Li Y, Li L, Xu WX. Increment of calcium-activated and delayed rectifier potassium current by hyposmotic swelling in gastric antral circular myocytes of guinea pig. *Acta Pharmacol Sin* 2001; 22: 566–72.
- 16 Piao L, Li Y, Li L, Ng J, Li ZL, Xu WX. The involvement of calcium mobilization in the calcium-activated potassium currents activated by hyposmotic swelling in gastric antral circular myocytes of the guinea-pig. *Jpn J Physiol* 2001; 51: 223–30.
- 17 Bolton TB, Lim SP. Properties of calcium stores and transient outward currents in single smooth muscle cells of rabbit. *J Physiol (Lond)* 1988; 409: 385–400.
- 18 Sutko JL, Airey J. Ryanodine receptor Ca²⁺ release channels. *Physiol Rev* 1996; 76: 1027–71.
- 19 Ling BN, Webster CL, Eaton DC. Eicosanoids modulate apical Ca²⁺-dependent K⁺ channels in cultured rabbit principal cells. *Am J Physiol* 1992; 63: F116–26.
- 20 Civan MM, Coca-Prados M, Peterson-Yantorno K. Pathways signaling the regulatory volume decrease of cultured nonpigmented ciliary epithelial cells. *Invest Ophthalmol Vis Sci* 1994; 5: 2876–86.
- 21 Strange K, Morrison R, Shrode L, Putnam R. Mechanism and regulation of swelling-activated inositol efflux in brain glial cells. *Am J Physiol* 1993; 265: C244–56.

- 22 Hoffmann EK, Dunham PB. Membrane mechanisms and intracellular signalling in cell volume regulation. *Int Rev Cytol* 1995; 161: 173–262.
- 23 Luo D, Broad LM, Bird GSJ, Putney Jr WJ. Signaling pathways underlying muscarinic receptor-induced $[Ca^{2+}]_i$ oscillations in HEK293 cells. *J Biol Chem* 2001; 276: 5613–21.
- 24 Schaloske R, Sonnemann J, Malchow D, Schlatterer C. Fatty acids induce release of Ca^{2+} from acidosomal stores and activate capacitative Ca^{2+} entry in *Dictyostelium discoideum*. *Biochem J* 1998; 332: 541–8.
- 25 McCarty NA, O'Neil RG. Calcium signaling in cell volume regulation. *Physiol Rev* 1992; 2: 1037–61.
- 26 Yamamoto Y, Suzuki H. Two types of stretch-activated channel activities in guinea-pig gastric smooth muscle cells. *Jpn J Physiol* 1996; 6: 337–45.
- 27 Sutko JL, Airey J. Ryanodine receptor Ca^{2+} release channels. *Physiol Rev* 1996; 6: 1027–71.
- 28 Yu YC, Guo HS, Piao L, Li L, Lee ZL, Xu WX. Intracellular calcium involved in muscarinic currents increased by hyposmotic membrane stretch in gastric myocytes of guinea-pig. *Acta Pharmacol Sin* 2002; 23: 961–1056.
-

The 15th World Congress of Pharmacology (IUPHAR-2006)

2006, July 2-7 Beijing International Convention Center, China

Info: Ms Xiao-dan Zhao
Chinese Pharmacological Society
1, Xian Nong Tan St
Beijing 100050, China
Phn/Fax 86-10-6316-5211
E-mail zhxd@imm.ac.cn
<http://www.cnphars.org>