

## Full-length article

# GLB prevents tumor metastasis of Lewis lung carcinoma by inhibiting tumor adhesion actions<sup>1</sup>

Yan PAN<sup>2</sup>, Qian-liu SONG<sup>2</sup>, Yan-hua LIN<sup>2</sup>, Ning LU<sup>2</sup>, He-ming YU<sup>3</sup>, Xue-jun LI<sup>2,4</sup><sup>2</sup>Department of Pharmacology, School of Basic Medical Sciences and State Key Laboratory of Natural Biomimetic Drugs, Peking University, Beijing 100083, China; <sup>3</sup>National Research Institute for Family Planning, Beijing 100081, China

## Key words

neoplasm metastasis; adhesion; carcinoma, Lewis lung

<sup>1</sup> Project supported by the National Natural Science Foundation of China (No 39770286, 30171090) and 973 Program of the Ministry of Science and Technology (No 2004CB518902).<sup>4</sup> Correspondence to Prof Xue-jun LI.

Phn 86-10-8280-2863

Fax 86-10-6217-9119.

E-mail xjli@bjmu.edu.cn

Received 2005-01-29

Accepted 2005-03-14

doi: 10.1111/j.1745-7254.2005.00125.x

## Abstract

**Aim:** To investigate the inhibitory effect of a new compound of GLB on tumor metastasis *in vivo* and analyze its actions on tumor cell adhesion to clarify its mechanism. **Methods:** The effect of GLB on tumor metastasis was analyzed by Lewis lung carcinoma model. The pathological morphology of lung alveolar was evaluated by hematoxylin-eosin staining. The effect of GLB on the proliferation of human prostate cancer cell (PC-3M, with a high metastatic characteristic) was studied using the MTT method, and its actions on PC-3M cell adhesion to human umbilical vein endothelial cells (HUVEC) and laminin were analyzed *in vitro*. **Results:** GLB (100 mg·kg<sup>-1</sup>·d<sup>-1</sup> for 28 d, ig) reduced the number of lung colonies of Lewis lung carcinoma metastasis significantly ( $P < 0.05$ ). Simultaneously, GLB could mitigate the damage of lung alveolar caused by metastatic tumor deposits. *In vitro*, GLB inhibited dramatically the adhesion of PC-3M cells to HUVEC ( $P < 0.01$ ) and laminin ( $P < 0.05$ ), without cytotoxic or anti-proliferative action on PC-3M cells. **Conclusion:** GLB has anti-tumor metastatic activity, which partly depends on its inhibition of tumor adhesion.

## Introduction

The ability to metastasis is the most fearsome aspect of cancer and most cancer deaths are the sequel of metastatic diseases rather than primary tumor growth. In order to form overt metastases, a cell must complete the metastatic cascade, a series of well-defined steps including local invasiveness, intravasation, circulation, adhesion and extravasation, survival, proliferation and angiogenesis<sup>[1]</sup>. Within this multistep process, adhesive ability of metastatic tumor cells is a critical factor in extravasation and formation of new tumor foci<sup>[2-6]</sup>.

Cancer continues to represent the largest cause of mortality in the world and claims over 6 million lives every year<sup>[7]</sup>. An extremely promising strategy for the management of cancer today is chemoprevention, which is defined as the use of synthetic or natural agents (alone or in combination) to block the development of cancer in humans<sup>[8,9]</sup>. Studies on the pharmacological mechanisms and the search for new anticancer and anti-metastasis drugs are necessary and hold

great interest for scientists. The compound GLB, (2''S,3aR, 6S,7aR)-3''-acetyl-3a,7a-dihydro-2,2,2',2'-tetramethyl-5''-(4-bromophenyl)-spiro{spiro[1,3-dioxolo(4,5-D)-pyrane-6(7H), 5'(4'H)-1',3'-dioxolo]-7(6H),2''(3'H)-(1,3,4)-oxadiazole} (Figure 1), was synthesized by Prof Zhong-jun LI (School of Pharmacy, Peking University) in 2000 and was patented by National Patent of China (No 03119612.8). It is structurally based on fructose. Previous studies showed that the com-




Figure 1. Chemical structure of GLB.

pound could inhibit the growth of S180 carcinoma in ICR and LACA mice. However, its precise function and mechanism of action on tumor and tumor metastasis have not been determined in full. In the present study, we investigated the effect of GLB on tumor metastasis in mice treated orally and examined its effect on the adhesion of tumor cells *in vitro* to clarify its possible mechanism of anti-metastasis.

## Materials and methods

**Cell culture** Human umbilical vein endothelial cells (HUVEC) were isolated by collagenase I (1 g/L, Invitrogen, USA) digestion of umbilical veins from undamaged sections of fresh cords. HUVEC were grown in Medium 199 (Gibco, USA) supplemented with 20% heat-inactivated fetal bovine serum (FBS), 20 mg/L endothelial cell growth supplement (ECGS, Sigma, USA),  $1 \times 10^5$  IU/L penicillin, 100 mg/L streptomycin and 2 mmol/L *L*-glutamine. Replicate cultures were obtained by trypsinization and cells of passages 3–6 were used in the experiments. The identification of HUVEC was confirmed by their polygonal morphology and by detecting their immunoreactivity for factor VIII-related antigens<sup>[10]</sup>. Human prostate carcinoma cells (PC-3M) were cultivated in RPMI-1640 containing 10% heat-inactivated FBS,  $1 \times 10^5$  IU/L penicillin and 100 mg/L streptomycin in a humidified incubator with 5% CO<sub>2</sub> in air at 37 °C.

**Assay of PC-3M cell adhesion to HUVEC** PC-3M cells were labeled with CFDA (carboxyfluorescein diacetate, Sigma, 100 mg/L) for 30 min at 37 °C<sup>[11]</sup>. Cells were washed twice with phosphate-buffered saline (PBS) to remove residual fluorescent dye. The viability of the cells was not compromised by this labeling protocol, as indicated by Trypan blue exclusion.

HUVEC were cultured to confluence in 96-well dishes coated with 2% gelatin. Confluence was confirmed using microscopy and CFDA-labeled PC-3M cells were washed and resuspended in culture medium. One hundred microliters of the tumor-cell suspension (about  $1 \times 10^4$  cells) was added to HUVEC monolayers in 96-well dishes and co-cultured for 30 min at 37 °C. At the end of the experiment, the wells were washed twice with PBS to remove non-adhering PC-3M cells. The adherent cells were counted under a fluorescence microscope using an excitation wavelength of 492 nm, using five fields in each well<sup>[10]</sup>.

**Assay of PC-3M cell adhesion to laminin** The 96-well plates were coated with laminin (extracted by the Department of Cytobiology, Peking University) and incubated in RPMI-1640 containing 0.02% bovine serum albumin (BSA) for 15 min at 37 °C. In a total volume of 100  $\mu$ L RPMI-1640,

$8 \times 10^4$  PC-3M cells were added to each well and incubated for 4 h at 37 °C in the presence of GLB. After removing unattached cells with two washes with PBS, the attached cells were incubated with 20  $\mu$ L sterile MTT dye (Sigma) for a further 2 h at 37 °C. The medium was then removed, and 100  $\mu$ L dimethyl sulfoxide (Me<sub>2</sub>SO) was added and mixed thoroughly. Spectrometric absorbance at 540 nm (for formazan dye) and 690 nm (as background level) was measured using a microplate reader (Bio-Rad, USA)<sup>[12]</sup>.

**Proliferation assay** PC-3M cells ( $1 \times 10^4$  cells/well) were cultured in 24-well plates and incubated for 1, 2, 3, 4, 5, and 6 d in the presence of GLB at various concentrations. The medium with GLB was changed every 48 h. After incubation the cells were harvested and washed with PBS. The number of viable cells in each well was then determined and counted using the Trypan blue exclusion assay<sup>[13]</sup>.

**Lewis lung carcinoma *in vivo* model** Female C57BL/6 mice weighing 18 g–20 g were used and purchased from the Experimental Animal Center of Peking University (Grade II, Certificate No 11-00-0004). Lewis lung carcinoma (provided by the Chinese Medical Science Institute) was maintained in C57BL/6 mice by subcutaneous injection in the axillary's region of 0.2 mL of homogenized tumor tissue [tumor tissue (g): 0.9% sodium chloride (mL), 1:3] prepared from donors similarly inoculated for experimental tumor transplantation<sup>[14]</sup>.

**Metastasis *in vivo*** GLB was suspended in 0.5% (w/v) sodium carboxyl methylcellulose (CMC-Na) in distilled water and administered orally for 27 d (25, 50, and 100 mg·kg<sup>-1</sup>·d<sup>-1</sup>) from the next day of tumor cell injection. After the d 6, the tumor volumes of mice were measured every 3 d. At the d 28, the mice were killed and the lungs and primary tumors were removed and weighed. The number of metastasized pulmonary colonies was counted and lung tissues were fixed in formalin for further analysis. The inhibitory rate of lung metastasis (%) was calculated using the equation  $(W_{\text{model}} - W_{\text{treatment}}) / (W_{\text{model}} - W_{\text{control}}) \times 100\%$ , where *W* is lung wet weight<sup>[14–16]</sup>.


**Pathological evaluation of lung** The fixed lung tissue samples were embedded in paraffin, sectioned at 5  $\mu$ m and stained with hematoxylin and eosin to evaluate alveolar integrity by counting area percentage of alveolus in lung. A total of 5 fields from each lung sample were screened randomly; the mean value was accepted as representative of the sample<sup>[17]</sup>.

**Statistical analysis** The experimental results were expressed as mean  $\pm$  standard deviation (SD). Statistical analysis was carried out using one-way analysis of variance (ANOVA) followed by Dunnett's test with SPSS version

10.0.  $P < 0.05$  was considered to be statistically significant.

## Results

**Inhibitory effects of GLB on spontaneous metastasis of Lewis lung carcinoma in mice** We investigated the effect of oral administration of GLB on the spontaneous metastasis of Lewis lung carcinoma. GLB at the dose of 100 mg/kg significantly reduced the number of pulmonary metastatic colonies of Lewis lung carcinoma cells ( $P < 0.05$ ). The inhibitory rate of colony formation was approximately 31%. During the drug treatment, we found neither body weight loss nor toxic death in the GLB-treated mice. In addition to a reduction in the number of lung colonies by GLB, the survival rate of mice treated with GLB was significantly higher than that of untreated mice bearing Lewis lung carcinoma (Table 1). Although the mean weight and volume of the primary tumor in GLB-treated mice were less than those of vehicle-treated mice, there was no statistical difference ( $P > 0.05$ , Table 1, Figure 2).



**Figure 2.** Effect of GLB on Lewis lung carcinoma primary tumor of C57BL/6 mice. C57BL/6 mice were implanted with Lewis lung carcinoma cells and systemic treated with vehicle (model) and doses of 25 mg/kg GLB (GLBL), 50 mg/kg GLB (GLBM) or 100 mg/kg GLB (GLBH) intragastrically once a day. After the 6 d, the mice tumor volume was measured every 3 d.  $n > 7$  mice. Mean  $\pm$  SD.

**Effect of GLB on alveolar integrity in mice with Lewis lung carcinoma cells** Alveolar integrity of lungs with metastatic tumor deposits in mice bearing Lewis lung carcinoma cells was significantly worse than that of normal lung tissue (area percentage of alveolus in lungs:  $37.8\% \pm 7.44\%$  vs  $44.9\% \pm 7.6\%$ ,  $P < 0.05$ ). The lung tissue of mice treated with 100 mg/kg GLB was clearly better than that of tumor transplanted mice (area percentage of alveolus in lungs:  $45.1\% \pm 6.13\%$  vs  $37.8\% \pm 7.4\%$ ,  $P < 0.05$ ) (Figure 3).

**Inhibitory effects of GLB on metastatic adhesion function of tumor cells** We evaluated the effects of GLB on the metastatic adhesion functions of tumor cells by examining its effect on PC-3M cell adhesion to endothelial monolayers and laminin. GLB, at the concentrations of  $1 \times 10^{-5}$  mol/L and  $10^{-6}$  mol/L, inhibited significantly the adhesion of PC-3M cells to HUVEC ( $P < 0.01$ , Figure 4). GLB at these concentrations also inhibited the attachment of PC-3M cells to the laminin ( $P < 0.05$ , Figure 5). To clarify whether the inhibitory action on adhesion function could be due to a cytotoxic effect, GLB was tested for its effect on PC-3M cell proliferation (Figure 6). GLB had no substantial effect on the growth of tumor cells after up to 5 d incubation. Until the latter period of cell growth (d 6 and 7), the proliferation of PC-3M cells treated with  $1 \times 10^{-5}$  mol/L and  $1 \times 10^{-6}$  mol/L GLB showed signs of a slight decrease.


## Discussion

As has been demonstrated, the sequential process of micrometastasis formation by Lewis lung carcinoma includes single cells in the blood circulation escaping from the immune system, adhering to epithelial cells, migrating into the tissue, proliferating and making colonies in the lung<sup>[18]</sup>. Investigate the earliest events during micrometastasis formation is useful and could be simply applied for the estimation of anti-metastatic and anti-adhesive effects of various anti-cancer agents<sup>[12]</sup>.

**Table 1.** Effect of GLB on lung colonization of Lewis lung carcinoma cells. Mean  $\pm$  SD. <sup>b</sup> $P < 0.05$  vs model.

Group	Dose (mg/kg, ig)	Body weight (g)	Survival (n)	Tumor weight (g)	Lung colonies (n)	Inhibitory rate of lung colonies (%)
Model	Vehicle	19.8 $\pm$ 1.96	9/13	7.42 $\pm$ 1.78	16 $\pm$ 2.6	
GLBL	25	20.25 $\pm$ 1.54	7/9	6.41 $\pm$ 1.51	15 $\pm$ 5.0	6
GLBM	50	19.08 $\pm$ 2.67	7/9	6.18 $\pm$ 1.11	12 $\pm$ 5.2	26
GLBH	100	19.67 $\pm$ 2.64	8/9	6.04 $\pm$ 1.37	11 $\pm$ 3.1 <sup>b</sup>	31


GLBL, GLB 25 mg  $\cdot$  kg<sup>-1</sup>  $\cdot$  d<sup>-1</sup>; GLBM, GLB 50 mg  $\cdot$  kg<sup>-1</sup>  $\cdot$  d<sup>-1</sup>; GLBH, GLB 100 mg  $\cdot$  kg<sup>-1</sup>  $\cdot$  d<sup>-1</sup>.



**Figure 3.** Effect of GLB on alveolar integrity in mice with Lewis lung carcinoma cells. The area percentage of alveolus in lung tissue of normal C57BL/6 mice was almost 50% (A), and that in lung of mice bearing Lewis lung carcinoma (model) was decreased (B). Lung tissues of C57BL/6 mice bearing Lewis lung carcinoma treated with (C) 25 mg/kg, (D) 50 mg/kg and (E) 100 mg/kg GLB were significantly better than the model. Hematoxylin-eosin stain,  $\times 100$ . (F) The statistical result of alveolus area percentage. Control, normal; M, model; GLBL, GLB 25  $\text{mg}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$ ; GLBM, GLB 50  $\text{mg}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$ ; GLBH, GLB 100  $\text{mg}\cdot\text{kg}^{-1}\cdot\text{d}^{-1}$ .  $n=5$  mice. Mean $\pm$ SD. <sup>b</sup> $P<0.05$  vs model. <sup>r</sup> $P<0.05$  vs control.

The antitumor activities of the derivatives of oxadiazoles, one-membered heterocycles, have been reported and are found to be related to their chelating abilities or non-polarity,


which is essential to penetrate through intracellular sites<sup>[19]</sup>. These include the new 5-(2-amino-3-pyridyl)-2-thioxo-3H-1,3,4-oxadiazole derivatives that have shown cytotoxic




**Figure 4.** Inhibitory effect of GLB on adhesion of PC-3M to human umbilical vein endothelial cells (HUVEC). Carboxyfluorescein diacetate (CFDA)-labelled PC-3M cells were added to the HUVEC monolayer cultured at 37 °C for 30 min. The adherent tumor cells were counted in five fields in each plate, using a fluorescence microscope. (A) Control; (B) 1×10<sup>-7</sup> mol/L GLB; (C) 1×10<sup>-6</sup> mol/L GLB; (D) 1×10<sup>-5</sup> mol/L GLB. Magnification ×400. (E) Inhibition of PC-3M adhesion to HUVEC by GLB. n=4 wells. Mean±SD. <sup>c</sup>P<0.01 vs control.

activity against the cells of 4 human tumor cell lines<sup>[20]</sup> and 2, 5-disubstituted-1,3,4-oxadiazole compounds, which are potential anticancer agents<sup>[21]</sup>. The present study demonstrated that oral administration of GLB inhibited the spontaneous pulmonary metastasis of tumor cells in C57BL/6 mice and increased the survival rate. GLB significantly inhibited the adhesion by PC-3M cells without affecting cell proliferation *in vitro*. The present data suggest that GLB prevents tumor metastasis partly by inhibiting the metastatic adhesion of tumor cells.

Tumor cell adhesion to endothelial monolayers is a critical step in tumor metastasis<sup>[22]</sup>, as is laminin expression in



**Figure 5.** Inhibitory effect of GLB on adhesion of PC-3M to laminin. PC-3M cells were added to wells coated with laminin and incubated at 37 °C for 4 h in the presence of GLB. The number of adhesive cells per well was determined using the MTT assay and are shown as absorbance. n=6 wells. Mean±SD. <sup>b</sup>P<0.05 vs control.



**Figure 6.** Effect of GLB on proliferation of PC-3M cells. PC-3M cells were cultured for 7 d in the absence (■) or presence of either 1×10<sup>-6</sup> mol/L (□) or 1×10<sup>-5</sup> mol/L (▲) GLB. n=4 wells. Mean±SD. <sup>b</sup>P<0.05 vs control.

endothelial cells, being one of the earliest extracellular matrix (ECM) proteins. Laminin interaction with the β-1 integrin has been shown to have an important role in the regulation of tumor-to-endothelial cell adhesion<sup>[10]</sup>. We demonstrated that GLB significantly inhibited the adhesion of tumor cells to laminin, using the same doses of GLB as had been found to inhibit adhesion of tumor cells to endothelial cells. This suggests that the anti-adhesion activity of GLB might be due to its effect on laminin, one of many other factors involved in adhesion.

β-1 Integrin is known to recognize many ECM proteins as ligands, such as laminin, collagen and fibronectin. Laminin has been identified as the most important of these ligands for β-1 integrin in the cell adhesion process<sup>[23,24]</sup>. Whether GLB exerts its function through β-1 integrin or not is yet to be determined.

In the present study, GLB showed no effect on the weight of the primary tumor site in C57BL/6 mice subcutaneously injected with a Lewis lung carcinoma tumor cell suspension (Table 1). In a preliminary experiment where S180 carcinomas were implanted subcutaneously into ICR and LACA mice, GLB-treated mice (50 mg·kg<sup>-1</sup>·d<sup>-1</sup> for 12 d, ip) showed a significant decrease in tumor weight, with inhibitory rates of 43% and 32% (tumor weight at d 12, mean±SD: 1.16±0.67 g vs 2.05 ± 0.64 g, *P*<0.01 for ICR mice; 1.81±0.68 g vs 2.65±0.81 g, *P*<0.05 for LACA mice; *n*=10 mice). The different effects of GLB on these *in vivo* assay models may be dependent on the kind of tumor, the period of the experiment and route of administration. We speculate that there might be other mechanisms of action for GLB that require further study, including the relationship between its effects on tumor growth or metastasis and its inhibition of angiogenesis.

In conclusion, the compound GLB prevents tumor metastasis by inhibiting tumor adhesion. This suggests it has potential for future anti-metastatic development.

## References

- 1 Gassmann P, Enns A, Haier J. Role of tumor cell adhesion and migration in organ-specific metastasis formation. *Onkologie* 2004; 27: 577–82.
- 2 Nicolson GL. Metastatic tumor cell interactions with endothelium, basement membrane and tissue. *Curr Opin Cell Biol* 1989; 1: 1009–19.
- 3 Lewalle JM, Cataldo D, Bajou K, Lambert CA, Foidart JM. Endothelial cell intracellular Ca<sup>2+</sup> concentration is increased upon breast tumor cell contact and mediates tumor cell transendothelial migration. *Clin Exp Metastasis* 1998; 16: 21–9.
- 4 Honn KV, Tang DG. Adhesion molecules and tumor cell interaction with endothelium and subendothelial extracellular matrix. *Cancer Metastasis Rev* 1992; 11: 353–75.
- 5 Chambers AF, MacDonald IC, Schmidt EE, Koop S, Morris VL, Khokha R, *et al*. Steps in tumor metastasis: new concept for intravital videomicroscopy. *Cancer Metastasis Rev* 1995; 14: 279–301.
- 6 Tressler RJ, Yeatman T, Nicolson GL. Extracellular annexin VI expression is associated with divalent cation-dependent endothelial cell adhesion of metastatic RAW117 large-cell lymphoma cells. *Exp Cell Res* 1994; 215: 395–400.
- 7 Abdullaev FI, Luna RR, Roitenburd BV, Espinosa AJ. Pattern of childhood cancer mortality in Mexico. *Arch Med Res* 2000; 31: 526–31.
- 8 Gupta M, Mazumder UK, Kumar RS, Kumar TS. Antitumor activity and antioxidant role of *Bauhinia racemosa* against Ehrlich ascites carcinoma in Swiss albino mice. *Acta Pharmacol Sin* 2004; 25: 1070–6.
- 9 Rao KVK, Schwartz SA, Nair HK, Aalinkeel R, Mahajan S, Chawda R, *et al*. Plant derived products as a source of cellular growth inhibitory phytochemicals on PC-3M, DU-145 and LNCaP prostate cancer cell lines. *Curr Sci* 2004; 87: 1585–8.
- 10 Andrews EJ, Wang JH, Winter DC, Laug WE, Redmond HP. Tumor cell adhesion to endothelial cells is increased by endotoxin via an upregulation of β-1 integrin expression. *J Surg Res* 2001; 97: 14–9.
- 11 Woodward JK, Nichols CE, Rennie IG, Parsons MA, Murray AK, Sisley K. An *in vitro* assay to assess uveal melanoma invasion across endothelial and basement membrane barriers. *Invest Ophthalmol Vis Sci* 2002; 43: 1708–14.
- 12 Yoshitomi Y, Nakanishi H, Kusano Y, Muneshue S, Oguri K, Tatematsu M, *et al*. Inhibition of experimental lung metastases of Lewis lung carcinoma cells by chemically modified heparin with reduced anticoagulant activity. *Cancer Lett* 2004; 207: 165–74.
- 13 Tozawa K, Kawai N, Hayashi Y, Sasaki S, Kohri K, Okamoto T. Gold compounds inhibit adhesion of human cancer cells to vascular endothelial cells. *Cancer Lett* 2003; 196: 93–100.
- 14 Xiang Y, Ma B, Li T, Yu HM, Li XJ. Acetazolamide suppresses tumor metastasis and related protein expression in mice bearing Lewis lung carcinoma. *Acta Pharmacol Sin* 2002; 23: 745–51.
- 15 Prontera C, Mariani B, Rossi C, Poggi A, Rotilio D. Inhibition of gelatinase A (MMP-2) by batimastat and captopril reduces tumor growth and lung metastases in mice bearing Lewis lung carcinoma. *Int J Cancer* 1999; 81: 761–6.
- 16 Ma B, Xiang Y, Li T, Yu HM, Li XJ. Inhibitory effect of topiramater on Lewis lung carcinoma metastasis and its relation with AQP1 water channel. *Acta Pharmacol Sin* 2004; 25: 54–60.
- 17 Ustundag N, Bozkurt AK, Demirkaya A, Koksak C, Mayda AS. Histopathological and immunohistochemical detection of protective effects of University of Wisconsin solution supplemented with iloprost on donor lung damage. *Transplant Proc* 2004; 36: 1271–4.
- 18 Hirano S. *In vitro* and *in vivo* cytotoxic effects of nitric oxide on metastatic cells. *Cancer Lett* 1997; 115: 57–62.
- 19 Mishra L, Said MK, Itokawa H, Takeya K. Antitumor and antimicrobial activities of Fe(II)/Fe(III) complexes derived from some heterocyclic compounds. *Bioorg Med Chem* 1995; 3: 1241–5.
- 20 Liszkiewicz H, Kowalska MW, Wietrzyk J, Opolski A. Synthesis and anti-proliferative activity *in vitro* of new 5-(2-amino-3-pyridyl)-2-thioxo-3H-1,3,4-oxadiazole derivatives. *Indian J Chem, B* 2003; 42B: 2846–52.
- 21 Shah HP, Shah BR, Bhatt JJ, Desai NC, Trivedi PB, Undavia NK. Synthesis of 2,5-disubstituted 1,3,4-oxadiazoles as potential antimicrobial, anticancer and anti-HIV agents. *Indian J Chem, B* 1998; 37B: 180–2.
- 22 Danen EH, Marcinkiewicz C, Cornelissen IM, van Kraats AA, Pachter JA, Ruiter DJ, *et al*. The disintegrin eristostatin interferes with integrin α4β1 function and with experimental metastasis of human melanoma cells. *Exp Cell Res* 1998; 238: 188–96.
- 23 Dejana E. Endothelial cell adhesive receptors. *J Cardiovasc Pharmacol* 1993; 21: S18–21.
- 24 Kitayama J, Nagawa H, Tsuno N, Osada T, Hatano K, Sunami E, *et al*. Laminin mediates tethering and spreading of colon cancer cells in physiological shear flow. *Br J Cancer* 1999; 80: 1927–34.