

Diastereoselective Synthesis of Enantioenriched, Annulated Tetrahydrofurans by Simultaneous Formation of the O-1–C-5 and the C-5–C-4 Bonds

Seda Ünalı, ^a Mustafa Özlügedik, ^{a,b} Roland Fröhlich, ^{a,c} Dieter Hoppe ^{a,*}

^a Organisch-Chemisches Institut, Westfälische Wilhelms-Universität Münster, Corrensstraße 40, 48149 Münster, Germany
Fax: (+49)-251-83-36531, e-mail: dhoppe@uni-muenster.de

^b Present address: DyStar Textilfarben GmbH & Co. Deutschland KG, F & F VE, P. O. Box 100 480, 51304 Leverkusen, Germany

Fax: (+49)-214-30-55970, e-mail: ozluedik@gmx.de

^c Author to whom correspondence about the X-ray analysis should be addressed

Received: April 6, 2005; Accepted: June 13, 2005

Abstract: Enantiomerically enriched 1-[2-(1-hydroxy-alkyl)cycloalkylidene]methyl *N,N*-diisopropylcarbamates, which are easily available by an asymmetric homoaldol reaction, condense with aldehydes under the influence of boron trifluoride etherate to form cyclohexano- or cyclopentano-annulated tetrahydrofuran-3-carboxaldehydes, bearing two quaternary carbon

atoms and four adjacent stereogenic centers, with complete diastereoselectivity.

Keywords: diastereoselective synthesis; enantioenriched tetrahydrofurans; α,β -enediolate synthons; enol carbamates; Mukaiyama reaction; oxabicyclic compounds


Introduction

Tetrahydrofurans are substructures of numerous natural products,^[1] and a great variety of methods for their synthesis exists.^[2] In 1989 we published a highly efficient and surprisingly simple general method for the synthesis of monocyclic stereohomogeneous, 2,3,4,5-tetrasubstituted tetrahydrofurans **4** (Scheme 1),^[3] from homoaldol products^[1,4] **1** and aldehydes **2**. The assumed decisive in-

termediate is the (*E*)-oxonium ion **3**, which is formed under the influence of the strong Lewis acid BF₃; it undergoes an intramolecular Mukaiyama-type^[5] addition of the enolic moiety onto the former carbonyl carbon atom from the least hindered conformation. Subsequently, the carbamoyl group is extruded. The reaction is also applicable to ketals^[1] and for intramolecular condensations.^[3b] Some mechanistically related syntheses were published later.^[6]

By applying these methods we expected to realise a synthesis of tetrahydrofurans **10** by formal combination of the dianion **C** with two different aldehydes **A** and **B** (Figure 1).

We now report the realization of the above-mentioned pathway to provide a synthesis of highly substituted bicyclic tetrahydrofurans.



Scheme 1. Synthesis of tetrasubstituted tetrahydrofurans.








Figure 1. Formation of the bicycles **10** in a highly regio- and stereoselective manner.

Table 1. Diastereomerically pure bicyclic tetrahydrofurans **10**.


Starting materials 7 ee [%] ^[a]	Product 10	Yield [%]	ee [%] ^[b]	Alcohol 11 (yield, %)	<i>p</i> -Nitrobenzoate 12 (yield, %)
7a [76]	 10a	55	[78]	11a (70)	12a (79)
7b [78]	 10b	54	[81]	11b (72)	12b (81)
7c [86]	 10c	62	[91]	11c (73)	12c (84)
7d [93]	 10d	63	[96]	11d (74)	12d (85)

^[a] Enantiomeric excess was determined by ¹H NMR shift experiments.^[7c]

^[b] Enantiomeric excess was determined by chiral HPLC.

Results and Discussion


We recently observed that the carbanionic species, obtained by (–)-sparteine-mediated deprotonation of 1-(2-alkylcycloalk-1-enyl)methyl carbamates of type **5** exhibit considerable configurational stability and, hence, can be converted efficiently to optically active homoaldol products **7** with up to 96% ee (Scheme 2).^[7] The condensation of **7** with aromatic or aliphatic aldehydes^[8] or 2-alkenals in the presence of 1 equivalent of boron trifluoride etherate at 0 °C or 20 °C proceeds smoothly and, after aqueous work-up, diastereomerically pure tetrahydrofurans **10** can be isolated (Table 1). Further diastereomers could not be detected by ¹H NMR. The enantiomeric ratios of aldehydes **10a**, **b**, and **d** were determined by chiral HPLC; the analysis of **10c** was carried out at the stage of ester **12c**. The ee values, as expected, correspond well with those of the starting homoaldol products **7**. For safe assignments, the racemates *rac*-**10** and *rac*-**12** were synthesised from *rac*-**7** and analysed (see Experimental Section). The high diastereoselectivity originates from the (*E*)-oxonium intermediate **8**, being highly favoured over the (*Z*) diastereomer due to allylic 1,3-strain,^[9] which combines with the enolic double bond with high selectivity from the less shielded upper face and leads to a *cis*-annulation.

**Figure 2.** Solid-state structure of **12c**.^[10]

For the sake of obtaining shelf-stable products, the aldehydes **10** were reduced by LiAlH₄ to produce the alcohols **11** (Table 1) and these converted to *p*-nitrobenzoates **12** in order to enhance the chance for crystalline compounds. Fortunately, ester **12c** gave suitable crystals for an X-ray analysis, applying anomalous dispersion; it proves the assumed relative and absolute configurations (Figure 2).^[10]

Conclusion

Overall, the allyl carbamate **5** was successfully applied as a chiral 1-formylcycloalkane 2,3-dianion **C** which as-



- (a) $R^3CH=O$ (1.2 equivs.)/ $BF_3 \cdot OEt_2$ (1.2 equivs.), 1 h at $0^\circ C$
 (b) $LiAlH_4$ (1 M solution in Et_2O), 30 min at $0^\circ C$
 (c) *p*-Nitrobenzoyl chloride/DMAP, 15 h at room temperature

7	n	R ¹	R ²	R ³ -CH=O
A	1	Bu	4-BrC ₆ H ₄	4-BrC ₆ H ₄ CH=O
B	1	Bu	C ₆ H ₅	4-BrC ₆ H ₄ CH=O
C	1	Me	4-BrC ₆ H ₄	(<i>E</i>)-CH ₃ CH=CH-CH=O
D	0	Me	4-BrC ₆ H ₄	(CH ₃) ₂ CH-CH=O

Scheme 2. Synthesis of diastereomerically pure tetrahydrofurans **10**.

sembles two different or equal aldehydes **A** and **B** to form the bicyclic species **10** in a highly regio- and stereo-selective manner (see Figure 1). The expected relative and absolute configurations of compounds were **10** established vigorously by X-ray analysis. The configuration of all four consecutive stereogenic centres in **10** arises, directly or indirectly, from the chiral induction in the (–)-sparteine-mediated deprotonation step.

Experimental Section

General Remarks

All reactions, which are sensitive to moisture or air, were carried out under Ar using septum and syringe techniques. All solvents were purified by distillation or dried (Et_2O , CH_2Cl_2) prior to use. Flash chromatography was carried out with silica gel (40–63 μm) using an Ar pressure of 1.2–1.4 bar. Chiral HPLC was carried out with a chiral column chiragrom-1 and chiragrom-2, 60 $\times$ 2 mm, purchased from Grom Analytic and HPLC GmbH, Herrenberg. The solvent systems used for the measurement were hexane:*i*-PrOH (100:1) and (200:1). 1H and ^{13}C NMR spectra were recorded on ARX 300 and AMX 400 Bruker spectrometers. 2D NMR experiments were carried out on a Varian Unity Plus 600. $CDCl_3$ was used as solvent for

normal NMR measurements, IR absorption spectra were recorded using an IFS 28 purchased from Bruker and a PE 298 purchased from Perkin-Elmer & Co GmbH, Überlingen. The melting point was measured on an SMP3 melting point apparatus purchased from Stuart Scientific, UK. The optical rotations were measured in a 10-cm cuvette on a polarimeter 241 purchased from Perkin-Elmer & Co GmbH, Überlingen. Elemental analysis were performed at the Microanalytical Section of the Organisch-Chemisches Institut, WWU Münster, on a Vario El III purchased from Elementar Analysen Systeme GmbH. Exact mass measurements were carried out on Micro ToF (Bruker Daltonics, Bremen), calibrations were done directly before the measurements of samples with sodium formate clusters.

General Procedure for Synthesis of Aldehydes **10** (GP1)

To a cooled ($0^\circ C$) solution of **7** (0.50 mmol) and aldehyde $R^3CH=O$ in CH_2Cl_2 (5 mL) under an argon atmosphere, $BF_3 \cdot OEt_2$ (0.60 mmol) was added in small portions with a syringe. After stirring the reaction mixture for a period of 1 h at $0^\circ C$ and 30 min at $20^\circ C$, a saturated aqueous solution of $NaHCO_3$ (10 mL) was added and the mixture was extracted with Et_2O (3×20 mL). The combined ether solutions were dried with $MgSO_4$ and concentrated under vacuum. The crude residue was purified by silica gel flash chromatography (petroleum ether: Et_2O = 4:1). Correct microanalyses were obtained at the stage of corresponding *p*-nitrobenzoates **12**.

(1*S*,6*S*,7*R*,9*S*)-7,9-Di(4-bromophenyl)-6-butyl-8-oxabicyclo[4.3.0]nonane-1-carboxaldehyde (**10a**): According to GP1, **10a** was obtained from allyl carbamate **7a** (240 mg, 0.50 mmol) and *p*-bromobenzaldehyde (111 mg, 0.60 mmol) as a colourless oil; yield: 143 mg (55%); R_F = 0.71 (petroleum ether: Et_2O = 4:1); $[\alpha]_D^{20}$: -82.1 (c = 0.61, $CHCl_3$). Chiral HPLC: chiragrom-2, hexane:*i*-PrOH = 200:1, 78% ee, major enantiomer appears at higher retention time.

rac-**10a** was obtained from allyl carbamate *rac*-**7a** (240 mg, 0.50 mmol) and *p*-bromobenzaldehyde (111 mg, 0.60 mmol) as a colourless oil; yield: 137 mg (53%); 1H NMR (300 MHz, $CDCl_3$): δ = 0.84 (t, 3H, CH_3), 1.13–2.16 (m, 14H, CH_2), 4.57 (s, 1H, CH), 5.03 (s, 1H, CH), 7.08–7.49 (m, 8H, CH), 9.54 (s, 1H, CHO); ^{13}C NMR (75 MHz, $CDCl_3$): δ = 12.8 (CH_3), 20.8 (CH_2), 21.9 (CH_2), 23.0 (CH_2), 26.6 (CH_2), 27.0 (CH_2), 28.6 (CH_2), 30.4 (CH_2), 46.9 (C_q), 61.9 (C_q), 80.8 (CH), 84.8 (CH), 100.9/124.2/125.2/126.3/126.8/127.3/128.0/130.7 (CH and C_q of aryl carbons), 206.4 (C=O); IR (film): ν = 2958, 2951, 2853 (C_{aliph} -H), 1700 cm^{-1} (C=O).

(1*S*,6*S*,7*R*,9*S*)-9-(4-Bromophenyl)-6-butyl-7-phenyl-8-oxabicyclo[4.3.0]nonane-1-carboxaldehyde (**10b**): According to GP1, **10b** was obtained from allyl carbamate **7b** (193 mg, 0.50 mmol) and *p*-bromobenzaldehyde (111 mg, 0.60 mmol) as a colourless oil; yield: 119 mg (54%); R_F = 0.70 (petroleum ether: Et_2O = 4:1); $[\alpha]_D^{20}$: -59.3 (c 0.61, $CHCl_3$). Chiral HPLC: chiragrom-2, hexane:*i*-PrOH = 200:1, e.r. = 91.5:9.5 (81% ee), major enantiomer appears at lower retention time.

rac-**10b** was obtained from allyl carbamate *rac*-**7b** (193 mg, 0.5 mmol) and *p*-bromobenzaldehyde (111 mg, 0.6 mmol) as a colourless oil; yield: 121 mg (55%); 1H NMR (300 MHz, $CDCl_3$): δ = 0.95 (t, 3H, CH_3), 1.13–1.86 (m, 14H, CH_2), 4.56 (s, 1H, CH), 5.02 (s, 1H, CH), 7.11–7.46 (m, 9H, CH), 9.53 (s,

1H, CHO); ^{13}C NMR (75 MHz, CDCl_3): δ = 12.3 (CH_3), 20.4 (CH_2), 21.4 (CH_2), 23.3 (CH_2), 28.6 (CH_2), 30.0 (CH_2), 32.1 (CH_2), 33.1 (CH_2), 46.5 (C_q), 60.9 (C_q), 79.8 (CH), 83.8 (CH), 110.9/124.2/125.2/126.3/127.2/128.0/130.7 (CH and C_q of aryl carbons), 206.1 ($\text{C}=\text{O}$); IR (film): ν = 2958, 2916, 2860 ($\text{C}_{\text{aliph}}-\text{H}$), 1721 cm^{-1} ($\text{C}=\text{O}$).

(1S,6S,7R,9S)-7-(4-Bromophenyl)-6-methyl-9-[(E)-prop-1-enyl]-8-oxabicyclo[4.3.0]nonane-1-carboxaldehyde (**10c**): According to GP1, **10c** was obtained from allyl carbamate **7c** (123 mg, 0.28 mmol) and (E)-2-butenal (28 mg, 0.40 mmol) as a colourless oil; yield: 63 mg (62%); R_F = 0.70 (petroleum ether:Et₂O = 4:1); $[\alpha]_D^{20}$: -94.2 (c 0.50, CHCl_3). Chiral HPLC: chiragrom-1, hexane:i-PrOH = 200:1, 91% ee, major enantiomer appears at higher retention time.

rac-**10c** was obtained from allyl carbamate **7c** (123 mg, 0.28 mmol) and (E)-2-butenal (28 mg, 0.40 mmol) as a colourless oil; yield: 66 mg (65%); ^1H NMR (300 MHz, CDCl_3): δ = 1.18 (s, 3H, CH_3), 1.33–2.16 (m, 11H, CH_2 and CH_3), 4.45 (d, 1H, CH), 4.97 (s, 1H, CH), 5.55 (m, 1H, CH), 5.81 (m, 1H, CH), 7.18 (m, 2H, CH), 7.39 (m, 2H, CH), 9.64 (s, 1H, CHO); ^{13}C NMR (75 MHz, CDCl_3): δ = 17.8 (CH_3), 19.8 (CH_3), 21.2 (CH_2), 21.4 (CH_2), 26.8 (CH_2), 36.1 (CH_2), 47.7 (C_q), 61.2 (C_q), 85.2 (CH), 86.8 (CH), 121.6 (C_q), 127.3 (CH), 127.8 (CH), 131.3 (CH), 132.2 (CH), 138.4 (C_q), 203.3 ($\text{C}=\text{O}$); IR (film): ν = 2958, 2920, 2853 ($\text{C}_{\text{aliph}}-\text{H}$), 1719 cm^{-1} ($\text{C}=\text{O}$).

(1S,2S,4R,5S)-4-(4-Bromophenyl)-5-methyl-2-(1-methyl-ethyl)-3-oxabicyclo[3.3.0]octane-1-carboxaldehyde (**10d**): According to GP1, **10d** was obtained from allyl carbamate **7d** (106 mg, 0.25 mmol) and 2-methylpropanal (29 mg, 0.40 mmol) as a colourless oil; yield: 55 mg (63%); R_F = 0.71 (petroleum ether:Et₂O = 4:1); $[\alpha]_D^{20}$: -119.1 (c 0.51, CHCl_3). Chiral HPLC: chiragrom-2, hexane:i-PrOH = 200:1, 96% ee, major enantiomer appears at lower retention time.

rac-**10d** was obtained from allyl carbamate *rac*-**7d** (106 mg, 0.25 mmol) and 2-methylpropanal (29 mg, 0.40 mmol) as a colourless oil; yield: 54 mg (61%); ^1H NMR (300 MHz, CDCl_3): δ = 0.95 (s, 3H, CH_3), 1.02 (d, 6H, CH_3), 1.35–2.19 (m, 6H, CH_2), 2.15 (m, 1H, CH), 4.14 (d, J = 6.5 Hz, 1H, CH), 4.43 (s, 1H, CH), 7.14 (m, 2H, CH), 7.34 (m, 2H, CH), 9.59 (s, 1H, CHO); ^{13}C NMR (75 MHz, CDCl_3): δ = 17.8 (CH_3), 19.5 (CH_3), 19.6 (CH_3), 21.2 (CH_2), 21.5 (CH_2), 23.6 (CH_2), 27.3 (CH), 47.7 (C_q), 61.5 (C_q), 75.5 (CH), 86.3 (CH), 121.4 (C_q), 131.4 (CH), 132.5 (CH), 138.9 (C_q), 203.5 ($\text{C}=\text{O}$); IR (film): ν = 2958, 2951, 2853 ($\text{C}_{\text{aliph}}-\text{H}$), 1721 cm^{-1} ($\text{C}=\text{O}$).

General Procedure for Synthesis of Alcohols 11 (GP2)

To a solution of aldehyde **10** (0.25 mmol) in Et₂O (5 mL), LiAlH₄ (1 M solution in Et₂O; 0.25 mmol) was added in small portions at 0 °C. After stirring at 0 °C for 30 min and subsequently, at room temperature for 3 h, Et₂O (20 mL) and a saturated aqueous solution of NH₄Cl (10 mL) were added. The aqueous phase was extracted (Et₂O, 2 × 20 mL), the combined solutions were dried over anhydrous MgSO₄ and concentrated under vacuum. The residue was purified by silica gel flash column chromatography (petroleum ether:Et₂O = 4:1).

(1R,6S,7R,9S)-{7,9-Di(4-bromophenyl)-6-butyl-8-oxabicyclo[4.3.0]non-1-yl}-methanol (**11a**): According to GP2, **11a** was obtained from **10a** (130 mg, 0.25 mmol) as a colourless oil; yield: 91 mg (70%); R_F = 0.18 (petroleum ether:Et₂O = 4:1); $[\alpha]_D^{20}$: -36.5 (c 0.65, CHCl_3). Chiral HPLC: chiragrom-2, hex-

ane:i-PrOH = 200:1, 78% ee, major enantiomer appears at higher retention time.

rac-**11a** was obtained from **10a** (130 mg, 0.25 mmol) as a colourless oil; yield: 94 mg (72%); ^1H NMR (300 MHz, CDCl_3): δ = 0.93 (t, 3H, CH_3), 1.16–1.86 (m, 14H, CH_2), 3.54 (d, 1H, J = 11.7 Hz, CH), 3.69 (d, J = 11.6 Hz, 1H, CH), 4.54 (s, 1H, CH), 4.58 (s, 1H, CH), 7.12–7.40 (m, 8H, CH); ^{13}C NMR (75 MHz, CDCl_3): δ = 13.0 (CH_3), 20.4 (CH_2), 21.8 (CH_2), 23.7 (CH_2), 28.5 (CH_2), 30.3 (CH_2), 32.7 (CH_2), 33.9 (CH_2), 46.8 (C_q), 49.4 (C_q), 63.9 (CH_2), 79.6 (CH), 83.8 (CH), 111.5/124.4/125.8/126.6/127.3/128.2/130.4/130.5 (CH and C_q of aryl carbons); IR (film): ν = 3370 (OH), 2958, 2930, 2867 cm^{-1} ($\text{C}_{\text{aliph}}-\text{H}$).

(1R,6S,7R,9S)-{9-(4-Bromophenyl)-6-butyl-7-phenyl-8-oxabicyclo[4.3.0]non-1-yl}-methanol (**11b**): According to GP2, **11b** was obtained from **10b** (110 mg, 0.25 mmol) as a colourless oil; yield: 80 mg (72%); R_F = 0.18 (petroleum ether:Et₂O = 4:1); $[\alpha]_D^{20}$: +30.1 (c 0.48, CHCl_3). Chiral HPLC: chiragrom-2, hexane:i-PrOH = 200:1, 81% ee, major enantiomer appears at higher retention time.

rac-**11b** was obtained from **10b** (110 mg, 0.25 mmol) as a colourless oil; yield: 77 mg (70%); ^1H NMR (300 MHz, CDCl_3): δ = 0.93 (t, 3H, CH_3), 1.14–1.86 (m, 14H, CH_2), 3.58 (d, 1H, J = 11.7 Hz, CH), 3.71 (d, J = 11.8 Hz, 1H, CH), 4.84 (s, 1H, CH), 4.99 (s, 1H, CH), 7.11–7.28 (m, 9H, CH); ^{13}C NMR (75 MHz, CDCl_3): δ = 12.7 (CH_3), 21.2 (CH_2), 21.9 (CH_2), 23.8 (CH_2), 26.2 (CH_2), 29.6 (CH_2), 31.7 (CH_2), 33.5 (CH_2), 45.4 (C_q), 49.9 (C_q), 64.6 (CH_2), 79.9 (CH), 83.6 (CH), 111.9/124.4/125.7/126.3/127.2/127.6/128.9/130.5 (CH and C_q of aryl carbons); IR (film): ν = 3426 (OH), 2979, 2923, 2853 cm^{-1} ($\text{C}_{\text{aliph}}-\text{H}$).

(1R,6S,7R,9S)-{7-(4-Bromophenyl)-6-methyl-9-[(E)-prop-1-enyl]-8-oxabicyclo[4.3.0]non-1-yl}-methanol (**11c**): According to GP2, **11c** was obtained from **10c** (61 mg, 0.17 mmol) as a colourless oil; yield: 45 mg (73%); R_F = 0.16 (petroleum ether:Et₂O = 4:1); $[\alpha]_D^{20}$: -80.0 (c 0.25, CHCl_3). Chiral HPLC: chiragrom-1, hexane:i-PrOH = 200:1, 91% ee, major enantiomer appears at higher retention time.

rac-**11c** was obtained from **10c** (61 mg, 0.17 mmol) as a colourless oil; yield: 45 mg (74%); ^1H NMR (300 MHz, CDCl_3): δ = 0.40 (s, 3H, CH_3), 1.16–2.26 (m, 11H, CH_2 and CH_3), 3.56 (d, 1H, J = 11.7 Hz, CH), 3.73 (d, J = 11.7 Hz, 1H, CH), 4.06 (m, 1H, CH), 4.55 (s, 1H, CH), 5.77 (m, 2H, CH), 7.11 (m, 2H, CH), 7.38 (m, 2H, CH); ^{13}C NMR (75 MHz, CDCl_3): δ = 17.8 (CH_3), 19.8 (CH_3), 21.8 (CH_2), 21.9 (CH_2), 30.5 (CH_2), 32.1 (CH_2), 45.6 (C_q), 50.5 (C_q), 64.3 (CH_2), 83.0 (CH), 88.5 (CH), 120.8 (C_q), 128.2/128.4/129.9/130.7/131.3/131.5 (CH and C_q), 138.0 (C_q); IR (film): ν = 3372 (OH), 2956, 2932, 2869 cm^{-1} ($\text{C}_{\text{aliph}}-\text{H}$).

(1R,2S,4R,5S)-{4-(4-Bromophenyl)-5-methyl-2-(1-methyl-ethyl)-3-oxabicyclo[3.3.0]oct-1-yl}-methanol (**11d**): According to GP2, **11d** was obtained from **10d** (50 mg, 0.14 mmol) as a colourless oil; yield: 37 mg (74%); R_F = 0.16 (petroleum ether:Et₂O = 4:1); $[\alpha]_D^{20}$: -42.7 (c 0.55, CHCl_3). Chiral HPLC: chiragrom-2, hexane:i-PrOH = 200:1, 96% ee, major enantiomer appears at lower retention time.

rac-**11d** was obtained from **10d** (50 mg, 0.14 mmol) as a colourless oil; yield: 38 mg (75%); ^1H NMR (300 MHz, CDCl_3): δ = 0.43 (s, 3H, CH_3), 1.04 (d, 6H, CH_3), 1.33–2.16 (m, 6H, CH_2), 2.10 (m, 1H, CH), 3.53 (d, J = 11.7 Hz, 1H, CH), 3.75 (d, J = 1.7 Hz, 1H, CH), 4.13 (d, 1H, CH), 4.46 (s, 1H, CH), 7.13–7.42 (m, 4H, CH); ^{13}C NMR (75 MHz, CDCl_3): δ = 17.4

(CH₃), 19.8 (CH₃), 21.3 (CH₂), 21.6 (CH₂), 32.8 (CH₂), 27.7 (CH), 47.5 (C_q), 61.6 (C_q), 64.7 (CH₂), 73.1 (CH), 88.7 (CH), 128.3 (C_q), 129.6 (CH), 131.3 (CH), 138.2 (C_q); IR (film): ν = 3372 (OH), 2954, 2931, 2864 cm⁻¹ (C_{aliph}-H); anal. calcd. for C₁₈H₂₅O₂Br (352.29): C 61.19, H 7.13; found: C 61.44, H 7.35.

General Procedure for *p*-Nitrobenzoates **12** (GP3)

A solution of alcohol **11** (0.20 mmol), *p*-nitrobenzoyl chloride (0.30 mmol) and DMAP (0.08 mmol) in CH₂Cl₂ (5 mL) was stirred for 15 h at room temperature. Aqueous saturated solution of NaHCO₃ (10 mL) and Et₂O (20 mL) were added, the phases separated and the aqueous phase extracted twice with Et₂O (20 mL, each). The combined organic extracts were concentrated under vacuum and the residue purified by flash column chromatography on silica gel (petroleum ether: Et₂O = 4:1).

(1*R*,6*S*,7*R*,9*S*)-[7,9-Di(4-bromophenyl)-6-butyl-8-oxabicyclo[4.3.0]non-1-yl]methyl 4-Nitrobenzoate (**12a**): According to GP3, **12a** was obtained from **11a** (90 mg, 0.17 mmol) as a colourless solid; yield: 91 mg (79%); R_F = 0.30 (petroleum ether: Et₂O = 4:1); [α]_D²⁰: -41.1 (c 0.59, CHCl₃). Chiral HPLC: chiragrom-2, hexane:*i*-PrOH = 200:1, 78% ee, major enantiomer appears at lower retention time.

rac-**12a** was obtained from **11a** (90 mg, 0.17 mmol) as a colourless solid; yield: 94 mg (81%); ¹H NMR (300 MHz, CDCl₃): δ = 0.94 (t, 3H, CH₃), 1.16–1.64 (m, 14H, CH₂), 3.66 (d, J = 11.5 Hz, 1H, CH), 3.72 (d, J = 11.6 Hz, 1H, CH), 4.34 (s, 1H, CH), 5.19 (s, 1H, CH), 7.19–7.36 (m, 8H, CH), 8.13–8.24 (m, 4H, CH); ¹³C NMR (75 MHz, CDCl₃): δ = 14.2 (CH₃), 21.0 (CH₂), 21.6 (CH₂), 26.2 (CH₂), 28.8 (CH₂), 29.4 (CH₂), 30.8 (CH₂), 31.5 (CH₂), 52.3 (C_q), 56.6 (C_q), 64.8 (CH₂), 80.3 (CH), 83.2 (CH), 121.5/123.7/125.6/126.2/127.2/127.6/129.4/130.9/130.7/131.8/132.3/135.7 (CH and C_q of aryl carbons), 166.0 (C=O); IR (KBr): ν = 3021, 2958, 2853 (C_{aliph}-H), 1703 cm⁻¹ (C=O); anal. calcd. for C₃₂H₃₃Br₂NO₅ (671.42): C 57.24, H 4.95, N 2.09; found: C 57.52, H 4.80, N 2.19.

(1*R*,6*S*,7*R*,9*S*)-[9-(4-Bromophenyl)-6-butyl-7-phenyl-8-oxabicyclo[4.3.0]non-1-yl]methyl 4-Nitrobenzoate (**12b**): According to GP3, **12b** was obtained from **11b** (75 mg, 0.17 mmol) as a colourless oil; yield: 81 mg (81%); R_F = 0.30 (petroleum ether: Et₂O = 4:1); [α]_D²⁰: -72.2 (c 0.52, CHCl₃). Chiral HPLC: chiragrom-2, hexane:*i*-PrOH = 200:1, 81% ee, major enantiomer appears at higher retention time.

rac-**12b** was obtained from **11b** (75 mg, 0.17 mmol) as a colourless oil; yield: 80 mg (80%); ¹H NMR (300 MHz, CDCl₃): δ = 0.98 (t, 3H, CH₃), 1.16–1.66 (m, 14H, CH₂), 3.53 (d, J = 11.6 Hz, 1H, CH), 3.72 (d, J = 11.7 Hz, 1H, CH), 4.98 (s, 1H, CH), 5.11 (s, 1H, CH), 7.11–7.28 (m, 9H, CH), 8.12–8.27 (m, 4H, CH); ¹³C NMR (75 MHz, CDCl₃): δ = 14.2 (CH₃), 21.0 (CH₂), 21.8 (CH₂), 26.4 (CH₂), 26.2 (CH₂), 26.4 (CH₂), 28.6 (CH₂), 32.0 (CH₂), 52.1 (C_q), 56.4 (C_q), 64.8 (CH₂), 80.6 (CH), 83.8 (CH), 122.2/123.4/124.5/125.9/126.8/127.9/129.2/130.4/130.6/131.5/132.4/135.6, (CH and C_q of aryl carbons), 166.0 (C=O); IR (KBr): ν = 3014, 2951, 2853 (C_{aliph}-H), 1707 cm⁻¹ (C=O); anal. calcd. for C₃₂H₃₄NO₅Br (592.52): C 64.87, H 5.78, N 2.36; found: C 64.60, H 5.56, N 2.31.

(1*R*,6*S*,7*R*,9*S*)-[7-(4-Bromophenyl)-6-methyl-9-[(*E*)-propenyl]-8-oxabicyclo[4.3.0]non-1-yl]methyl 4-Nitrobenzoate (**12c**): According to GP3, **12c** was obtained from **11c** (40 mg, 0.11 mmol) as a colourless solid; yield: 47 mg (84%); R_F =

0.30 (petroleum ether: Et₂O = 4:1), mp 214.5 °C (petroleum ether: Et₂O = 4:1); [α]_D²⁰: -62.2 (c 0.59, CHCl₃). Chiral HPLC: chiragrom-2, hexane:*i*-PrOH = 200:1, 91% ee, major enantiomer appears at higher retention time.

rac-**12c** was obtained from **11c** (40 mg, 0.11 mmol) as a colourless solid; yield: 48 mg (85%); ¹H NMR (300 MHz, CDCl₃): δ = 0.53 (s, 3H, CH₃), 1.06–2.15 (m, 11H, CH₂ and CH₃), 3.52 (d, J = 11.5 Hz, 1H, CH), 3.72 (d, J = 11.5 Hz, 1H, CH), 4.10 (m, 1H, CH), 4.58 (s, 1H, CH), 5.78 (m, 1H, CH), 5.58 (m, 1H, CH), 7.11 (m, 2H, CH), 7.39 (m, 2H, CH), 8.13 (m, 2H, CH), 8.24 (m, 2H, CH); ¹³C NMR (75 MHz, CDCl₃): δ = 17.9 (CH₃), 20.2 (CH₃), 21.8 (CH₂), 22.12 (CH₂), 31.8 (CH₂), 32.5 (CH₂), 46.1 (C_q), 49.1 (C_q), 64.6 (CH₂), 83.3 (CH), 88.2 (CH), 121.5 (C_q), 123.7 (CH), 128.4 (CH), 129.6 (CH), 130.3 (CH), 130.6 (CH), 131.0 (CH), 135.5 (C_q), 137.8 (C_q), 150.7 (C_q), 164.5 (C=O); IR (KBr): ν = 3024, 2954, 2857 (C_{aliph}-H), 1708 cm⁻¹ (C=O); anal. calcd. for C₂₆H₂₈BrNO₅ (514.41): C 60.71, H 5.49, N 2.72; found: C 60.99, H 5.39, N 2.67.

(1*R*,2*S*,4*R*,5*S*)-[4-(4-Bromophenyl)-5-methyl-2-(1-methyl-ethyl)-3-oxabicyclo[3.3.0]oct-1-yl]methyl 4-Nitrobenzoate (**12d**): According to GP3, **12d** was obtained from **11d** (35 mg, 0.10 mmol) as a colourless oil; yield: 30 mg (85%) R_F = 0.29 (petroleum ether: Et₂O = 4:1); [α]_D²⁰: -111.1 (c 0.50, CHCl₃). Chiral HPLC: chiragrom-1, hexane:*i*-PrOH = 200:1, 96% ee, major enantiomer appears at higher retention time.

rac-**12d** was obtained from **11d** (35 mg, 0.10 mmol) as a colourless oil; yield: 29 mg (83%); ¹H NMR (300 MHz, CDCl₃): δ = 0.56 (s, 3H, CH₃), 1.03 (d, 6H, CH₃), 1.10–2.16 (m, 6H, CH₂), 2.13 (m, 1H, CH), 3.53 (d, J = 11.7 Hz, 1H, CH), 3.75 (d, J = 11.7 Hz, 1H, CH), 4.56 (d, 1H, CH), 5.04 (s, 1H, CH), 7.17 (m, 2H, CH), 7.49 (m, 2H, CH), 8.12 (m, 2H, CH), 8.25 (m, 2H, CH); ¹³C NMR (75 MHz, CDCl₃): δ = 17.5 (CH₃), 19.2 (CH₃), 21.1 (CH₂), 21.2 (CH₂), 23.8 (CH₂), 27.2 (CH), 47.4 (C_q), 61.3 (C_q), 64.7 (CH₂), 73.4 (CH), 88.9 (CH), 121.1 (C_q), 123.3 (C_q), 128.6 (CH), 130.3 (CH), 130.2 (CH), 131.4 (CH), 137.3 (C_q), 150.8 (C_q), 164.2 (C=O); IR (film): ν = 3023, 2962, 2852 (C_{aliph}-H), 1706 cm⁻¹ (C=O); anal. calcd. for C₂₅H₂₉BrO₅N (502.40): C 59.77, H 5.62, N 2.79; found: C 60.09, H 5.71, N 2.86; exact mass for C₂₅H₃₀BrO₅N: calcd.: 502.1236; found: 502.1230.

X-Ray Crystallographic Study^[10]

X-ray crystal structure analysis HOP1432: formula C₂₆H₂₈NO₅Br, M = 514.40, colourless crystal 0.40 × 0.35 × 0.25 mm, a = 10.353(3), b = 12.682(2), c = 18.842(5) Å, a = 98.57(1), β = 93.55(1), γ = 102.88(1)°, V = 2373.1(10) Å³, ρ_{calc} = 1.440 g cm⁻³, μ = 17.70 cm⁻¹, empirical absorption correction (0.538 ≤ T ≤ 0.666), Z = 4, triclinic, space group $P\bar{1}$ (No. 2), R = 0.071073 Å, T = 198 K, ω and ϕ scans, 15773 reflections collected ($\pm h$, $\pm k$, $\pm l$), [(sin θ)/ λ] = 0.59 E⁻¹, 7394 independent (R_{int} = 0.075) and 3613 observed reflections [$I \geq 2 \sigma(I)$], 599 refined parameters, R = 0.068, wR^2 = 0.168, max. residual electron density 0.43 (−0.42) e Å⁻³, contains two almost identical molecules in the asymmetric unit, hydrogen atoms calculated and refined as riding atoms.

Acknowledgements

This work was supported by the Deutsche Forschungsgemeinschaft, (SFB 424) and the Fonds der Chemischen Industrie.

References and Notes

- [1] Review: X. L. Hou, Z. Yang, H. N. C. Wong, *Progress in Heterocyclic Chemistry* **2002**, *14*, 139–179.
- [2] Review: T. L. B. Boivin, *Tetrahedron* **1997**, *43*, 3309–3362.
- [3] a) D. Hoppe, T. Krämer, C. Freire Erdrügger, E. Egert, *Tetrahedron Lett.* **1989**, *30*, 1233–1236; b) H. Paulsen, C. Graeve, D. Hoppe, *Synthesis* **1996**, 141–144.
- [4] Reviews: a) D. Hoppe, T. Hense, *Angew. Chem.* **1997**, *109*, 2376–2410; *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 2282–2316; b) D. Hoppe, F. Marr, M. Brüggemann, *Organolithiums in Enantioselective Synthesis, Topics in Organometallic Chemistry*, Vol 5, (Ed.: D. M. Hodgson), Springer-Verlag, Berlin, **2003**, *5*, 61–138; c) D. Hoppe, G. Christoph, *The Chemistry of Organolithium Compounds*, Vol 2, (Eds.: Z. Rappoport, I. Marek), John Wiley & Sons, Chichester, **2004**, 1055–1164.
- [5] a) T. Mukaiyama, S. Kobayashi, *Organic Reactions* **1994**, *46*, 1–104; b) T. Mukaiyama, *Tetrahedron* **1999**, *55*, 8609–8670; c) T. Mukaiyama, *Angew. Chem. Int. Ed. Engl.* **2004**, *43*, 5590–5614; *Angew. Chem. Engl.* **2004**, *116*, 5708–5733.
- [6] For related approaches see: a) R. W. Hoffmann, V. Giesen, M. Fuest, *Liebigs Ann. Chem.* **1993**, 629–639; b) P. Mohr, *Tetrahedron Lett.* **1993**, *34*, 6251–6254; c) N. Hakanı, J. T. Link, D. W. C. McMillan, L. E. Overman, W. G. Trankle, J. A. Wurster, *Org. Lett.* **2000**, *2*, 223–226.
- [7] a) M. Özlügedik, J. Kristensen, B. Wibbeling, R. Fröhlich, D. Hoppe, *Eur. J. Org. Chem.* **2002**, 414–427, b) M. Özlügedik, J. Kristensen, J. Reuber, R. Fröhlich, D. Hoppe, *Synthesis* **2004**, *14*, 2303–2316; c) S. Ünalı, R. Fröhlich, D. Hoppe, *Synthesis* **2005**, in press.
- [8] *p*-Bromobenzaldehyde in most cases was used for R²CH=O or R³CH=O in order to improve the crystallization tendency and for introduction of a heavy atom for applying anomalous X-ray dispersion.
- [9] Review: R. W. Hoffmann, *Chem. Rev.* **1989**, *89*, 1841–1860.
- [10] Data set was collected with a Nonius Kappa CCD diffractometer, equipped with a rotating anode generator. Programs used: data collection COLLECT (Nonius B. V., 1998), data reduction Denzo-SMN (Z. Otwinowski, W. Minor, *Methods in Enzymology*, **1997**, *276*, 307–326), absorption correction SORTAV (R. H. Blessing, *Acta Cryst.* **1995**, *A51*, 33–37; R. H. Blessing, *J. Appl. Cryst.* **1997**, *30*, 421–426), structure solution SHELXS-97 (G. M. Sheldrick, *Acta Cryst.* **1990**, *A46*, 467–473), structure refinement SHELXL-97 (G. M. Sheldrick, Universität Göttingen, **1997**), graphics MO-PICT 3.2 (M. Brüggemann, Universität Münster, **2001**). CCDC 182785 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge at www.ccdc.cam.ac.uk/conts/retrieving.html [or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: (internat.) +44(1223)336–033, E-mail: deposit@ccdc.cam.ac.uk].