

C₃H₅N or H₂O), $M_r = 7084.54$, cubic, $Fm\bar{3}m$, $a = b = c = 26.5200(13)$ Å, $V = 18651.8(16)$ Å³, $Z = 2$, $\rho_{\text{calcd}} = 1.261$ g cm⁻³, $2\theta_{\text{max}} = 50.0^\circ$ ($-31 \leq h \leq 31$, $-31 \leq k \leq 18$, $-28 \leq l \leq 31$), $T = 173$ K, 22842 measured reflections, $RI = 0.0602$ for 670 reflections ($I > 2\sigma(I)$), $wR2 = 0.1840$ for 891 independent reflections (all data) and 63 parameters, $GOF = 0.990$.

Data were collected on a Bruker SMART-APEX CCD diffractometer using MoK α radiation ($\lambda = 0.71073$ Å), operating in the Ω and φ scan mode. All crystal data were corrected for Lorentz and polarization effects, and the SADABS^[25] program was used for absorption correction. The structures were solved by direct methods and the structure solutions and refinements were based on $|F^2|$. All non-hydrogen atoms were refined with anisotropic displacement parameters, whereas hydrogen atoms were placed in calculated positions and given isotropic U values 20% higher than the atom to which they are bonded. All crystallographic calculations were conducted with the SHELXTL^[26] software suite. Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication nos. CCDC-156776 (1) and CCDC-156777 (2). Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB21EZ, UK (fax: (+44)1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).

Received: January 25, 2001 [Z16502]

- [1] A. Kaur Sra, M. Andruh, O. Kahn, S. Golhen, L. Ouahab, J. V. Yakhmi, *Angew. Chem.* **1999**, *111*, 2768–2771; *Angew. Chem. Int. Ed.* **1999**, *38*, 2606–2609.
- [2] X. X. Zhang, S. S. Y. Chui, I. D. Williams, *J. Appl. Phys.* **2000**, *87*, 6007–6009.
- [3] S.-i. Noro, S. Kitagawa, M. Kondo, K. Seki, *Angew. Chem.* **2000**, *112*, 2161–2164; *Angew. Chem. Int. Ed.* **2000**, *39*, 2081–2084.
- [4] H. Li, M. Eddaoudi, M. O'Keefe, O. M. Yaghi, *Nature* **1999**, *402*, 276–279.
- [5] J. S. Seo, D. Whang, H. Lee, S. I. Jun, J. Oh, Y. J. Jeon, K. Kim, *Nature* **2000**, *404*, 982–986.
- [6] G. W. Orr, L. J. Barbour, J. L. Atwood, *Science* **1999**, *285*, 1049–1052.
- [7] L. R. MacGillivray, J. L. Atwood, *Nature* **1997**, *389*, 469–472.
- [8] L. R. MacGillivray, J. L. Atwood, *Angew. Chem.* **1999**, *111*, 1080–1096; *Angew. Chem. Int. Ed.* **1999**, *38*, 1018–1033.
- [9] S. Leininger, B. Olenyuk, P. J. Stang, *Chem. Rev.* **2000**, *100*, 853–907.
- [10] P. J. Stang, B. Olenyuk, *Acc. Chem. Res.* **1997**, *30*, 502–518.
- [11] N. Takeda, K. Umemoto, K. Yamaguchi, M. Fujita, *Nature* **1999**, *398*, 794–796.
- [12] M. Fujita, D. Oguro, M. Miyazawa, H. Oka, K. Yamaguchi, K. Ogura, *Nature* **1995**, *378*, 469–471.
- [13] K. Umemoto, K. Yamaguchi, M. Fujita, *J. Am. Chem. Soc.* **2000**, *122*, 7150–7151.
- [14] H. S. M. Coxeter, M. S. Longuet-Higgins, J. C. P. Miller, *Philos. Trans. R. Soc. London A* **1954**, *246*, 401–450.
- [15] M. J. Wenninger, *Polyhedron Models*, Cambridge University Press, Cambridge, **1989**.
- [16] A. Holden, *Shapes, Space, and Symmetry*, Columbia University Press, New York, **1971**.
- [17] O. M. Yaghi, H. L. Li, C. Davis, D. Richardson, T. L. Groy, *Acc. Chem. Res.* **1998**, *31*, 474–484.
- [18] M. Eddaoudi, H. L. Li, O. M. Yaghi, *J. Am. Chem. Soc.* **2000**, *122*, 1391–1397.
- [19] D. Wells, *The Penguin Dictionary of Curious and Interesting Geometry*, Penguin, London **1991**.
- [20] E. W. Weisstein, *The CRC Concise Encyclopedia of Mathematics*, CRC, Boca Raton, FL, **1999**.
- [21] A. Andreini, *Mem. Soc. Ital. Sci.* **1907**, *14*, 75.
- [22] A. F. Wells, *Three-dimensional Nets and Polyhedra*, Wiley, New York, **1977**.
- [23] S. S. Y. Chui, S. M. F. Lo, J. P. H. Charmant, A. G. Orpen, I. D. Williams, *Science* **1999**, *283*, 1148–1150.
- [24] S. J. Cantrill, A. R. Pease, J. F. Stoddart, *Dalton Trans.* **2000**, 3715–3734.
- [25] SADABS: Area-Detector Absorption Correction, Siemens Industrial Automation, Inc., Madison, WI, **1996**.
- [26] SHELXTL v5.1, Bruker-AXS, Inc., Madison, WI, **1998**.

A Rationally Designed NH₄⁺ Receptor Based on Cation– π Interaction and Hydrogen Bonding**

Sang Yong Jon, Jeongho Kim, Minkyung Kim, Sang-Hyun Park, Woo Sung Jeon, Jeongsuk Heo, and Kimoon Kim*

Design and synthesis of ammonium ion receptors have received much attention, not only because ammonium binding is one of the fundamental problems of molecular recognition, but also because such receptors serve as an essential component of sensors used in clinical^[1] and environmental analyses.^[2] Most of the NH₄⁺ receptors known to date harness hydrogen bonding in binding an ammonium ion.^[3] For example, nonactin (see Scheme 1), a natural antibiotic agent and one of the most effective NH₄⁺ receptors, utilizes four ethereal and four carbonyl oxygen atoms to bind NH₄⁺ through hydrogen bonding.^[4] While hydrogen bonding is most commonly involved in molecular recognition, cation– π interaction has drawn much attention in recent years.^[5] In fact, cation– π interaction has been successfully employed in designing synthetic receptors for alkylammonium ions and alkali metal ions.^[5, 6] Despite extensive theoretical studies on NH₄⁺– π interaction, however, no synthetic NH₄⁺ receptor exploiting cation– π interaction had been reported until recently. We recently reported a novel NH₄⁺ receptor based on tris(pyrazol-1-ylmethyl)benzene that is highly selective for binding NH₄⁺ over alkali metal ions.^[7] Subsequent theoretical studies on this receptor by us^[8] and others^[9] suggest that the cation– π interaction between the substrate and the central benzene ring of the receptor plays a significant role in the binding affinity and selectivity of the tripodal receptor toward NH₄⁺. Despite the high selectivity, however, the ion-selective electrode (ISE) based on this receptor shows low sensitivity and strong pH dependence of its performance in comparison to a nonactin-based ISE, and this hampers practical applications as a sensor. Here we present a rationally designed cage-type NH₄⁺ receptor that binds the cation by both cation– π interaction and hydrogen bonding, with high sensitivity and selectivity comparable or superior to those of nonactin over a wide range of pH.


In designing the receptor, we considered the following stereoelectronic features: 1) spatial distribution of lone-pair

[*] Prof. K. Kim, Dr. S. Y. Jon, J. Kim, M. Kim, Dr. S.-H. Park, W. S. Jeon, J. Heo
National Creative Research Initiative Center for Smart Supramolecules and Department of Chemistry
Division of Molecular and Life Sciences
Pohang University of Science and Technology (POSTECH)
San 31 Hyojadong, Pohang 790-784 (Republic of Korea)
Fax: (+82)54-279-8129
E-mail: kkim@postech.ac.kr

[**] We gratefully acknowledge the Korean Ministry of Science and Technology (Creative Research Initiative Program) for support of this work, and the Korean Ministry of Education (Brain Korea 21 program) for graduate studentships to J.H. and W.S.J. M.K. is grateful to POSTECH for an Undergraduate Research Fellowship. We also thank Professor P. K. Bharadwaj for helpful discussions.

 Supporting information for this article is available on the WWW under <http://www.angewandte.com> or from the author.

electrons for effective hydrogen bonding with the tetrahedral NH_4^+ ion, 2) benzene rings ideally positioned for strong cation- π interaction, and 3) a rigid framework with a cavity of correct size for binding NH_4^+ . With this in mind, we designed receptor **1** in which two benzene rings are rigidly held together by three dialkoxybenzene units (Scheme 1). We chose phenolic oxygen atoms as



Scheme 1. Chemical structures of the ammonium receptors discussed here.

hydrogen bonding acceptors because their $\text{p}K_a$ values are larger than those of nitrogen atoms, and thus we can lower the pH dependence of the receptor performance. Molecular mechanics calculations^[10] on **1** and its NH_4^+ complex showed that each dialkoxybenzene unit is almost ideally positioned to “bite” NH_4^+ through hydrogen bonding, while the two benzene rings are properly positioned at the top and bottom of the cage for cation- π interaction with NH_4^+ , with an $\text{N} \cdots$ benzene centroid distance of about 3.0 Å. In addition, the cavity size of **1** is perfect for NH_4^+ but too large for smaller ions such as Li^+ and Na^+ .

Receptor **1** is easily synthesized from the reaction of catechol with 1,3,5-tris(bromomethyl)-2,4,6-triethylbenzene^[11] in two steps in good yield. The X-ray crystal structures^[12] of **1** (see Supporting Information) and its NH_4^+ complex $[\mathbf{1}\text{-NH}_4]\text{PF}_6$ (Figure 1) are in good agreement with those calculated by molecular modeling. These structures reveal that the only significant structural change in the receptor associated with NH_4^+ binding is that the mean distance between the two benzene rings increases from 5.588 to 5.966 Å. In $[\mathbf{1}\text{-NH}_4]\text{PF}_6$, as expected, the NH_4^+ ion is located at the center of the receptor and forms hydrogen bonds with six oxygen atoms. A set of oxygen atoms (O2, O2A, and O2B) interacts with the NH_4^+ ion with $\text{O} \cdots \text{N}$ distances of 3.224(5) Å and $\text{N-H} \cdots \text{O}$ angles of 168(12)°, whereas the other set (O1, O1A, and O1B) has $\text{O} \cdots \text{N}$ distances of 3.226(5) Å and $\text{N-H} \cdots \text{O}$ angles of 122(7)°.^[13] The average distance between the nitrogen atom of NH_4^+ and the center of the top or bottom benzene ring is 2.985 Å, which is in good agreement with the optimal distance for cation- π interaction between NH_4^+ and benzene suggested by theoretical studies.^[5, 14] The PF_6^- counterion is located outside the cage with no apparent interactions.




Figure 1. a) Side and b) top views of $[\mathbf{1}\text{-NH}_4]\text{PF}_6$ in the solid state. Hydrogen atoms (except for those of NH_4^+) and the PF_6^- counterion are omitted for clarity. The ethyl groups of the “bottom” benzene ring are disordered over two sites (occupancy: C16 0.64, C16' 0.36). Selected interatomic distances [Å] and angles [°]: $\text{O1} \cdots \text{N1}$ 3.226(5), $\text{N1-H2} \cdots \text{O1}$ 122(7), $\text{O2} \cdots \text{N1}$ 3.224(5), $\text{N1-H2} \cdots \text{O2}$ 168(12), centroid of “top” benzene $\cdots \text{N1}$ 2.991, centroid of “bottom” benzene $\cdots \text{N1}$ 2.980.

The performance of an ISE based on **1**^[15] is comparable or even superior to that of a nonactin-based ISE in terms of sensitivity and selectivity (Figure 2): detection limit $3.2 \times 10^{-6} \text{ M}$ (**1**) versus $1.5 \times 10^{-6} \text{ M}$ (nonactin); selectivity coefficients $\lg K_{\text{NH}_4^+/\text{K}^+} = -0.97$ (**1**) versus -0.88 (nonactin), and $\lg K_{\text{NH}_4^+/\text{Na}^+} = -3.00$ (**1**) versus -2.85 (nonactin). Furthermore, it displays only a small pH dependence (see Supporting Information) which is also comparable to that of the nonactin-based ISE.




Figure 2. Responses of ISEs to NH_4^+ , K^+ , and Na^+ (background electrolyte: 0.05 M Tris-HCl, pH 7.2). a) Responses of an ISE based on **1**. b) Selectivity coefficients of ISEs based on nonactin, **1**, and **2**.

To understand the role of cation– π interaction in **1** in binding NH_4^+ , we synthesized the modified receptor **2**, which does not have the “top” benzene ring (see Scheme 1). As expected, the binding affinity of **1** toward NH_4^+ is at least ten times higher than that of **2**: the association constants for binding of NH_4^+ to **1** and **2**, determined by the extraction method,^[16] are 3.3×10^7 and $1.9 \times 10^6 \text{ M}^{-1}$, respectively.^[17] This difference in binding affinity was also confirmed by a theoretical study. A gas-phase calculation^[18] showed that the binding energies $-E_{\text{BSSE}}$ of **1** and **2** for NH_4^+ are 62.3 and 58.2 kcal mol⁻¹, respectively. The difference in binding energy between the two receptors (4.1 kcal mol⁻¹) is in reasonable agreement with the experimental value (ca. 1.7 kcal mol⁻¹)^[19] calculated from the association constants if we consider solvation effects.^[5, 20] These experimental and theoretical studies support a prominent role of cation– π interaction in the recognition of NH_4^+ by **1**.

In conclusion, we have presented a rationally designed cage-type receptor that binds NH_4^+ by cation– π interaction and hydrogen bonding, with high sensitivity and selectivity comparable or even superior to those of nonactin over a wide range of pH. In particular, this work demonstrates that an appropriate combination of hydrogen bonding and cation– π interaction can produce a large synergic effect in recognizing NH_4^+ . The high performance, low cost, and easy synthesis may warrant practical applications of this receptor as an ammonium ion sensor.

Experimental Section

1: A solution of 1,3,5-tris(bromomethyl)-2,4,6-triethylbenzene (58 mg, 0.13 mmol) and 1,3,5-tris[(2-methoxyphenoxy)methyl]-2,4,6-triethylbenzene (69 mg, 0.13 mmol) in DMF (26 mL) was added to a suspension of K_2CO_3 (109 mg, 0.79 mmol) and [18]crown-6 (7 mg, 0.026 mmol) in DMF (6 mL) at 85 °C over 12 h. The resulting mixture was stirred for an additional 4 h and then cooled to room temperature. After addition of water and chloroform to the mixture, the aqueous layer was separated and extracted twice with chloroform. The combined organic layers were concentrated and purified by chromatography on silica to afford **1** (43 mg, 46 %). ¹H NMR (CDCl_3): δ = 6.92 (s, 2H), 4.96 (s, 2H), 2.60 (q, 2H), 1.03 (t, 3H); ¹³C NMR (CDCl_3): δ = 149.6, 145.6, 131.8, 120.2, 111.1, 63.9, 23.1, 16.6; MS (FAB): m/z : 727.3 [M^+]; satisfactory elemental analysis (C, H).

Received: January 19, 2001 [Z16463]

- [1] a) P. Bühlmann, E. Pretsch, E. Bakker, *Chem. Rev.* **1998**, 98, 1593; b) M. Dobler in *Comprehensive Supramolecular Chemistry*, Vol. 1 (Eds.: J. L. Atwood, J. E. D. Davies, D. D. MacNicol, F. Vögtle), Pergamon, Oxford, **1996**.
- [2] a) S. Ozawa, P. C. Hauser, K. Seiler, S. S. S. Tan, W. E. Morf, W. Simon, *Anal. Chem.* **1991**, 53, 640; b) S. J. West, S. Ozawa, K. Seiler, S. S. S. Tan, W. Simon, *Anal. Chem.* **1992**, 54, 633.
- [3] a) C. J. Pedersen, *J. Am. Chem. Soc.* **1967**, 89, 7017; b) J. M. Lehn, *Acc. Chem. Res.* **1978**, 11, 49; c) D. Siswanta, H. Hisamoto, H. Tohma, N. Yamamoto, K. Suzuki, *Chem. Lett.* **1994**, 945; d) T. Moriuchi-Kawakami, S. Nakazawa, M. Ota, M. Nishihira, H. Hayashi, Y. Shibutani, T. Shono, *Anal. Sci.* **1998**, 14, 1065; e) K. Suzuki, D. Siswanta, T. Otsuka, T. Amano, T. Ikeda, H. Hisamoto, R. Yoshihara, S. Ohba, *Anal. Chem.* **2000**, 72, 2200; f) H.-S. Kim, H. J. Park, H. J. Oh, Y. K. Koh, J.-H. Choi, D.-H. Lee, G. S. Cha, H. Nam, *Anal. Chem.* **2000**, 72, 4683.
- [4] a) M. Dobler, J. D. Dunitz, B. T. Kilburn, *Helv. Chim. Acta* **1969**, 52, 2573; b) K. Neupport-Laves, M. Dobler, *Helv. Chim. Acta* **1976**, 59, 614.
- [5] J. C. Ma, D. A. Dougherty, *Chem. Rev.* **1997**, 97, 1303.
- [6] a) S. L. De Wall, E. S. Meadows, L. J. Barbour, G. W. Gokel, *J. Am. Chem. Soc.* **1999**, 121, 5613; b) S. L. De Wall, L. J. Barbour, G. W. Gokel, *J. Am. Chem. Soc.* **1999**, 121, 8405; c) S. L. De Wall, E. S. Meadows, L. J. Barbour, G. W. Gokel, *Proc. Natl. Acad. Sci. USA* **2000**, 97, 6271.
- [7] J. Chin, C. Walsdorff, B. Stranix, J. Oh, H. J. Chung, S.-M. Park, K. Kim, *Angew. Chem.* **1999**, 111, 2923; *Angew. Chem. Int. Ed.* **1999**, 38, 2756.
- [8] S. H. Park, J. Chin, K. Kim, unpublished results.
- [9] K. S. Oh, C.-W. Lee, H. S. Choi, S. J. Lee, K. S. Kim, *Org. Lett.* **2000**, 2, 2679.
- [10] Molecular mechanics calculations were carried out by using Cerius2 (version 3.8, force field: universal 1.02) from Molecular Simulations Inc.
- [11] C. Walsdorff, W. Saak, S. Pohl, *J. Chem. Res. Synop.* **1996**, 282; *J. Chem. Res. Miniprint* **1996**, 1601.
- [12] Crystal data for **1**·0.5 CHCl_3 : $\text{C}_{48}\text{H}_{54}\text{O}_6(\text{CHCl}_3)_{0.5}$, M_r = 786.60, monoclinic, space group $P2_1/n$, a = 13.5223(2), b = 21.9568(4), c = 14.0568(10) Å, β = 94.261(10)°, V = 4162.0(10) Å³, Z = 4, T = 223 K, ρ_{calcd} = 1.255 g cm⁻³, Siemens SMART CCD diffractometer, $\text{Mo}_{\text{K}\alpha}$ radiation, 6485 independent reflections, R_1 = 0.0808 ($I > 2\sigma(I)$), wR_2 = 0.2037 (all data), GOF: 1.110. Crystal data for **[1-NH₄]₂PF₆·CH₃CN**: $\text{C}_{48}\text{H}_{58}\text{F}_6\text{N}_6\text{O}_6\text{P}(\text{CH}_3\text{CN})$, M_r = 930.98, trigonal, space group $P\bar{3}c1$, a = b = 16.6234(2), c = 19.6564(4) Å, V = 4704.08(12) Å³, Z = 4, T = 228 K, ρ_{calcd} = 1.315 g cm⁻³, 2520 independent reflections, R_1 = 0.0955 ($I > 2\sigma(I)$), wR_2 = 0.2576 (all data), GOF: 1.087. Hydrogen atoms of NH_4^+ were located from the difference electron density map, and their positions and thermal parameters were refined. The ethyl groups of the “bottom” benzene ring are disordered over two sites. The PF_6^- counterion is also twofold disordered. Crystallographic data (excluding structure factors) for the structures reported in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no. CCDC-156039 (**1**) and CCDC-156040 (**[1-NH₄]₂PF₆**). Copies of the data can be obtained free of charge on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: (+44) 1223-336-033; e-mail: deposit@ccdc.cam.ac.uk).
- [13] Although the distances between the ammonium nitrogen atom and receptor oxygen atoms O1 and O2 are essentially the same (3.226(5) and 3.224(5) Å, respectively), the different N–H···O angles (122(7) and 168(12)°, respectively) suggest that the ammonium ion forms a stronger hydrogen bond with O2 than with O1 in the solid state.
- [14] K. S. Kim, J. Y. Lee, S. J. Lee, T.-K. Ha, D. H. Kim, *J. Am. Chem. Soc.* **1994**, 116, 7399.
- [15] E. Bakker, P. Bühlmann, E. Pretsch, *Chem. Rev.* **1997**, 97, 3083.
- [16] S. S. Moore, T. L. Tarnowski, M. Newcomb, D. J. Cram, *J. Am. Chem. Soc.* **1977**, 99, 6398.
- [17] Furthermore, an ISE based on **1** is more selective and sensitive than that based on **2** for binding NH_4^+ over K^+ or Na^+ (see Figure 2b): detection limit $4.4 \times 10^{-6} \text{ M}$ (**2**).
- [18] Ab initio calculations were performed with Gaussian 98.^[21] The geometries were fully optimized with C_3 symmetry conditions at the B3LYP/3-21G* level. The binding energies of complexes were corrected for basis set superposition error (BSSE). The BSSE was calculated by using the counterpoise method^[22] with a modification that takes account of the energy effect due to the change of geometries upon complexation.^[23] Binding energies and BSSE were calculated at the B3LYP/6-31G** level.
- [19] A similar gain in host–guest binding energy through interactions between alkylammonium ions and aromatic rings has been observed: a) H.-J. Schneider, T. Blatter, S. Simova, I. Theis, *J. Chem. Soc. Chem. Commun.* **1989**, 580; b) H.-J. Schneider, T. Blatter, P. Zimmermann, *Angew. Chem.* **1990**, 102, 1194; *Angew. Chem. Int. Ed. Engl.* **1990**, 29, 1161.
- [20] O. M. Cabarcos, C. J. Weinheimer, J. M. Lisy, *J. Chem. Phys.* **1999**, 110, 8429.
- [21] Gaussian 98 (Revision A.7), M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Montgomery, R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo,

- S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, C. Gonzalez, M. Challacombe, P. M. W. Gill, B. G. Johnson, W. Chen, M. W. Wong, J. L. Andres, M. Head-Gordon, E. S. Replogle, J. A. Pople, Gaussian, Inc., Pittsburgh, PA, 1998.
- [22] S. F. Boys, F. Bernardi, *Mol. Phys.* **1970**, *19*, 553.
- [23] P. I. Nagy, D. A. Smith, G. Alagona, C. Ghio, *J. Phys. Chem.* **1994**, *98*, 486.

Macrocycles within Macrocycles: Cyclen, Cyclam, and Their Transition Metal Complexes Encapsulated in Cucurbit[8]uril**

Soo-Young Kim, In-Sun Jung, Eunsung Lee, Jaheon Kim, Shigeru Sakamoto, Kentaro Yamaguchi, and Kimoon Kim*

*Dedicated to Professor Eiichi Kimura
on the occasion of his retirement*

Inclusion of a macrocycle in a macrocycle, reminiscent of Russian Matryoshka dolls, is a rare phenomenon, although inclusion of crown ethers, cryptands, and their alkali metal complexes inside the cavity of γ -cyclodextrin has been known for many years.^[1] Recently, the encapsulation of cryptands, crown ethers, and their alkali metal complexes in supramolecular capsules and cages held together by hydrogen bonds and coordination bonds, respectively, has also been reported.^[2, 3] However, few examples of the inclusion of transition metal macrocyclic complexes in synthetic molecular or supramolecular host systems are available in the literature^[4] even though such systems are potentially useful as catalysts with high specificity as a result of the unique microenvironment around the transition metal ion which is reminiscent of metalloenzymes.

Cucurbituril (CB[6]), a macrocycle comprising six glycoluril units, forms stable host–guest complexes with small

molecules, such as aliphatic and aromatic amines.^[5] Our recent discovery^[6] of new cucurbituril homologues, cucurbit-[*n*]uril (CB[*n*]; *n* = 5, 7, and 8), containing five, seven, and eight glycoluril units has opened up new opportunities to expand the host–guest chemistry of cucurbituril. For example, the largest member of the cucurbituril family, CB[8], has a cavity comparable to that of γ -cyclodextrin and can accommodate two aromatic guest molecules to form 1:2 host–guest complexes,^[6] or 1:1:1 ternary complexes.^[7] The large cavity prompted us to explore the inclusion of macrocycles and their transition metal complexes in CB[8]. Of the many potential candidates for the guest macrocycles we chose tetraazamacrocycles, not only because they are effective hosts for transition metal ions^[8] but also because their transition metal complexes exhibit catalytic activities in many reactions, such as epoxidation and DNA hydrolysis.^[9] Here we present novel macrocycles within macrocycles, in which cyclen (1,4,7,10-tetraazacyclododecane), cyclam (1,4,8,11-tetraazacyclotetradecane), and their transition metal complexes are encapsulated in CB[8].^[10]

Heating an aqueous solution of CB[8] and cyclen tetrahydrochloride at 100 °C for 2 h, followed by slow cooling to room temperature produces colorless crystals of complex **1a**. The ¹H NMR spectrum of **1a** in D₂O reveals a shift of the cyclen signal to higher field relative to that of free cyclen, which is consistent with the formation of a 1:1 host–guest complex of CB[8], and cyclen.^[11] The X-ray crystal structure of **1a** (Figure 1) confirms the encapsulation of a fully protonated cyclen macrocycle in the cavity of CB[8] with four Cl[−] counterions residing outside CB[8]. The inner macrocycle is significantly tilted with respect to the outer macrocycle: the angle between the average planes of the inner




Figure 1. X-ray crystal structure of **1a**: a) top view and b) side view. Color code: oxygen: red, nitrogen: blue, carbon: gray. Hydrogen atoms, counterions (Cl[−]), and solvent (H₂O) molecules are omitted for clarity.

[*] Prof. Dr. K. Kim, S.-Y. Kim, I.-S. Jung, E. Lee, Dr. J. Kim
National Creative Research Initiative Center for
Smart Supramolecules
and Department of Chemistry
Division of Molecular and Life Sciences
Pohang University of Science and Technology
San 31 Hyojadong, Pohang 790-784 (Republic of Korea)
Fax: (+82) 54-279-8129
E-mail: kkim@postech.ac.kr

S. Sakamoto, Prof. Dr. K. Yamaguchi
Chemical Analysis Center, Chiba University
1-33 Yayoicho, Inageku, Chiba 263-8522 (Japan)

[**] We gratefully acknowledge the Korean Ministry of Science and Technology (Creative Research Initiative Program) for support of this work, and the Korean Ministry of Education (Brain Korea 21 program) for graduate studentships to S.-Y.K. and E.L. We also thank Mr. Woo Sung Jeon for his help with the electrochemistry and Professor P. K. Bharadwaj for helpful discussions.

Supporting information for this article is available on the WWW under <http://www.angewandte.com> or from the author.