


It is now clear that the chemoselectivities in the Lewis acid promoted reactions of certain aldehydes and ketones are opposite to those in the reactions in the absence of Lewis acid. By proper choice of the reaction conditions, either by carrying out the reaction in the presence or in the absence of Lewis acids, secondary and primary alcohols, homoallylic alcohols, and Diels–Alder adducts can be synthesized chemoselectively. A further extension of the above concept is being investigated in our laboratories.

Received: March 20, 2001 [Z16808]

The Fate of Bis(η^3 -allyl)palladium Complexes in the Presence of Aldehydes (or Imines) and Allylic Chlorides: Stille Coupling versus Allylation of Aldehydes (or Imines)

Hiroyuki Nakamura, Ming Bao, and Yoshinori Yamamoto*

Palladium-catalyzed Stille coupling between allylic chlorides **1** and allyltributyltin proceeds through bis(η^3 -allyl)palladium intermediates **2** to give the allylic–allylic 1,5-hexadiene coupling products **3** and **3'** in moderate to high yields (Scheme 1).^[1] Recently, we found that **2** (R = H), generated



Scheme 1. Different reaction pathways of bis(η^3 -allyl)palladium intermediates **2**.

from allyl chloride and allyltributyltin in the presence of a palladium catalyst, reacts with aldehydes **4** and imines **5** to produce the corresponding homoallyl alcohols **6** and amines **7**, respectively, in high yields.^[2, 3] Also, we found that unsymmetrical bis(η^3 -allyl)palladium complexes **2** can selectively transfer the unsubstituted allyl group to aldehydes and imines.^[4] Important questions are what factors control the reaction pathways of bis(η^3 -allyl)palladium intermediates **2**, and why does the Stille coupling of **2** cease in the presence of aldehydes **4** or imines **5**?


We now report that phosphine ligands play a key role in this process; the Stille coupling reaction takes place in the presence of triphenylphosphine,^[5] even if aldehydes **4** and imines **5** are present, whereas the allylation of aldehydes and imines occurs in the absence of the phosphine, even in the presence of allylic chlorides **1** (Scheme 2).

The palladium-catalyzed reactions of various aldehydes **4** and imines **5** with allylic chlorides were investigated in the presence or absence of PPh₃ (Table 1). In the absence of a phosphine ligand, the reaction of **1a** and allyltributyltin with benzaldehyde (**4a**) gave the homoallyl alcohol **6a** in 94% yield, and cinnamyl chloride **1a** was recovered in essentially

- [1] For reviews, see: a) *Modern Carbonyl Chemistry* (Ed.: J. Otera), Wiley-VCH, Weinheim, **2000**; b) *Lewis Acids in Organic Synthesis, Vol. 1–2* (Ed.: H. Yamamoto), Wiley-VCH, Weinheim, **2000**; c) *Lewis Acid Reagents* (Ed.: H. Yamamoto), Oxford University Press, New York, **1999**; d) R. Mahrwald, *Chem. Rev.* **1999**, *99*, 1095–1120; e) A. Mengel, O. Reiser, *Chem. Rev.* **1999**, *99*, 1191–1223; f) M. Santelli, J. M. Pons, *Lewis Acids and Selectivity in Organic Synthesis*, CRC, Boca Raton, **1996**; g) S. Shambayati, S. L. Schreiber in *Comprehensive Organic Synthesis, Vol. 1* (Eds.: B. M. Trost, I. Fleming), Pergamon, Oxford, **1991**, pp. 283–324; h) M. Yamaguchi in *Comprehensive Organic Synthesis, Vol. 1* (Eds.: B. M. Trost, I. Fleming), Pergamon, Oxford, **1991**, pp. 325–353.
- [2] For reports of chemoselective aldol reactions mediated by special Lewis acids, for example, [Eu(dppm)₃] and Bu₃SnClO₄, see: a) K. Mikami, M. Terada, T. Nakai, *J. Org. Chem.* **1991**, *56*, 5456–5459; b) J. Chen, J. Otera, *Synlett* **1997**, 29–30; c) J. Chen, J. Otera, *Tetrahedron* **1997**, *53*, 14275–14286.
- [3] The discrimination between two different carbonyl compounds on the basis of steric effects has been reported; see: S. Saito, H. Yamamoto in *Modern Carbonyl Chemistry* (Ed.: J. Otera), Wiley-VCH, Weinheim, **2000**, pp. 33–67.
- [4] α -Alkoxy aldehydes can be selectively activated by bidentate Lewis acids in the presence of other aldehydes as a result of the chelation effect; see: M. T. Reetz, A. Gansäuer, *Tetrahedron* **1993**, *49*, 6025–6030, and references therein.
- [5] Y. Yamamoto, N. Asao, *Chem. Rev.* **1993**, *93*, 2207–2293.
- [6] The rather low yield of **9** is a result of the preferential dimerization of cyclopentadiene in the presence of BF₃·OEt₂.
- [7] Gaussian 94, M. J. Frisch, G. W. Trucks, H. B. Schlegel, P. M. W. Gill, B. G. Johnson, M. A. Robb, J. R. Cheeseman, T. A. Keith, J. A. Petersson, J. A. Montgomery, K. Raghavachari, M. A. Al-Laham, V. G. Zakrzewski, J. V. Ortiz, J. B. Foresman, J. Cioslowski, B. B. Stefanov, A. Nanayakkara, M. Challacombe, C. Y. Peng, P. Y. Ayala, W. Chen, M. W. Wong, J. L. Andres, E. S. Replogle, R. Gomperts, R. L. Martin, D. J. Fox, J. S. Binkley, D. J. DeFrees, J. Baker, J. J. P. Stewart, M. Head-Gordon, C. Gonzales, J. A. Pople, Gaussian, Inc., Pittsburgh, PA, **1995**.
- [8] a) B. W. Gung, *Tetrahedron Lett.* **1991**, *32*, 2867–2870; b) B. W. Gung, M. A. Wolf, *J. Org. Chem.* **1992**, *57*, 1370–1375.
- [9] The results of theoretical calculations of the LUMO in the presence, and absence, of BF₃ and transition-state calculations will be published in a full paper.
- [10] R. J. Gillespie, J. S. Hartman, *Can. J. Chem.* **1968**, *46*, 2147–2157.
- [11] W. P. Neumann, E. Heymann, *Justus Liebigs Ann. Chem.* **1965**, 683, 11.
- [12] a) A. J. Leusink, H. A. Budding, J. W. Marsman, *J. Organomet. Chem.* **1968**, *13*, 155–162; b) A. J. Leusink, H. A. Budding, W. Drenth, *J. Organomet. Chem.* **1968**, *13*, 163–168.

[*] Prof. Dr. Y. Yamamoto, Dr. H. Nakamura, Dr. M. Bao
Department of Chemistry
Graduate School of Science
Tohoku University
Sendai 980-8578 (Japan)
Fax: (+81)22-217-6784
E-mail: yoshi@yamamoto1.chem.tohoku.ac.jp

Supporting information for this article is available on the WWW under <http://www.angewandte.com> or from the author.



Scheme 2. The chemoselectivities of bis(η^3 -allyl)palladium complexes in the presence of aldehydes (or imines) and allylic chlorides. **4a–4e** and **5a–5d** were used for the reactions shown in Table 1.

quantitative yield (entry 1), whereas in the presence of one equivalent of PPh_3 (relative to Pd), the reaction gave a 2/1 mixture of **6a** and **3a** + **3a'**. When two equivalents of PPh_3 were used, the ratio of **6a** to **3a** + **3a'** became 1/1. With 3–5 equiv of PPh_3 , only the Stille coupling reaction took place; using four equivalents of PPh_3 gave the best result: a 92/8 mixture of **3a** and **3a'** in 90% yield, and benzaldehyde was recovered quantitatively (entry 1).^[6] Similar observations were made in the reactions of **1b**, **1c**, and **1a** (entries 2–4). β -Elimination took place with the aliphatic substrate **1d** in the presence of PPh_3 to give 1,3-tetradecadiene (entry 5).^[7] A key role of PPh_3 in controlling the chemoselectivity was also observed in the reactions of the imines. In the absence of the phosphine ligand, the reaction of **1a**, allyltributyltin, and **5a** gave the allylation product **7a** in 82% yield, and **1a** was recovered, whereas in the presence of 4 equiv of PPh_3 , a 92/8 mixture of **3a** and **3a'** was obtained in 91% yield, and the imine **5a** remained unchanged (entry 6). Similar results were obtained in the other cases (entries 7–9). Thus, highly chemoselective reactions can be accomplished by means of the presence or absence of PPh_3 .


We investigated the chemoselectivity on substrates **8a** and **8b**, which contain an allyl chloride unit and an aldehyde group in the same molecule (Scheme 3). In

Table 1. Palladium-catalyzed reaction of allylic chlorides **1** and allyltributyltin with aldehydes **4** or imines **5** in the presence or absence of PPh_3 .^[a]

Entry	1 , R	4 or 5	Ligand/ time [h]	Yield [%] of 6 or 7 ^[b]	Yield [%] of 3 + 3' (3/3') ^[b,c]
1	1a , Ph	4a	none/11 PPh_3 /19	94 –	– 90 (92/8)
2	1b , <i>p</i> -MeC ₆ H ₄	4b	none/37 PPh_3 /13	83 –	– 78 (100/0)
3	1c , <i>p</i> -BrC ₆ H ₄	4c	none/22 PPh_3 /22	98 –	– 80 (83/17)
4	1a , Ph	4d	none/26 PPh_3 /22	93 –	– 88 (92/8)
5	1d , CH ₃ (CH ₂) ₁₀	4e	none/29 PPh_3 /17	71 –	– – ^[c]
6	1a , Ph	5a	none/2 PPh_3 /4	82 –	– 91 (92/8)
7	1a , Ph	5b	none/64 PPh_3 /4	78 –	– 82 (91/9)
8	1a , Ph	5c	none/132 PPh_3 /5	92 –	– 79 (91/9)
9	1a , Ph	5d	none/132 PPh_3 /21	73 –	– 80 (91/9)

[a] All reactions were carried out in the presence of 5 mol% of $[\text{Pd}_2(\text{dba})_3] \cdot \text{CHCl}_3$ in THF at room temperature (dba = dibenzylideneacetone). In the Stille coupling, 40 mol% of PPh_3 was added. The aldehydes **4** and imines **5** are shown in Scheme 2. [b] Yields of isolated products based on **1**, **4**, or **5**. [c] β -Elimination took place to give the diene product CH₃(CH₂)₉CH=CHCH=CH₂ (89%, *cis/trans* = 1/1).

the absence of the phosphine ligand, the *ortho* isomer **8a** reacted with allyltributyltin to give the allylation product **9a** in 88% yield, and the allyl chloride unit remained unchanged.




Scheme 3. Investigation of chemoselectivity on substrates **8a** and **8b**, which contain an allyl chloride unit and an aldehyde group in the same molecule.

However, in the presence of 4 equiv of PPh_3 , only the Stille coupling reaction took place to give a 96/4 mixture of **10a** and **10a'** in 73% yield, and the aldehyde group remained unchanged. Similarly, the *para* isomer **8b** gave **9b** in 83% yield, whereas in the presence of PPh_3 a 90/10 mixture of **10b** and **10b'** was produced in 74% yield.

A plausible mechanism is shown in Scheme 4. Oxidative addition of **1a** to palladium(0) gives the η^3 -allyl(chloro)palladium complex **11**, and transmetalation of **11** with allyltributyltin produces the bis(η^3 -allyl)palladium complex **12**. In the

presence of PPh_3 (1 equiv) to give **3a**, whereas only a trace of **3a** was obtained in the absence of PPh_3 . Therefore, the coordination of PPh_3 to the complex **12** in the reaction is essential for the Stille coupling process.

In conclusion, the fate of bis(η^3 -allyl)palladium complexes in the presence of aldehydes (or imines) and allylic chlorides has been clarified for the first time. The PPh_3 ligand plays a key role in controlling the chemoselectivity. Now, we can perform chemoselective allylation of allylic halides and aldehydes (imines).



Scheme 4. Proposed mechanism for the chemoselective allylation in the presence of cinnamyl chloride **1a** and benzaldehyde **4a**.

absence of PPh_3 , benzaldehyde (**4a**) coordinates to palladium(II) to produce the homoallyloxypalladium complex **13**. The transmetalation of **13** with allyltributyltin produces the corresponding homoallyloxytin compound **15** and regenerates the bis(η^3 -allyl)palladium intermediate **12**. In the allylation reaction, the unsubstituted η^3 -allyl group of **12** undergoes nucleophilic addition to benzaldehyde, while the phenyl-substituted allyl group acts as a nontransferable η^3 -allyl ligand.^[4b] Therefore, only catalytic amounts of cinnamyl chloride **1a** react initially with allyltributyltin to afford the bis(η^3 -allyl)palladium intermediate **12**, and large amounts of **1a** remain in the reaction medium. However, in the presence of PPh_3 , the phosphine ligand coordinates to **12** to give **16** and/or **17**.^[8] Reductive elimination from **16** and/or **17** gives the Stille coupling product **3a** and regenerates the palladium(0) catalyst. Indeed, the Stille coupling reaction of allyltributyltin with the dimer of complex **11**, which was synthesized according to a literature procedure,^[9] proceeded in the

Received: March 20, 2001

Revised: June 8, 2001 [Z16809]

- [1] a) J. K. Stille, *Angew. Chem.* **1986**, 98, 504–520; *Angew. Chem. Int. Ed. Engl.* **1986**, 25, 508–524; b) T. N. Mitchell, *Synthesis* **1992**, 803–815; c) V. Farina, G. P. Roth, *Adv. Met. Org. Chem.* **1996**, 5, 1–53; d) V. Farina, V. Krishnamurthy, W. J. Scott, *Org. React.* **1997**, 50, 1–652.
- [2] a) H. Nakamura, K. Aoyagi, J.-G. Shim, Y. Yamamoto, *J. Am. Chem. Soc.* **2001**, 123, 372–377; b) H. Nakamura, H. Iwama, Y. Yamamoto, *Chem. Commun.* **1996**, 1459–1460.
- [3] Recently, a theoretical study on the reactivity of bis(η^3 -allyl)palladium complexes was reported: K. J. Szabo, *Chem. Eur. J.* **2000**, 6, 4413–4421.
- [4] a) H. Nakamura, H. Iwama, Y. Yamamoto, *J. Am. Chem. Soc.* **1996**, 118, 6641–6647; b) H. Nakamura, K. Nakamura, Y. Yamamoto, *J. Am. Chem. Soc.* **1998**, 120, 4242–4243.
- [5] In general, the Stille coupling reaction of allylic halides with allyl tin compounds requires ligands such as triphenylphosphine and maleic anhydride to induce reductive elimination in the catalytic cycle: a) J. Godschalx, J. K. Stille, *Tetrahedron Lett.* **1980**, 21, 2599–2602; b) J. M. Cuerva, E. Gómez-Bengoa, M. Méndez, A. M. Echavarren, *J. Org. Chem.* **1997**, 62, 7540–7541; c) A. Goliaszewski, J. Schwartz, *Organometallics* **1985**, 4, 417–419.
- [6] Representative procedures for the palladium-catalyzed allylation with allyltributyltin in the presence of allylic chlorides and aldehydes (or imines) are given in the Supporting Information.
- [7] In the absence of allyltributyltin, under otherwise similar reaction conditions, β -hydrogen elimination did not occur, and the starting materials **1d** and **4e** were recovered. It is known that the presence of allyltributyltin facilitates β -hydrogen elimination: E. Keinan, S. Kumar, V. Dangur, J. Vaya, *J. Am. Chem. Soc.* **1994**, 116, 11151–11152.
- [8] O. Kuhn, H. Mayr, *Angew. Chem.* **1999**, 111, 356–358; *Angew. Chem. Int. Ed.* **1999**, 38, 343–346.
- [9] P. R. Auburn, P. B. Mackenzie, B. Bosnich, *J. Am. Chem. Soc.* **1985**, 107, 2033–2046.