

- [6] a) C. M. Jensen, S. Nemeh, W. C. Kaska, presented at the 1980 Biennial Inorganic Chemistry Symposium, (Guelph, Ontario, Canada), **1980**; b) S. Nemeh, C. Jensen, E. Binamira-Soriaga, W. C. Kaska, *Organometallics* **1983**, 2, 1442–1447; c) M. Gupta, C. Hagen, W. C. Kaska, R. E. Cramer, C. M. Jensen, *J. Am. Chem. Soc.* **1997**, 119, 840–841.
- [7] C. M. Jensen, *Chem. Commun.* **1999**, 2443–2449, and refs. therein.
- [8] “Anthraphos” has previously been used to refer to the chiral ligand *trans*-11,12-bis(diphenylphosphino)-9,10-dihydro-9,10-ethanoanthracene: a) T. Y. Fu, Z. Liu, J. R. Scheffer, J. Trotter, *Tetrahedron Lett.* **1994**, 35, 7593–7596; b) T. Y. Fu, Z. Liu, S. T. Rettig, J. R. Scheffer, J. Trotter, *Acta Crystallogr. Sect. C* **1997**, 53, 1577–1579. We thank Prof. J. R. Scheffer for the permission to reuse this name.
- [9] a) M. W. Haenel, D. Jakubik, C. Krüger, P. Betz, *Chem. Ber.* **1991**, 124, 333–336; b) M. W. Haenel, S. Oevers, J. Bruckmann, J. Kuhnigk, C. Krüger, *Synlett* **1998**, 301–303.
- [10] *t*Bu₂PSiMe₃^[11] with *t*BuOK in analogy to: F. Uhlig, R. Hummelt, *J. Organomet. Chem.* **1993**, 452, C9–C10.
- [11] *t*Bu₂PSiMe₃: H. Schumann, L. Rösch, *Chem. Ber.* **1974**, 107, 854–868. Alternatively it was prepared from *t*Bu₂PH in THF by consecutive addition of *n*BuLi in hexane and Me₃SiCl.
- [12] In cyclododecane (b.p. 250 °C) the ring strain of 2.4 kcal mol^{−1} is much lower than in cyclodecane (12.0 kcal mol^{−1}) or cyclooctane (10.0 kcal mol^{−1}) which frequently are used as test substrate for catalytic dehydrogenations. The enthalpy of about 33 kcal mol^{−1} necessary to dehydrogenate alkanes to alkenes and hydrogen is reduced by relief of ring strain.
- [13] a) W.-W. Xu, G. P. Rosini, M. Gupta, C. M. Jensen, W. C. Kaska, K. Krogh-Jespersen, A. S. Goldman, *Chem. Commun.* **1997**, 2273–2274; b) F. Liu, A. S. Goldman, *Chem. Commun.* **1999**, 655–656.
- [14] M. T. Reetz, G. Haderlein, K. Angermund, *J. Am. Chem. Soc.* **2000**, 122, 996–997.
- [15] Computations were carried out using a Silicon Graphics Octane workstation and the modeling software SYBYL 6.7 (Tripos Assoc., St. Louis, USA, **2001**). Important parameters for the *ams* calculation: geometries of the catalyst fragments without the phosphorus and iridium substituents are based on X-ray structures of **3**^[6c] and of the IrCl₃ complex of **9** (C. W. Lehmann, M.-A. Schneider, S. Oevers, W. C. Kaska, M. W. Haenel, unpublished results). Systematic search about the P–C_{alkyl} bonds with 5° and the C_{alkyl}–C_{methyl} bonds with 30° (**3**, **6**) or 60° (**4**, **7**) torsion angle increments. All conformations with *E* < 3 kcal mol^{−1} (force field: TAFF, no charges) higher than the energy minimum are superimposed according to Figure 2, forming a “super-molecular” fragment. The *ams* was determined using Connolly’s method^[16] analogous to the previously published procedure^[14] by “rolling” a sphere of 1.8 Å diameter over the Ir part of the molecular surface of these super-molecular fragments.
- [16] M. L. Connolly, *Science* **1983**, 221, 709–713.
- [17] Computational Details: All of the calculations were carried out by using DFT^[18] with either B3LYP or BP86^[19, 20, 21] as implemented in Gaussian 98.^[22] TS are confirmed by separated frequency calculations. The Ir basis set was a modified LANL2DZ basis.^[23] For the P atoms, the LANL2DZ basis set was augmented by a d-type polarization function.^[23b, 24] The carbon and hydrogen atoms bonded to Ir and those in ethane, ethene, and dihydrogen were described by the Dunning–Huzinaga double-zeta basis sets with polarization functions (D95*), while STO-3G was used for the uncoordinated carbon and hydrogen atoms in the ligand of **8**.
- [18] R. G. Parr, W. Yang, *Density-functional theory of atoms and molecules*, Oxford University Press, Oxford, **1989**.
- [19] a) A. D. Becke, *Phys. Rev.* **1988**, A38, 3098–3100; b) A. D. Becke, *J. Chem. Phys.* **1993**, 98, 1372–1377; c) A. D. Becke, *J. Chem. Phys.* **1993**, 98, 5648–5652.
- [20] C. T. Lee, W. T. Yang, R. G. Parr, *Phys. Rev.* **1988**, B37, 785–789.
- [21] J. P. Perdew, *Phys. Rev.* **1986**, B33, 8822–8824.
- [22] Gaussian 98 (Revision A.6), M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Montgomery, Jr., R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, C. Gonzalez, M. Challacombe, P. M. W. Gill, B. G. Johnson, W. Chen, M. W. Wong, J. L. Andres, M. Head-Gordon, E. S. Replogle, J. A. Pople, Gaussian, Inc., Pittsburgh, PA, **1998**.
- [23] a) LANL2DZ basis set is Dunning D95 basis sets on the first row, Los Alamos ECP plus double-zeta basis sets on Na–Bi; b) P. J. Hay, W. R. Wadt, *J. Chem. Phys.* **1985**, 82, 299–310; c) W. R. Wadt, P. J. Hay, *J. Chem. Phys.* **1985**, 82, 284–298; d) M. Couty, M. B. Hall, *J. Comput. Chem.* **1996**, 17, 1359–1370.
- [24] A. Hollwarth, M. Bohme, S. Dapprich, A. W. Ehlers, A. Gobbi, V. Jonas, K. F. Kohler, R. Stegmann, A. Veldkamp, G. Frenking, *Chem. Phys. Lett.* **1993**, 208, 237–240.
- [25] H.-J. Fan, E. Webster, M. B. Hall, unpublished results.
- [26] a) S. Niu, M. B. Hall, *J. Am. Chem. Soc.* **1999**, 121, 3992–3999; b) S. Li, M. B. Hall, *Organometallics* **2001**, 20, 2153–2160; c) S. Li, M. B. Hall, *Organometallics* **2001**, 20, 3210; d) M. Kanzelberger, B. Singh, M. Czerw, K. Krogh-Jespersen, A. S. Goldman, *J. Am. Chem. Soc.* **2000**, 122, 11017–11018.
- [27] K. Krogh-Jespersen, M. Czerw, M. Kanzelberger, A. S. Goldman, *J. Chem. Inf. Comput. Sci.* **2001**, 41, 56–63.

Nickel-Catalyzed Homoallylation of Aldehydes in the Presence of Water and Alcohols**

Masanari Kimura, Akihiro Ezoe, Shuji Tanaka, and Yoshinao Tamaru*


By virtue of their high polarity, a wide variety of allylic metal and metalloid (e.g., B, Si, Sn) complexes can serve as nucleophilic reagents toward carbonyl compounds;^[1] some allylations can be successfully performed even in the presence of water under Barbier conditions.^[2] On the other hand, because of their low polarity, the homoallylic metals that are capable of undergoing a nucleophilic addition reaction are likely to be limited to some electropositive main group metals (e.g., Li, Mg).^[3] Accordingly, reactions with these homoallylic metals should be performed with care under strictly water-free conditions and could be applied effectively to those compounds that do not possess an acidic hydrogen.

Here, we disclose that even in the presence of water and alcohols, nickel complexes serve as a catalyst to promote homoallylation of aldehydes with 1,3-dienes. The reaction proceeds under essentially the same conditions as those reported recently from these laboratories for a nickel-catalyzed homoallylation of carbonyl compounds with dienes in the presence of a stoichiometric amount of Et₃B.^[4–6] These two reactions, that is, in the presence and in the absence of water, provide bishomoallyl alcohols in comparable yields

[*] Prof. Dr. Y. Tamaru, Dr. M. Kimura, A. Ezoe, Dr. S. Tanaka
Department of Applied Chemistry, Faculty of Engineering
Nagasaki University
1-14 Bunkyo-machi, Nagasaki 852-8521 (Japan)
Fax: (+81)95-847-9008
E-mail: tamaru@net.nagasaki-u.ac.jp


[**] We thank Mr. Y. Ohhama, NMR Facility, for his outstanding technical assistance. Financial support from the Ministry of Education, Science, Sports, and Culture, Japanese Government, Grant-in-Aid for Scientific Research B, is acknowledged.

with the same sense of stereoselectivity and regioselectivity. For example, isoprene (4.0 mmol) reacted with glutaraldehyde (1.0 mmol) regioselectively at the C1 position and stereoselectively to provide **1a** in 57% yield as a single




isomer, which is 1,3-*anti* with respect to a tetrahydropyranyl oxygen and a methyl group, when exposed to $[\text{Ni}(\text{acac})_2]$ (0.1 mmol, acac = acetylacetonato) and Et_3B (2.4 mmol) in THF (5 mL) at room temperature for 40 h [Eq. (a)]. Glutaraldehyde is stable in water and commercially available as an aqueous solution.^[7] In the above reaction, a 50% aqueous solution was used as received; it follows that the homoallylation could be performed successfully in the presence of about six equivalents of water.

Butadiene reacted with glutaraldehyde (50% aqueous) similarly well and provided the expected homoallylation product **1b** in reasonable yield [Eq. (b)]; however, as noted



previously,^[4] butadiene was the only exception among the dienes examined so far that furnished such an allylation product as **2a** as a minor component.

Encouraged by the compatibility of the $[\text{Ni}(\text{acac})_2]/\text{Et}_3\text{B}/\text{diene}/\text{aldehyde}$ system with water, we next examined the applicability of the system to the homoallylation of cyclic hemiacetals **4** [Eq. (c)]. To achieve homoallylation of cyclic



hemiacetals with a homoallylic metal, one would anticipate that two equivalents of the reagent would be required; the first equivalent to abstract the hydrogen atom, the second to effect a nucleophilic addition to the aldehyde of ω -metalloxyaldehyde, which might be present as a component of an equilibrium mixture with 2-metalloxy-1-oxacycloalkane. To our surprise, the homoallylation of cyclic hemiacetals proceeded successfully under identical conditions to those for the homoallylation of aldehydes,^[4] that is, use of extra amounts of $[\text{Ni}(\text{acac})_2]$, Et_3B , and dienes was not necessary (Table 1). Furthermore, unsymmetrical dienes, such as isoprene (**3b**) and methyl sorbate (**3d**), reacted regioselectively at the C1 and C2 positions, respectively. The stereoselectivity was also

excellent; **3b** provided 1,3-*anti* isomers **5b** and **5e** exclusively (Table 1, runs 2 and 5) and **3d** afforded 1,2-*anti* isomers **5d** and **5g** as a single diastereomer (Table 1, runs 4 and 7), irrespective of the ring size of **4**. The reactivity decreases in the order of 2-hydroxytetrahydrofuran (**4a**), 2-hydroxytetrahydropyran (**4b**), and 2-hydroxy-5,6-benzo-1-oxacyclohexane (**4c**) (cf., Table 1, runs 3, 6, and 8). Indeed, a small amount of **4c** (13%) remained unchanged even after 78 h at room temperature (Table 1, run 8).

Reduction of **1a** with NaBH_4 afforded a diol **5e**, the IR and ^1H NMR spectra of which were identical in all respects to those of the product obtained in run 5, Table 1. The 1,3-*anti* structure of **5e** was derived from that of a lactone derivative **6a** on the basis of NOE experiments (Figure 1, See Exper-




Figure 1. Selected NOE data (% increment) of **6a** and **6b** derived from **5e** and **5g**, respectively.











imental Section). The 1,2-*anti* structure of **5g** was also established based on that of the lactone derivative **6b** (Figure 1).

In conclusion, we have demonstrated that a catalytic system of $[\text{Ni}(\text{acac})_2]/\text{diene}/\text{Et}_3\text{B}$ efficiently promotes homoallylation of a 50% aqueous solution of glutaraldehyde and cyclic hemiacetals at room temperature. Unsymmetrical dienes react regio- and stereoselectively and provide bishomoallyl alcohols with excellent 1,2- and 1,3-asymmetric induction. These reaction features may be ideal for modification of sugars and greatly contribute to carbohydrate and related chemistry. The low toxicity and low cost of the reagents and the mild reaction conditions are of particular interest from an environmental and an economic point of view.

Experimental Section

anti-5-Hydroxy-7-methyl-8-nonen-1-ol (5e) (run 5, Table 1): Into a flask containing $[\text{Ni}(\text{acac})_2]$ (25.7 mg, 0.1 mmol) purged with N_2 were successively added dry THF (5 mL), isoprene (400 μL , 4.0 mmol), 2-hydroxytetrahydrofuran (102.1 mg, 1.0 mmol),^[8] and Et_3B (2.4 mmol, 1M hexane) by syringe. The homogeneous mixture was stirred at room temperature for 62 h. After dilution with EtOAc , the mixture was washed with 2M HCl , saturated NaHCO_3 , and then with brine. The aqueous phases were combined and saturated with NaCl , and then extracted three times with EtOAc . The combined organic phase was dried (MgSO_4) and concentrated in vacuo. The residue was purified by column chromatography over silica gel (hexane/ EtOAc , 2/1, v/v) to provide **5e** (108.5 mg, 63%). IR (neat film): $\tilde{\nu} = 3332$ (s), 1639 (m), 1057 (m), 995 (m), 910 (s) cm^{-1} ; ^1H NMR (CDCl_3 , 400 MHz, TMS): $\delta = 1.02$ (d, $J = 6.6$ Hz, 3H), 1.36–1.63 (m, 10H), 2.33 (m, 1H), 3.65 (t, $J = 6.2$ Hz, 2H), 4.94 (dd, $J = 1.5, 10.3$ Hz, 1H), 5.03 (br dd, $J = 1.5, 17.2$ Hz, 1H), 5.78 (ddd, $J = 7.7, 10.3, 17.2$ Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz, TMS): $\delta = 20.3, 21.7, 32.7, 35.6, 37.3, 44.5, 62.9, 70.4, 112.9, 145.2$; HRMS calcd for $\text{C}_{10}\text{H}_{20}\text{O}_2$: 172.1463; found m/z (%): 172.1462 ($[M^+]$, 6), 157 (3), 154 (3), 129 (13), 116 (100); elemental analysis calcd (%) for $\text{C}_{10}\text{H}_{20}\text{O}_2$: C 69.72, H 11.70; found: C 69.32, H 11.52.

Table 1. Et₃B-promoted, Ni⁰-catalyzed homoallylation of cyclic hemiacetals **4a–c** with 1,3-dienes **3a–d**.^[a]

Run	Diene 3	Hemiacetal 4	<i>t</i> [h]	Product 5	Yield [%]
1	 3a	 4a	24	 5a	63 ^[b]
2	 3b	4a	22	 5b	83
3	 3c	4a	23	 5c	88
4	 3d	4a	46	 5d	64
5	3b	 4b	62	 5e	63
6	3c	4b	40	 5f	99
7	3d	4b	72	 5g	51
8	3c	 4c	78	 5h	82 ^[c]

[a] Reaction conditions: A homogeneous solution of **3** (4 mmol), **4** (1 mmol), [Ni(acac)₂] (0.1 mmol), and Et₃B (2.4 mmol in 1 M hexane) in THF (5 mL) was stirred under N₂ at room temperature for the period of time indicated. [b] In addition to **5a**, an allylation product, *trans*-6-octene-1,4-diol, was isolated in 23 % yield. [c] Yield based on 87 % conversion of **4c**.

cis- γ -(4-Acetoxybutyl)- α -methyl- γ -butyrolactone (**6a**): A solution of **5e** (86.1 mg, 0.5 mmol) in dichloromethane (5 mL) was cooled to -78°C , and ozone was bubbled through until a blue color appeared (ca. 10 min). The excess of ozone was removed by a flow of N₂. The mixture was allowed to warm to room temperature. Into this solution were added AcOH (2 mL), H₂O (0.2 mL), concentrated H₂SO₄ (0.2 mL), and 30 % H₂O₂ (1 mL). The mixture was stirred at room temperature for 1 h and then at 60°C for 5 h. The mixture was diluted with EtOAc, washed with saturated NaHCO₃ and then brine, and dried (MgSO₄) and concentrated in vacuo. The residue was purified by column chromatography over silica gel (hexane/EtOAc, 8/1, v/v) to provide **6a** (61.1 mg, 57 %). IR (neat film): $\tilde{\nu}$ = 2939 (s), 1755 (s), 1737 (s), 1244 (s), 1168 (s), 1002 (m), 927 (m) cm⁻¹; ¹H NMR (CDCl₃, 400 MHz, TMS): δ = 1.27 (d, *J* = 7.0 Hz, 3H), 1.51 (td, *J* = 12.5, 10.6 Hz, 1H), 1.42–1.80 (m, 6H), 2.05 (s, 3H), 2.49 (ddd, *J* = 5.3, 8.8, 12.5 Hz, 1H), 2.68 (ddq, *J* = 8.8, 12.5, 7.0 Hz, 1H), 4.07 (t, *J* = 6.6 Hz, 1H), 4.34 (ddt, *J* = 7.7, 10.6, 5.3 Hz, 1H); ¹³C NMR (CDCl₃, 100 MHz): δ = 15.1, 20.9, 22.0, 28.4, 35.2, 35.9, 37.4, 64.1, 78.2, 171.1, 179.3; HRMS calcd for C₁₁H₁₈O₄: 214.1205; found *m/z* (%): 214.1227 ([*M*⁺], 17), 196 (59), 125 (72), 95 (100).

Received: May 29, 2001 [Z17199]

- [1] Y. Yamamoto, A. Asao, *Chem. Rev.* **1993**, 93, 2207–2293.
 [2] Allylborane: a) G.-M. Chen, P. V. Ramachandran, H. C. Brown, *Angew. Chem.* **1999**, 111, 828–829; *Angew. Chem. Int. Ed.* **1999**, 38, 825–826; allylmagnesium: b) C.-J. Li, W.-C. Zhang, *J. Am. Chem. Soc.* **1998**, 120, 9102–9103; allylzinc: c) F. Marquez, A. Llebaria, A. Delgado, *Org. Lett.* **2000**, 2, 547–549; d) S. Hanessian, R.-Y. Yang, *Tetrahedron Lett.* **1966**, 37, 5273–5276; allyltin: e) T. P. Loh, J. Xu, *Tetrahedron Lett.* **1999**, 40, 2431–2434; allylmercury: f) T. H. Chan, Y. Yang, *Tetrahedron Lett.* **1999**, 40, 3863–3866; allylindium: g) W. Lu, J. Ma, Y. Yang, T. H. Chan, *Org. Lett.* **2000**, 2, 3469–3471, and references

therein; h) S. Araki, T. Kamei, T. Hirashita, H. Yamamura, K. Kawai, *Org. Lett.* **2000**, 2, 847–849; i) A. Melekhov, A. G. Fallis, *Tetrahedron Lett.* **1999**, 40, 7867–7870, and references therein; j) L. A. Paquette, J. Mendez-Andino, *Tetrahedron Lett.* **1999**, 40, 4301–4304, and references therein.

- [3] Nucleophilic reaction of homoallylzirconium is one exception: C. A. Bertelo, J. Schwartz, *J. Am. Chem. Soc.* **1976**, 98, 262–264.
 [4] M. Kimura, A. Ezoe, K. Shibata, Y. Tamaru, *J. Am. Chem. Soc.* **1998**, 120, 4033–4034.
 [5] Similar homoallylation of ketones by the use of Et₂Zn; M. Kimura, H. Fujimatsu, A. Ezoe, K. Shibata, M. Shimizu, S. Matsumoto, Y. Tamaru, *Angew. Chem.* **1999**, 111, 410–413; *Angew. Chem. Int. Ed.* **1999**, 38, 397–400.
 [6] Ni-catalyzed intramolecular homoallylation of ω -dienyl aldehyde: a) Y. Sato, M. Takimoto, M. Mori, *J. Am. Chem. Soc.* **2000**, 122, 1624–1634, and references therein; b) K. Shibata, M. Kimura, M. Shimizu, Y. Tamaru, *Org. Lett.* **2001**, 3, 2181–2183.
 [7] Pure glutaraldehyde may be prepared by removing water from an aqueous solution and subjected to the Grignard reaction: a) M. Rosenberger, D. Andrews, F. DiMaria, A. J. Duggan, G. Saucy, *Helv. Chim. Acta* **1972**, 55, 249–255; see, also: b) M. J. Kukla, H. J. Breslin, *J. Org. Chem.* **1987**, 52, 5046–5048.
 [8] S. Kodato, M. Nakagawa, K. Nakayama, T. Hino, *Tetrahedron*, **1989**, 45, 7247–7262.