

DOI: 10.1002/ange.200502409


# Reactions of Nitrogen Oxides with Heme Models: Spectral Characterization of an Elusive Five-Coordinate Fe<sup>III</sup>(porphyrin) Nitrito Intermediate\*\*

Tigran S. Kurtikyan\* and Peter C. Ford\*

The roles of nitrogen dioxide (NO<sub>2</sub>) and of the nitrite ion (NO<sub>2</sub><sup>−</sup>) in mammalian biology are drawing increasing attention.<sup>[1–4]</sup> NO<sub>2</sub> is a strong oxidant and nitrating species that initiates destructive pathways in living systems.<sup>[1]</sup> Several disease states have been suggested to be connected with exogenously and endogenously formed NO<sub>2</sub>.<sup>[2,3]</sup> Likely endogenous sources of NO<sub>2</sub> are the reactions of nitric oxide (NO) with oxyglobins and with the superoxide ion (O<sub>2</sub><sup>−</sup>), which leads to coordinated and free peroxynitrite (ONOO<sup>−</sup>),<sup>[5,6]</sup> subsequent decay of which affords both the nitrate ion and NO<sub>2</sub>.<sup>[7]</sup> Furthermore, there is increasing evidence that NO<sub>2</sub> may also be formed through enzymatic pathways with the participation of heme peroxidases.<sup>[1a]</sup> NO<sub>2</sub> has been shown to be a product of the reaction of NO and nitrate, as mediated by a heme center.<sup>[8]</sup> With regard to nitrite, the NO<sub>2</sub><sup>−</sup> ion is ubiquitous in mammalian tissues and fluids and is the largest reservoir of accessible NO equivalents in the cardiovascular system.<sup>[4a,b]</sup> Recent studies have suggested that NO<sub>2</sub><sup>−</sup> itself is either a vasodilator or forms the known vasodilator NO through reaction with heme centers.<sup>[4c]</sup> Clearly, there is a continuing incentive to elucidate the interactions of these ligands with heme models to understand the fundamental chemistry relevant to their biological properties.


Reported herein are studies of the reaction of NO<sub>2</sub> with ferroheme models Fe<sup>II</sup>(por) (por = *meso*-tetraphenylporphyrinato dianion (TPP) or *meso*-tetra-*p*-tolylporphyrinato dianion (TTP)) in porous, sublimed layers. In an earlier study,<sup>[9]</sup> it was shown that the reaction of Fe(tpp) leads to the formation

of the nitrate complex Fe(tpp)(η<sup>2</sup>-O<sub>2</sub>NO). Herein we demonstrate FTIR and UV/Vis spectroscopic evidence that manifest the formation of the elusive five-coordinate nitrito complex Fe(por)(η<sup>1</sup>-ONO) [Eq. (1)] as an intermediate in the



first stage of this reaction. Notably, the only well-characterized iron porphyrin complexes of NO<sub>2</sub> are those with sterically protecting porphyrins (e.g. “picket fence” porphyrin)<sup>[10]</sup> and the nitro–nitrosyl complexes Fe(por)(NO<sub>2</sub>)(NO) that are stable only under NO.<sup>[11]</sup> No five-coordinate nitrito complexes of the type Fe<sup>III</sup>(por)(NO<sub>2</sub>) (or Fe<sup>III</sup>(por)(ONO)) have been characterized previously, even with picket fence porphyrins.

The introduction of small increments of NO<sub>2</sub> gas into a cryostat, at room temperature, that contains sublimed layers of Fe(tpp) (see Experimental Section) leads to the appearance of three new IR spectroscopic bands at  $\tilde{\nu}$  = 1528, 901, and 751 cm<sup>−1</sup> that have their isotope counterparts at  $\tilde{\nu}$  = 1495, 878, and 747 cm<sup>−1</sup> when <sup>15</sup>NO<sub>2</sub> is used (Figure 1). For the



**Figure 1.** FTIR spectra of Fe(tpp) sublimed layers (solid line) after short ( $\approx 30$  s) exposure to NO<sub>2</sub> ( $5 \times 10^{-2}$  mmHg) followed by evacuation (dashed line). The same procedure with <sup>15</sup>NO<sub>2</sub> (dotted line).

analogous complex with Fe(tpp), only small deviations from these values (1–2 cm<sup>−1</sup>) are seen (see Table 1), but the last band is masked by an intense TPP band and can be registered only in the difference spectra. The intensity growth of these bands correlates directly with the addition of each portion of NO<sub>2</sub> (<sup>15</sup>NO<sub>2</sub>) (Figure 2) and demonstrates the formation of a new species that clearly contains a coordinated NO<sub>2</sub> ligand. This new complex is fairly stable in the solid state at room temperature, but decomposes under vacuum within a few days to give mostly the nitrosyl complex Fe(tpp)(NO) with a strong  $\nu(\text{NO})$  band at  $\tilde{\nu}$  = 1677 cm<sup>−1</sup>.

[\*] Prof. Dr. T. S. Kurtikyan  
Research Institute of Applied Chemistry (ARIAC)  
375005 Yerevan (Armenia)  
and  
Molecule Structure Research Centre (MSRC) NAS  
375014 Yerevan (Armenia)  
Fax: (+37) 410-28-2667  
E-mail: tkurt@msrc.am

Prof. Dr. P. C. Ford  
Department of Chemistry and Biochemistry  
University of California  
Santa Barbara, CA 93106 (USA)  
Fax: (+1) 806-893-4120  
E-mail: ford@chem.ucsb.edu

[\*\*] This work was supported by the U.S. Civilian Research and Development Foundation (CRDF Grant No. AC2-2520-TB-03) and by the US Department of Energy (DE-FG02-04ER15506). We thank M. E. Hakobyan for providing us with pure samples of iron(II) porphyrins.





six-coordinate structures are formed, but this was not the case. Furthermore, formation of  $\text{Fe}(\text{por})(\eta^1\text{-ONO})_2$  would involve oxidation either of the iron to give a  $\text{Fe}^{\text{IV}}$  species or of the porphyrin ring to give the porphyrin radical. For the latter, an intense  $\pi$  cation radical marker band in the vicinity of  $\tilde{\nu} = 1300\text{ cm}^{-1}$ <sup>[19]</sup> should be observed as was seen for the bisnitro complex  $\text{Co}^{\text{III}}(\text{tpp}^+)(\text{NO}_2)_2$ ,<sup>[20]</sup> which was obtained in sublimed layers. No such band was seen in the spectra reported herein.

The reaction of  $\text{Fe}(\text{ttp})(\text{ONO})$  with NO was investigated through preparation of a sample of the layered  $\text{Fe}(\text{ttp})(\text{ONO})$  material as described above, followed by cooling to 77 K and introducing excess NO. At 77 K there was no apparent reaction, but heating to 150 K led to the appearance of new bands at  $\tilde{\nu} = 1886\text{ cm}^{-1}$ , 1507, and  $934\text{ cm}^{-1}$ , whereas the bands at  $\tilde{\nu} = 1528$ , 901, and  $757\text{ cm}^{-1}$ , which were assigned to  $\text{Fe}(\text{ttp})(\text{ONO})$ , disappeared (Figure 3, bottom curve). The

conclude that the low-temperature reaction of NO with  $\text{Fe}(\text{ttp})(\text{ONO})$  is a simple addition to give  $\text{Fe}(\text{ttp})(\eta^1\text{-ONO})(\text{NO})$ , which supports the view that  $\text{Fe}(\text{por})(\text{ONO})$  is a five-coordinate nitrito complex.

Upon warming the initially formed nitrite–nitrosyl complex, isomerization to the more stable nitro–nitrosyl form  $(\text{Fe}(\text{por})(\text{NO}_2)(\text{NO}))^{[10,11]}$  as characterized by IR spectroscopic bands at  $\tilde{\nu} = 1864\text{ cm}^{-1}$  [ $\nu(\text{N}=\text{O})$ ],  $\approx 1455\text{ cm}^{-1}$  [ $\nu_\alpha(\text{NO}_2)$ ], and  $1295\text{ cm}^{-1}$  [ $\nu_s(\text{NO}_2)$ ] was observed [Figure 3, upper curve, Eq. (2)]. For both the nitrite–nitrosyl complex and its nitro–




**Figure 3.** FTIR spectra of sublimed layer containing  $\text{Fe}(\text{ttp})(\text{ONO})$  after introducing 2 mm NO into cryostat at 77 K, warming to 140 K and recooling (bottom curve); after additional warming to 240 K (upper curve) and recooling.

former bands have their isotope counterparts at  $\tilde{\nu} = 1851$ ,  $\approx 1472$ , and  $914\text{ cm}^{-1}$  when  $^{15}\text{NO}$  interacts with layers containing  $\text{Fe}(\text{ttp})(\text{O}^{15}\text{NO})$ . Notably, the product spectra are very similar to those reported<sup>[17]</sup> for metastable species obtained when a low temperature KBr pellet, which contained  $\text{Fe}(\text{ttp})(\text{NO})(\text{NO}_2)$ , was photolyzed. The photochemical product displayed IR spectroscopic bands at  $\tilde{\nu} = 1507$  and  $934\text{ cm}^{-1}$  that were assigned to the  $\nu(\text{N}=\text{O})$  and  $\nu(\text{N}-\text{O})$  nitrito stretching modes of the nitrito–nitrosyl complex  $\text{Fe}(\text{ttp})(\text{ONO})(\text{NO})$ . The  $\nu(\text{N}=\text{O})$  mode of this transient species was not reported owing to overlap with the nitrosyl band of the starting material, although difference spectra indicate that the photoproduct nitrosyl band shifts to higher frequencies. The coincidence of the spectral data reported herein for the product of the NO reaction with  $\text{Fe}(\text{ttp})(\text{ONO})$  and the metastable photoproduct implies the formation of an analogous species, namely the  $\eta^1\text{-ONO}$  complexes. Thus, we

nitrosyl isomer (formed upon warming), the spin-sensitive bands appear at the higher frequencies characteristic of the low-spin iron state.

Formation of the five-coordinate nitrito complex  $\text{Fe}(\text{por})(\eta^1\text{-ONO})$  upon initial reaction of  $\text{NO}_2$  with  $\text{Fe}(\text{por})$  is also supported by changes in the UV/Vis spectra (Figure 4). The



**Figure 4.** UV/Vis spectrum of  $\text{Fe}(\text{ttp})$  sublimed layer (solid line) after short time exposure to a  $\text{NO}_2$  atmosphere ( $4 \times 10^{-2}\text{ mmHg}$ , dashed line). Spectrum was recorded after FTIR spectroscopic measurement of the same sample evidencing formation of the nitrito complex  $\text{Fe}(\text{ttp})(\text{ONO})$ .

reaction was first monitored by FTIR spectroscopy (by using  $\text{CaF}_2$  plates) and then the electronic absorption spectrum of the same sample was recorded. The spectral changes observed are characteristic of conversion from the ferrous into the ferric state, that is, going from  $\text{Fe}(\text{ttp})$  to  $\text{Fe}(\text{ttp})(\text{ONO})$ . Furthermore, despite the difference in media (solid vs. solution), the electronic spectrum of the latter is quite similar to that obtained by Scheidt and co-workers<sup>[21]</sup> at the first stage of the reaction between  $[\text{Fe}(\text{ttp})(\text{OClO}_3)]$  and  $\text{NO}_2^-$  in  $\text{CH}_2\text{Cl}_2$ . Although the spectrum obtained under these conditions was reasonably assigned as likely being due to an iron(III) nitrite complex, the exact nature of this complex was

not clear owing to uncertainty regarding stoichiometry (one or two nitrites), mode of ligation (O or N bound), and whether solvent is coordinated. However, the electronic spectral similarities with the thin-film experiments strongly suggest that the unstable species seen in the earlier solution study<sup>[21]</sup> was indeed the five-coordinate nitrito complex Fe(tpp)( $\eta^1$ -ONO) characterized here.

Continued addition of NO<sub>2</sub> increments to the layered nitrite complex Fe(por)( $\eta^1$ -ONO) at room temperature led to the formation of the known nitrate complexes Fe(por)( $\eta^2$ -O<sub>2</sub>NO). This is consistent with a previous study<sup>[9]</sup> that found that interaction of NO<sub>2</sub> gas with a sublimed layer of Fe(tpp) gave Fe(tpp)( $\eta^2$ -O<sub>2</sub>NO). Notably, the high-frequency  $\nu$ (N=O) band at  $\tilde{\nu} \approx 1530 \text{ cm}^{-1}$  for this complex is disposed in the same spectral range as the  $\nu$ (N=O) of O-coordinated nitrito ligand. This coincidence is likely the reason that the intermediacy of the nitrito complexes in such reactions was not detected in the previous study.

The presence of an intermediate O-coordinated nitrito ligand has been proposed previously<sup>[8,22]</sup> as a reasonable interpretation of kinetic and spectral data for various transformations that take place in the course of the NO<sub>x</sub> interaction with heme models in solids and in solution. The demonstration herein that such five-coordinate iron-porphyrin nitrito complexes are formed from the reaction of Fe(por) and NO<sub>2</sub> at room temperature confirms these suggestions.

In summary, the data reported herein indicate that the first stage of the reaction of NO<sub>2</sub> with the ferrous porphyrins Fe(por) leads to the formation of the previously unknown five-coordinate ferric nitrito complex Fe(por)( $\eta^1$ -ONO). The mechanism of subsequent transformations upon the addition of new NO<sub>2</sub> increments, which leads to the eventual formation of nitrate complexes, is now under detailed investigation.

## Experimental Section

The ferrous porphyrinates Fe(por) are very sensitive to oxygen and are readily oxidized to Fe<sup>III</sup> derivatives. For this reason the more stable hexacoordinate complexes, Fe(por)(B)<sub>2</sub>, with nitrogen bases (por = TPP or the *meso*-tetra-*p*-tolylporphyrinato dianion; B is Py or Pip) were used as the precursors. These were synthesized by a known procedure.<sup>[23]</sup> The low-temperature sublimate was prepared<sup>[24]</sup> by placing a Fe(por)(B)<sub>2</sub> sample in a Knudsen cell and heating to about 470 K under high vacuum ( $P = 3 \times 10^{-5}$  Torr). Evacuation for 3 h resulted in the complete elimination of the coordinated axial ligands B, as monitored by measurement of the pressure at the outlet of the cryostat. Liquid nitrogen was then poured into the cryostat, and the Knudsen cell was heated to 520 K, at which temperature sublimation of Fe(por) onto the KBr or CaF<sub>2</sub> substrate occurred. In the latter case, CaF<sub>2</sub> plates were also used as the optical windows of the cryostat. The metal-aryl porphyrins layers obtained in this manner by sublimation onto a low-temperature (77 K) surface are spongelike<sup>[25a,b]</sup> and have high microporosity.<sup>[25c]</sup> Potential reactants easily diffuse across these layers, and adducts thus formed can be studied spectroscopically without solvent interference.

Sublimation was typically carried out over periods of 0.3–2.0 h to build up layers of thickness sufficient for UV/Vis and IR spectral studies. The Fe(por) sublimed layer was then heated to room temperature under dynamic vacuum, and small increments of NO<sub>2</sub> gas were introduced for a short-time period followed by a period of pumping out. The FTIR and UV/Vis spectra were measured after each NO<sub>2</sub> introduction.

NO<sub>2</sub> (<sup>15</sup>NO<sub>2</sub>), obtained by oxidation of NO (<sup>15</sup>NO) with excess pure dioxygen, was dried over P<sub>2</sub>O<sub>5</sub> and then was purified by fractional distillation until a pure white solid was obtained. <sup>15</sup>NO with 98.5% enrichment was purchased from the Institute of Isotopes, Republic of Georgia.

Received: July 11, 2005

Revised: October 7, 2005

Published online: December 2, 2005

**Keywords:** heme · IR spectroscopy · iron · nitrogen oxides · porphyrinoids

- [1] a) M. Kirsch, H.-G. Korth, R. Sustmann, H. de Groot, *Biol. Chem.* **2002**, 383, 389–399; b) S. Pfeiffer, B. Mayer, B. Hemmens, *Angew. Chem.* **1999**, 111, 1824–1844; *Angew. Chem. Int. Ed.* **1999**, 38, 1714–1731.
- [2] M. G. Espey, S. Xavier, D. D. Thomas, K. M. Miranda, D. A. Wink, *Proc. Natl. Acad. Sci. USA* **2002**, 99, 3481–3486.
- [3] S. Pfeiffer, A. Lass, K. Schmidt, B. Mayer, *FASEB J.* **2001**, 15, 2355–2364.
- [4] a) N. S. Bryan, T. Rassaf, R. E. Maloney, C. M. Rodriguez, F. Saijo, J. R. Rodriguez, M. Feelisch, *Proc. Natl. Acad. Sci. USA* **2004**, 101, 4308–4313; b) T. Rassaf, M. Feelisch, M. Kelm, *Free Radical Biol. Med.* **2004**, 36, 413–422; c) K. Cosby, K. S. Partovi, J. H. Crawford, R. P. Patel, C. D. Reiter, S. Martyr, B. K. Yang, M. A. Wacławiw, G. Zalos, X. Xu, K. T. Huang, H. Shields, D. B. Kim-Shapiro, A. N. Schechter, R. O. Cannon, M. T. Gladwin, *Nat. Med.* **2003**, 9, 1498–1505.
- [5] M. P. Dohle, J. W. Hoekstra, *J. Inorg. Biochem.* **1981**, 14, 351–358.
- [6] a) S. Herold, M. Exner, T. Nauser, *Biochemistry* **2001**, 40, 3385; b) S. Herold, *FEBS Lett.* **1999**, 443, 81.
- [7] J. L. Bourassa, E. P. Ives, A. L. Marqueling, R. Shimanovich, J. T. Groves, *J. Am. Chem. Soc.* **2001**, 123, 5142–5143.
- [8] T. S. Kurtikyan, G. M. Gulyan, G. G. Martirosyan, M. D. Lim, P. C. Ford, *J. Am. Chem. Soc.* **2005**, 127, 6216–6224.
- [9] T. S. Kurtikyan, T. G. Stepanyan, M. E. Akopyan, *Russ. J. Coord. Chem.* **1999**, 25, 721–725.
- [10] G. R. A. Wyllie, W. R. Scheidt, *Chem. Rev.* **2002**, 102, 1067–1089.
- [11] a) T. Yoshimura, *Inorg. Chim. Acta* **1984**, 83, 17–21; b) M. K. Ellison, C. E. Schulz, W. R. Scheidt, *Inorg. Chem.* **1999**, 38, 100–108; c) T. S. Kurtikyan, G. G. Martirosyan, I. M. Lorkovic', P. C. Ford, *J. Am. Chem. Soc.* **2002**, 124, 10124–10129.
- [12] K. Nakamoto, *Infrared and Raman Spectra of Inorganic and Coordination Compounds*, 3rd ed., Wiley, New York, **1978**, pp. 244–247.
- [13] T. S. Kurtikyan, T. G. Stepanyan, A. V. Gasparyan, *Russ. J. Coord. Chem.* **1997**, 23, 563–567.
- [14] T. G. Stepanyan, M. E. Akopyan, T. S. Kurtikyan, *Russ. J. Coord. Chem.* **2000**, 26, 453–457.
- [15] K. M. Miranda, X. Bu, I. M. Lorkovic, P. C. Ford, *Inorg. Chem.* **1997**, 36, 1992.
- [16] G. G. Martirosyan, A. S. Azizyan, T. S. Kurtikyan, P. C. Ford, *Chem. Commun.* **2004**, 1488–1489.
- [17] J. Lee, A. Y. Kovalevsky, I. Novozhilova, K. Bagley, P. Coppens, G. B. Richter-Addo, *J. Am. Chem. Soc.* **2004**, 126, 7180–7181.
- [18] H. Oshio, T. Ama, T. Watanabe, J. Kincaid, K. Nakamoto, *Spectrochim. Acta Part A* **1984**, 40, 863–870.
- [19] E. T. Shimomura, M. A. Phillippi, H. M. Goff, W. F. Sholz, C. A. Reed, *J. Am. Chem. Soc.* **1981**, 103, 6778–6780.
- [20] T. S. Kurtikyan, T. H. Stepanyan, *Russ. Chem. Bull.* **1998**, 47, 695–698.
- [21] M. J. Finnegan, A. G. Lappin, W. R. Scheidt, *Inorg. Chem.* **1990**, 29, 181–185.

- [22] M. D. Lim, I. M. Lorkovic, K. Wedeking, A. W. Zanella, S. M. Massick, P. C. Ford, *J. Am. Chem. Soc.* **2002**, *124*, 9737–9743.
- [23] L. M. Epstein, D. K. Straub, C. Maricondi, *Inorg. Chem.* **1967**, *6*, 1720–1722.
- [24] K. Nakamoto, T. Watanabe, T. Ama, M. W. Urban, *J. Am. Chem. Soc.* **1982**, *104*, 3744–3745.
- [25] a) M. P. Byrn, C. J. Curtis, Y. Hsiou, S. I. Khan, P. A. Sawin, S. K. Tendick, A. Terzis, C. E. Strouse, *J. Am. Chem. Soc.* **1993**, *115*, 9480–9497; b) M. P. Byrn, C. J. Curtis, S. I. Khan, P. A. Sawin, R. Tsurumi, C. E. Strouse, *J. Am. Chem. Soc.* **1990**, *112*, 1865–1874; c) T. S. Kurtikyan, A. V. Gasparyan, G. G. Martirosyan, G. H. Zhamkochyan, *J. Appl. Spectrosc.* **1995**, *62*, 62–66. (Russ.)
-