


Synthetic Methods

DOI: 10.1002/ange.200600556

Enantiomerically Pure Cyclopropenylcarbinols as a Source of Chiral Alkylidenecyclopropane Derivatives**

Samah Simaan, Ahmad Masarwa, Philippe Bertus, and Ilan Marek*

Over the past few decades, racemic cyclopropene and alkylidenecyclopropane (ACP) derivatives have been established as powerful tools in synthetic chemistry. Indeed, on one hand, cyclopropene derivatives can be readily transformed into more complex molecules by hydro- or carbometalation reactions^[1] on the strained internal double bond,^[2] whereas ACP derivatives have proven their usefulness by their unique reactivity with transition-metal catalysts.^[3] These catalyzed transformations, also based on the release of the high level of strain, can be performed either on the distal or proximal bonds of the three-membered ring and the *exo*-alkylidene moiety.^[4] Despite this mounting interest, the synthesis of enantiomerically pure cyclopropenes, particularly 1,2-disubstituted (internal) cyclopropenes and chiral ACPs are still in their infancy.^[5] In the course of our study on the functionalization of cyclopropenyl derivatives, we have recently shown that cyclopropenylcarbinols (readily prepared in a one-pot operation from 1,1,2-trihaloencyclopropane^[6]) are regio- and diastereoselectively reduced with LiAlH₄ in Et₂O into cyclopropylcarbinols as single *trans* isomers (Scheme 1,



Scheme 1. General preparation of ACPs (entries 5–12).

path A).^[1a] Herein, we document the first preparation of racemic and then enantiomerically pure ACPs **2** from cyclopropenylcarbinol derivatives **1** (Scheme 1, path B) through a copper-catalyzed addition of various Grignard reagents. In our initial investigations, the regio- and stereochemistry of the reaction was first checked on racemic cyclopropenylcarbinol derivatives **1a–h**, and the results reported in Table 1 fulfilled our expectations.

Table 1. Synthesis of ACP derivatives.

Entry	Substrate	R ¹	R ²	R ³	R ⁴	R ⁵	ACP	E/Z ^[a]	Yield [%] ^[b]
1	1a	CH ₃	H	C ₂ H ₅	C ₂ H ₅	C ₄ H ₉	2a	–	76
2	1a	CH ₃	H	C ₂ H ₅	C ₂ H ₅	C ₆ H ₅	2b	–	81
3	1b	CH ₃	H	C ₆ H ₅	C ₆ H ₅	C ₆ H ₅	2c	–	61
4	1c	C ₄ H ₉	H	C ₆ H ₅	C ₆ H ₅	C ₆ H ₅	2d	–	62
5	1d	C ₄ H ₉	H	<i>i</i> Pr	H	CH ₃	2e	100:0	82
6	1e	CH ₃	H	<i>i</i> Pr	H	C ₄ H ₉	2e	100:0	91
7	1f	CH ₃	H	C ₆ H ₅	H	C ₄ H ₉	2f	97:3	70
8	1f	CH ₃	H	C ₆ H ₅	H	C ₆ H ₅	2g	87:13	66
9	1f	CH ₃	H	C ₆ H ₅	H	CH ₃	2h	92:8	81
10	1f	CH ₃	H	C ₆ H ₅	H	C ₂ H ₅	2i	96:4	72
11	1g	CH ₃	H	Ar ^[c]	H	C ₂ H ₅	2j	95:5	88
12	1h	H	CH ₃	<i>p</i> -tol	H	CH ₃	2k	12:88	91


[a] Determined on crude NMR spectroscopic and gas-chromatographic analysis. [b] Yield of isolated product after purification by column chromatography. [c] Ar = *p*-BrC₆H₄.

In all experiments, ACPs were obtained in good-to-excellent yields which resulted from a formal copper-catalyzed S_N2' reaction of alkyl and aryl magnesium halides R⁵MgX (X = Br, I) with the free allylic alcohol **1a–h**. Tertiary and secondary alcohol derivatives (Table 1, entries 1–4 and 5–12, respectively) led similarly to ACP derivatives. Substituents on the double bond of the cyclopropenyl ring (R¹) can be either alkyl groups (entries 1–11) or a hydrogen atom (entry 12). The same is true for substituent R². Substituents R³ and R⁴ can be either alkyl, aryl, or hydrogen groups. When secondary alcohols were used (R³ = alkyl or aryl, R⁴ = H; entries 5–12), the fate of the stereochemistry of the double bond was raised. In all these experiments, we always found that the unique or major isomer was *E* configured (in entry 12, the configuration of the double bond was identical but the configuration of the product became *Z* because of the interconversion of substituents R¹ and R²).^[7] Without a copper salt, the reaction did not proceed.

With a straightforward method for the preparation of ACPs from cyclopropenylcarbinol derivatives in hand, we then turned our attention to their asymmetric synthesis. Considering that cyclopropenylcarbinols are strained allylic alcohols, we envisaged the enantiomerically enriched preparation of the ACPs to be through the Sharpless kinetic resolution (Scheme 2).^[8] When the racemic allylic alcohol **1f** (Table 2, entry 1) was subjected to the epoxidation reaction conditions using (*R,R*)-(+)-diethyl tartrate as the chiral ligand, we were pleased to observe, despite the highly reactive nature of the strained double bond, a very efficient kinetic resolution at –20 °C. Only one enantiomer is epxi-

[*] Dr. S. Simaan, A. Masarwa, Prof. I. Marek
The Mallat Family Laboratory of Organic Chemistry
Department of Chemistry
Institute of Catalysis Science and Technology
The Lise Meitner-Minerva Center for Computational Quantum Chemistry
Technion-Israel Institute of Technology, Haifa 32000 (Israel)
Fax: (+972) 4-829-3709
E-mail: chilanm@tx.technion.ac.il
Dr. P. Bertus
UMR 6519
Reactions selectives et Applications
CNRS Université de Reims Champagne-Ardenne
BP 1039, 51687 Reims Cedex 2 (France)

[**] This research was supported by the German–Israeli Project Cooperation (DIP-F.6.2), the Fund for the Promotion of Research at the Technion, and a fellowship from the Lady Davis Foundation. The authors thank Prof. Amnon Stanger for his great help in the determination of absolute configuration by computational methods. I.M. is holder of the Sir Michael and Lady Sobell Academic Chair.



Scheme 2. Kinetic resolution of cyclopropenylcarbinol derivatives. TBHP = *tert*-butyl hydroperoxide.

Table 2: Kinetic resolution of cyclopropenylcarbinol derivatives.


Entry	R ¹	R ²	R ³	Products	Yield [%] ^[b]	ee [%] ^[a]
1	CH ₃	H	C ₆ H ₅	1 f	46	99
2	CH ₃	H	<i>p</i> -BrC ₆ H ₄	1 g	44	99
3	C ₄ H ₉	H	C ₆ H ₅	1 i	44	99
4	CH ₃	H	CH ₂ CH ₂ C ₆ H ₅	1 j	42	96
5	CH ₃	CH ₃	C ₆ H ₅	1 k	47	99
6	CH ₃	CH ₃	CH ₂ CH ₂ C ₆ H ₅	1 l	40	95

[a] Yield of the isolated product after kinetic resolution and purification by column chromatography. [b] Enantiomeric excess was determined by gas-chromatographic analysis on chiral column (cyclodextrin B; see the Supporting Information).

dized to lead to the putative unstable chiral 2-oxabicyclo-[1.1.0]butane **3**.

Two enantiomerically pure α,β -unsaturated ketols **4** and **5**, formed in equal amounts (each in 22–25 % yield of isolated product) as a result of the isomerization of the oxabicyclobutane by cleavage of the two peripheral σ bonds.^[9] Oxabicyclobutanes have been postulated several times to be intermediates in various thermal^[10a] and photochemical reactions.^[10b] In none of these cases, however, were oxabicyclobutanes detected. Although we were also not able to isolate this intermediate, the Sharpless kinetic resolution implies that the corresponding oxabicyclobutane was formed as a reactive intermediate. The remaining non-oxidized products, namely, the cyclopropenylcarbinols **1** were obtained with very high enantiomeric excess and yields (95–99 % *ee* and 40–47 % yield of isolated product). The scope of the kinetic resolution is broad, as several different alkyl groups can be present either on the double bond of the cyclopropenyl unit ($R^1 = \text{CH}_3$ and C_4H_9) or on the cyclopropene ring itself ($R^2 = \text{H}$ or CH_3 ; Table 2, entries 1–4 and 5 and 6, respectively).

Aryl and alkyl groups can be interchangeably used as secondary substituents at the allylic position, although the enantiomeric excess is slightly higher when R^3 is an aryl group (99 % *ee*; entries 1–3 and 5) relative to an alkyl group (≈ 95 % *ee*; entries 4 and 6). Following the strategy we described in Scheme 1, path B for the preparation of ACPs from cyclopropenylcarbinols, alcohols **1 f, g** were treated with various Grignard reagents in the presence of 20 mol % of CuI (Scheme 3 and Table 3). The reaction proceeds with an



Scheme 3. Preparation of enantiomerically pure ACPs.

excellent transfer of chirality from chiral cyclopropenylcarbinol to ACP regardless of the nature of the alkyl magnesium halides ($R^3 = \text{aryl}$ or *alkyl*; Table 3, entries 1–4).


The absolute configuration of **2 j** was determined as *S* by using both computational methods^[11] and X-ray analysis of functionalized **2 j** (see the Supporting Information).

Table 3: Enantioselective preparation of ACPs.

Entry	Substrate	R ¹	R ²	R ³	Product	<i>E/Z</i>	Yield [%] ^[a]	ee [%] ^[b]
1	1 f	CH ₃	C ₆ H ₅	C ₆ H ₅	2 g	87:13	66	97
2	1 f	CH ₃	C ₆ H ₅	C ₄ H ₉	2 f	97:3	70	97
3	1 f	CH ₃	C ₆ H ₅	C ₂ H ₅	2 i	96:4	72	95
4	1 g	CH ₃	<i>p</i> -BrC ₆ H ₄	C ₂ H ₅	2 j	97:3	88	99

[a] Yield of isolated product after purification by column chromatography. [b] Enantiomeric excess of the *E* isomer was determined by gas-chromatographic analysis on chiral column (cyclodextrin B; see the Supporting Information).

The absolute configurations of the starting cyclopropenylcarbinol **1 g** and the final ACP **2 j** imply an overall *syn* S_N2' displacement of the alcohol moiety. Starting at -50°C , the stirred mixture was slowly warmed to room temperature and carefully followed by analysis of hydrolyzed aliquots. Under these conditions, we were able to isolate the addition products (after hydrolysis of aliquots of $[\text{Cu}(\mathbf{6})]$). These cyclopropyl-metal complexes ($[\text{Cu}(\mathbf{6})]$ or $\mathbf{6MgX}$) disappear in favor of the ACP **2**, which is obtained in good yield. As the deprotonation of the alcohol precedes the addition, the most stable conformer of the cyclopropenylcarbinolate is given in Scheme 4, with the smallest substituent (hydrogen) at the pre-existing orientation “inside” and the aryl group “outside”, away from the allylic methyl substituent (minimum 1,3-strain). Thus, this catalytic reaction proceeds through a *syn* addition/*syn* elimination mechanism (Scheme 4).



Scheme 4. Mechanistic hypothesis for the overall *syn* S_N2' process.

In conclusion, the copper-catalyzed carbomagnesiation reaction of chiral cyclopropenylcarbinol derivatives, obtained through the kinetic resolution of secondary allylic alcohols, leads to a straightforward, new preparation of enantiomerically pure alkylidenecyclopropane derivatives.

Received: February 10, 2006

Published online: April 20, 2006

Keywords: cyclopropanes · elimination · Grignard reagents · homogeneous catalysis · strained molecules

- [1] Recent hydro- and carbometalation reactions of cyclopropenes: a) E. Zohar, I. Marek, *Org. Lett.* **2004**, 6, 341; b) E. Zohar, M. Ram, I. Marek, *Synlett* **2004**, 1288; c) J. M. Fox, N. Yan, *Curr. Org. Chem.* **2005**, 9, 719; d) M. Rubin, V. Gevorgyan, *Synthesis* **2004**, 796; e) M. Rubina, M. Rubin, V. Gevorgyan, *J. Am. Chem. Soc.* **2002**, 124, 11566; f) M. Rubina, M. Rubin, V. Gevorgyan, *J. Am. Chem. Soc.* **2003**, 125, 7198; g) M. Rubina, M. Rubin, V. Gevorgyan, *J. Am. Chem. Soc.* **2004**, 126, 3688; h) L. A. Liao, J. M. Fox, *J. Am. Chem. Soc.* **2002**, 124, 14322; i) M. Nakamura, H. Isobe, E. Nakamura, *Chem. Rev.* **2003**, 103, 1295; j) I. Nakamura, G. B. Bajracharya, Y. Yamamoto, *J. Org. Chem.* **2003**, 68, 2297; k) S. Araki, T. Tanaka, T. Hirashita, J.-i. Setsune, *Tetrahedron Lett.* **2003**, 44, 8001.
- [2] R. D. Bach, O. Dmitrenko, *J. Am. Chem. Soc.* **2004**, 126, 4444.
- [3] For reviews, see: a) A. Brandi, A. Goti, *Chem. Rev.* **1998**, 98, 589; b) M. Lautens, W. Klute, W. Tam, *Chem. Rev.* **1996**, 96, 49; c) A. Brandi, S. Cicchi, F. Cordero, A. Goti, *Chem. Rev.* **2003**, 103, 1213; d) I. Nakamura, Y. Yamamoto, *Adv. Synth. Catal.* **2002**, 344, 111; e) B. M. Trost, *Angew. Chem. Int. Ed. Engl.* **1986**, 25, 1; f) *Methods of Organic Chemistry, Houben-Weyl, Vol. E17*, (Ed.: A. de Meijere), Thieme, Stuttgart, **1996**.
- [4] P. Binger, U. Schuchardt, *Chem. Ber.* **1981**, 114, 3313.
- [5] For chiral cyclopropenes, see: a) L. A. Liao, F. Zhang, N. Yan, J. A. Golen, J. M. Fox, *Tetrahedron* **2004**, 60, 1803; b) L. A. Liao, F. Zhang, O. Dmitrenko, R. D. Bach, J. M. Fox, *J. Am. Chem. Soc.* **2004**, 126, 4490; c) L. A. Liao, N. Yan, J. M. Fox, *Org. Lett.* **2004**, 6, 4937; d) M. K. Pallerla, J. M. Fox, *Org. Lett.* **2005**, 7, 3593; e) M. Isaka, E. Nakamura, *J. Am. Chem. Soc.* **1990**, 112, 7428; f) K. Kubota, M. Nakamura, M. Isaka, E. Nakamura, *J. Am. Chem. Soc.* **1993**, 115, 5867; g) Y. Lou, T. P. Remarchuk, E. J. Corey, *J. Am. Chem. Soc.* **2005**, 127, 14223; h) Y. Lou, M. Horikawa, R. A. Kloster, N. A. Hawryluk, E. J. Corey, *J. Am. Chem. Soc.* **2004**, 126, 8916; for chiral ACPs, see: i) E. Zohar, A. Stanger, I. Marek, *Synlett* **2005**, 2239; j) M. Lautens, P. H. M. Delanghe, *J. Am. Chem. Soc.* **1994**, 116, 8526; k) M. Lautens, Y. Ren, P. Delanghe, P. Chiu, S. Ma, J. Colucci, *Can. J. Chem.* **1995**, 73, 1251; l) M. Lautens, P. H. M. Delanghe, *J. Org. Chem.* **1993**, 58, 5037; m) A. de Meijere, L. Wessjohann, *Synlett* **1990**, 20; n) S. Ramaswamy, K. Prasad, O. Repic, *J. Org. Chem.* **1992**, 57, 6344; o) B. Achmatowicz, M. M. Kabat, J. Krajewski, J. Wicha, *Tetrahedron* **1992**, 48, 10201; p) J. E. Baldwin, R. M. Adlington, D. Beddington, A. T. Russell, *J. Chem. Soc. Chem. Commun.* **1992**, 1249; q) T. Nemoto, M. Ojika, Y. Sakagami, *Tetrahedron Lett.* **1997**, 38, 5667.
- [6] a) M. S. Baird, W. Nethercott, *Tetrahedron Lett.* **1983**, 24, 605; b) M. S. Baird, H. H. Hussain, *Tetrahedron* **1987**, 43, 215; c) A. R. Al-Dulayymi, M. S. Baird, *J. Chem. Soc. Perkin Trans. 1* **1994**, 1547.
- [7] The stereochemistry of **2j** was determined and assigned by analogy with the other reaction products; see the Supporting Information and also: T. Harada, H. Wada, A. Oku, *J. Org. Chem.* **1995**, 60, 5370.
- [8] Y. Gao, R. M. Hanson, J. M. Klunder, S. Y. Ko, H. Masamune, K. B. Sharpless, *J. Am. Chem. Soc.* **1987**, 109, 5765.
- [9] a) J. Ciabattoni, P. J. Kocienski, *J. Am. Chem. Soc.* **1969**, 91, 6534; b) J. Ciabattoni, P. J. Kocienski, *J. Am. Chem. Soc.* **1971**, 93, 4902; c) L. E. Friedrich, R. A. Fiato, *J. Am. Chem. Soc.* **1974**, 96, 5783; d) P. J. Kocienski, J. Ciabattoni, *J. Org. Chem.* **1974**, 39, 388; e) L. E. Friedrich, R. A. Fiato, *J. Org. Chem.* **1974**, 39, 416; f) L. E. Friedrich, N. L. de Vera, Y.-S. P. Lam, *J. Org. Chem.* **1978**, 43, 34; g) J. K. Crandall, W. W. Conover II, *J. Org. Chem.* **1978**, 43, 1323; h) B. M. Branan, X. Wang, P. Jankowski, J. Wicha, L. A. Paquette, *J. Org. Chem.* **1994**, 59, 6874.
- [10] a) S. Marmor, M. M. Thomas, *J. Org. Chem.* **1967**, 32, 252; b) E. J. Corey, J. D. Bass, R. Le Mahieu, R. B. Mitra, *J. Am. Chem. Soc.* **1964**, 86, 5570; c) H. E. Zimmerman, R. G. Lewis, J. J. McCullough, A. Padwa, S. W. Staley, M. Semmelhack, *J. Am. Chem. Soc.* **1966**, 88, 1965.
- [11] a) P. L. Polavarapu, *Chirality* **2002**, 14, 768; b) C. Forzato, G. Furlan, P. Nitti, G. Pitacco, D. Marchesan, S. Coriani, E. Valentin, *Tetrahedron: Asymmetry* **2005**, 16, 3011.