


One-Pot Arylative Epoxidation of Ketones by Employing Amphoteric Bromoperfluoroarenes**

Zhou Li and Vladimir Gevorgyan*

Bromopentafluorobenzene (PFPBr) is an important building block for the preparation of polyfluorophenyl-containing compounds;^[1] the PFP-Br bond can be cleaved under a variety of conditions (Scheme 1). Through metalation reac-




Scheme 1. Diverse reactivity of PFPBr.

tions with alkali metals, alkyl lithium reagents, or oxidative additions to Pd⁰, PFPBr serves as an equivalent of the PFP anion^[2] (mode I), which participates in reactions with diverse electrophiles or in cross-coupling reactions. On the other hand, there have been reports on Br⁺ abstraction from PFPBr by nucleophiles (mode II).^[3] Herein, we present an unprecedented mode III, in which PFPBr functions as an equivalent of both Br⁺ and PFP⁻ in the same transformation: a one-pot arylative epoxidation of ketones into pentafluorophenyl oxiranes.

Our group has recently reported that alkynyl bromides and cyanogen bromide can act as equivalents of both Br⁺ and alkynyl- or cyanide anions in a highly efficient one-pot conversion of ketones into fully substituted alkynyl or cyanoepoxides.^[4,5] As the amphoteric reactivity of PFPBr, as an equivalent of either Br⁺ or PFP⁻ has been established (Scheme 1), we hypothesized that it may also serve as a competent reagent for the one-pot cascade epoxidation of ketones (Scheme 2). It was expected that an enolate would abstract Br⁺ from PFPBr to produce α -bromoketone **I** and a PFP anion. Nucleophilic addition of the latter at the carbonyl group would generate alkoxide **II**, which, upon intramolecular S_N2 reaction, would produce oxirane **2a**.

To test the above hypothesis, the reaction of isobutyrophenone (**1a**) and PFPBr (**3a**) was examined (Table 1). Gratifyingly, it was found that employing LiHMDS in THF



Scheme 2. Proposed reaction path for the epoxidation of ketones with PFPBr. NaHMDS = sodium hexamethyldisilazide.

Table 1: Screening of reaction conditions.

Entry	Base	Solvent	Result ^[a]
1	LiHMDS	DMF	trace
2	LiHMDS	Et ₂ O	N.R.
3	LiHMDS	THF	23
4	LiHMDS	1,4-dioxane	N.R.
5	LiOiPr	THF	N.R.
6	LiOiPr	1,4-dioxane	N.R.
7	NaHMDS	THF	40 %
8	NaHMDS	Et ₂ O	75 %
9	NaHMDS	1,4-dioxane	96 %
10	KOtBu	THF	decomp
11	KOtBu	1,4-Dioxane	decomp

[a] Yields determined by GC, using pentadecane as an internal standard.

led to a 23 % yield of tetrasubstituted epoxide **2a** (entry 3). Brief optimization of the reaction conditions revealed that isobutyrophenone in the presence of NaHMDS in 1,4-dioxane was almost quantitatively converted into epoxide **2a** (entry 9). However, under these reaction conditions, 1-bromoperfluorobutane and 1-iodoperfluorobutane did not undergo this cascade transformation.^[6]



Next, the generality of the cascade transformation of various ketones with PFPBr was examined (Table 2). It was found that α,α -disubstituted methyl aryl ketones are suitable substrates for this transformation. Thus, isopropyl, cyclobutyl, cyclopentyl, and cyclohexyl phenyl ketones **1a–d** smoothly reacted with PFPBr to produce epoxides **2a–d** in good to excellent yields. Pyran-4-yl ketone **1e** was also successfully converted into the corresponding product **2e**. Diverse substituents on the phenyl ring, such as 4-methoxy **1f** and 4-cyano **1g,h**, were tolerated in this reaction (entries 6–8). Moreover, different heteroaryl ketones, including pyridin-3-yl

[*] Z. Li, Prof. V. Gevorgyan
Department of Chemistry, University of Illinois at Chicago
845 W Taylor St., Room 4500, Chicago, IL 60607 (USA)
E-mail: vlad@uic.edu
Homepage: <http://www.chem.uic.edu/vggroup>

[**] The support of the National Science Foundation (CHE-1112055) and NIH (1P50 GM-086145) is gratefully acknowledged.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201106969>.

yields (Scheme 3). Second, it was shown that polyfluorophenyl oxiranes are excellent substrates for S_NAr reactions^[9] (Scheme 4). Thus, **21** underwent an efficient substitution reaction with piperidine and sodium methylthiolate to produce the corresponding N- and S-containing products **5a** and **5b**, respectively, in excellent yields and with no oxirane ring-opening^[10] products detected.



In conclusion, we have demonstrated that amphoteric bromopolyfluoroarenes (or heteroarenes) can serve as equivalents of both Br^+ and aryl (or hetaryl) anions in the same cascade transformation. The reaction of enolizable ketones with bromopolyfluoroarenes leads to a variety of valuable tri- and tetrasubstituted epoxides^[11] in good to excellent yields and diastereoselectivities. The synthetic utility of the polyfluorophenyl-containing oxirane products was demonstrated by their further transformation, including a semi-pinacol rearrangement and S_NAr reactions.

Experimental Section

Ketone (0.5 mmol) was added to an oven-dried, conical vial equipped with a magnetic stir bar and a PTFE-topped screw cap. The vessel was then evacuated and recharged three times with argon. Anhydrous 1,4-dioxane (2.5 mL) was added dropwise and the reaction mixture was stirred until the ketone was completely dissolved. NaHMDs (0.6 mL, 1 M in THF) was subsequently added to the solution and the resulting mixture was stirred for 5 min. PFPBr, or another aryl bromide, was then added dropwise to this mixture and stirred for another 15 min, at which time precipitation of NaBr was observed. The reaction mixture was filtered through a silica or zeolite pad and washed with 50 mL of diethyl ether. The filtrate was concentrated under reduced pressure and the residue was purified by column chromatography on silica gel with a gradient from pure hexanes to 20:1 (v/v) hexanes/ethyl acetate to obtain the final product.

Received: October 2, 2011

Published online: December 15, 2011

Keywords: bromopentafluorobenzene · cascade transformations · epoxidation · perfluorinated compounds · perfluoroarylepoxydes

- [1] a) H.-J. Frohn, H. Franke, P. Fritzen, V. V. Bardin, *J. Organomet. Chem.* **2000**, 598, 127; b) P. B. Glover, A. P. Bassett, P. Nockemann, B. M. Kariuki, R. Van Deun, Z. Pikramenou, *Chem. Eur. J.* **2007**, 13, 6308; c) Y. Brinkmann, R. J. Madhusaw, R. Jazsar, G. Bernardinelli, E. P. Kündig, *Tetrahedron* **2007**, 63, 8413; d) M. Kuprat, M. Lehmann, A. Schulz, A. Villinger, *Organometallics* **2010**, 29, 1421; e) P. Romanato, S. Duttwyler, A. Linden, K. K. Baldrige, J. S. Siegel, *J. Am. Chem. Soc.* **2010**, 132, 7828; f) A. M. Trzeciak, H. Bartosz-Bechowski, Z. Ciunik, K. Niesyty, J. J. Ziolkowski, *Can. J. Chem.* **2001**, 79, 752; g) J. Y. Lee (Albemarle Corporation (Richmond, VA) USA), U. S. Patent 6248265 B1, Jan. 2, 2001.
- [2] For examples on preparation of pentafluorophenyl lithium and magnesium reagents, see: a) R. J. De Pasquale, *J. Organomet. Chem.* **1968**, 15, 233; b) M. Abarbri, F. Dehmel, P. Knochel, *Tetrahedron Lett.* **1999**, 40, 7449; c) W. Tyrre, M. S. Wickleder, *J. Organomet. Chem.* **2003**, 677, 28; d) A. Franzke, A. Pfaltz, *Chem. Eur. J.* **2011**, 17, 4131. For examples of the metalation of PFPBr by oxidative addition with transition metals, see: e) B. B. Anderson, C. L. Behrens, L. J. Radonovich, K. J. Klabunde, *J. Am. Chem. Soc.* **1976**, 98, 5390; f) A. C. Albéniz, P. Espinet, B. Martín-Ruiz, D. Milstein, *J. Am. Chem. Soc.* **2001**, 123, 11504; g) M.-Y. Jin, N. Yoshikai, *J. Org. Chem.* **2011**, 76, 1972; h) N. R. Patel, R. L. Kirchmeier, *Inorg. Chem.* **1992**, 31, 2537; i) V. A. Ivushkin, P. K. Sazonov, G. A. Artamkina, I. P. Beletskaya, *J. Organomet. Chem.* **2000**, 597, 77. For examples of the generation of PFP anion through electrophilic bromine abstraction by phosphines, see: j) V. V. Bardin, L. S. Pressman, L. N. Rogoz, G. G. Furin, *J. Fluorine Chem.* **1991**, 53, 213.
- [3] For examples of electrophilic bromine abstraction, see: a) L. R. Falvello, J. Fornies, R. Navarro, A. Rueda, E. P. Urriolabeitia, *Organometallics* **1996**, 15, 309; b) R. Bolton, J. P. B. Sandall, *J. Fluorine Chem.* **1976**, 7, 540; c) See also ref. [2i].
- [4] A. Trofimov, N. Chernyak, V. Gevorgyan, *J. Am. Chem. Soc.* **2008**, 130, 13538.
- [5] Z. Li, V. Gevorgyan, *Angew. Chem.* **2011**, 123, 2860; *Angew. Chem. Int. Ed.* **2011**, 50, 2808.
- [6] For an example of electrophilic iodine abstraction from perfluoroalkyl iodides, see: J. L. Howell, B. J. Muzzi, N. L. Rider, E. M. Aly, M. K. Abouelmagd, *J. Fluorine Chem.* **1995**, 72, 61.
- [7] a) J. Y. Kang, B. T. Connell, *J. Org. Chem.* **2011**, 76, 2379; b) S. Alesi, E. Emer, M. G. Capdevila, D. Petruzzello, A. Gualandi, P. G. Cozzi, *Molecules* **2011**, 16, 5298.
- [8] For examples of metal-catalyzed semipinacol rearrangements, see: a) S. Kulasegaram, R. J. Kulawiec, *J. Org. Chem.* **1997**, 62, 6547; b) B. C. Ranu, U. Jana, *J. Org. Chem.* **1998**, 63, 8212.
- [9] For S_NAr reactions on polyfluorophenyl rings, see: a) K. Tanaka, T. Takeyama, R. Yukawa, S. Iwata, T. Kurushima, *Supramol. Chem.* **2010**, 22, 186; b) R. Ranjbar-Karimi, M. Mashak-Shostari, A. Darehkordi, *Ultrason. Sonochem.* **2011**, 18, 258; c) R. Cano, D. J. Ramón, M. Yus, *J. Org. Chem.* **2011**, 76, 654; d) J. Zhao, T. Song, S. Zhu, L. Xu, *Tetrahedron* **2011**, 67, 910.
- [10] For oxirane ring opening reactions with N- and S-nucleophiles, see: a) S. Rodríguez-Esrich, K. S. Reddy, C. Jimeno, G. Colet, C. Rodríguez-Esrich, L. Solà, A. Vidal-Ferran, M. A. Pericàs, *J. Org. Chem.* **2008**, 73, 5340; b) D. Acetti, E. Brenna, C. Fuganti, F. G. Gatti, S. Serra, *Tetrahedron: Asymmetry* **2009**, 20, 2413.
- [11] For the synthetic use of oxiranes, see: a) D. M. Hodgson, C. D. Bray in *Aziridines and Epoxides in Organic Synthesis* (Eds.: A. K. Yudin), Wiley-VCH, Weinheim, **2006**, pp. 169–170; b) R. M. Hanson in *Organic Reactions*, Wiley, Hoboken, **2004**, pp. 116, 118; c) A. Nagaki, E. Takizawa, J. Yoshida, *Chem. Eur. J.* **2010**, 16, 14149.