


Regio- and Enantioselective Cobalt-Catalyzed Reductive [3+2] Cycloaddition Reaction of Alkynes with Cyclic Enones: A Route to Bicyclic Tertiary Alcohols**

Chu-Hung Wei, Subramaniyan Mannathan, and Chien-Hong Cheng*

Cycloaddition reactions are among the most powerful and widely applied methods for the synthesis of carbo- and heterocycles in organic synthesis.^[1] Transition-metal-catalyzed cycloaddition reactions are particularly attractive because they can provide new opportunities for the synthesis of various cyclic molecules in one pot.^[2] These reactions become possible as the complexation of substrates to metal temporarily polarizes and activates otherwise unreactive species.^[3] Some recent developments in this field are metal-catalyzed reductive [3+2] cycloaddition reactions utilizing α,β -unsaturated carbonyl compounds as the three-carbon unit. Sato et al. reported a stoichiometric titanium-promoted addition of tethered alkynes and enoates to synthesize bicyclic ketones.^[4] The work of Montgomery and co-workers on stoichiometric nickel-promoted addition of tethered alkynes and enals showed that the [3+2] cyclization can also be achieved using enolate metallacycles.^[5] Later, they disclosed a catalytic intermolecular version of the cycloaddition of alkynes and enals to synthesize cyclopentenols and cyclopentenones using triethylborane as the reducing agent.^[6a,b] In 2007, we reported a cobalt-catalyzed diastereoselective reductive [3+2] cycloaddition of allenes and enones affording 3-methylene cyclopentanol derivatives.^[7a] Subsequently, Urabe and co-workers described an iron-mediated intramolecular reductive cyclization of dienedioates to synthesize bicyclic ketoesters.^[8] Very recently, Ogoshi and Ohashi developed a unique Ni^0 -catalyzed [3+2] cycloaddition reaction of α,β -unsaturated phenyl esters with alkynes in *i*PrOH to yield cyclopentenone derivatives.^[9]

We have been interested in the reductive coupling of π components^[10] and related cycloaddition reaction.^[11] Herein, we report a novel cobalt-catalyzed reductive [3+2] cycloaddition of cyclic enones and alkynes affording bicyclic tertiary alcohols^[12] in good to high regioselectivity. Moreover, the asymmetric version^[13] of this catalytic reaction leading to two stereogenic centers can also be achieved by the proper choice of a chiral ligand (Scheme 1). This catalytic reaction provides an atom-economic method^[14] for the synthesis of



Scheme 1. Cobalt-catalyzed intermolecular reductive coupling and [3+2] cycloaddition reaction of enones with alkynes. L = ligand.

sterically crowded bicyclic tertiary alcohols, an important motif present in *Galbulimima* alkaloids,^[15] under mild conditions using relatively inexpensive cobalt metal complexes as catalysts.

Recently, we reported the enantioselective reductive coupling of alkynes with cyclic enones to synthesize β -substituted ketones in good yields and with high *ee* values.^[16] In the course of the study, we found that the reaction of diphenylacetylene (**1a**) with 2-cyclohexenone (**2a**) in the presence of CoBr_2 (5.0 mol %), 1,2-bis(diphenylphosphino)ethane (dppe; 5.0 mol %), Mn (1.10 mmol), ZnCl_2 (0.20 mmol), and CH_3CN at 40 °C for 24 h, gave the bicyclic product **3a** in 86% yield (Table 1, entry 1). Solvent played a vital role for the success of the reaction: CH_3CN and 1,4-dioxane afforded **3a** in 86% and 31% yield, respectively, whereas THF, CH_2Cl_2 , and toluene were totally inactive. Control experiments revealed that the reaction did not proceed without CoBr_2 /dppe, Mn, or ZnCl_2 . Furthermore, we found that **4a**, the reductive coupling product from **1a** and **2a**, could not be converted into product **3a** under the reaction conditions, which suggests that **4** is not an intermediate of **3a** (Scheme 1). To further understand the catalytic conditions, we examined this catalytic reaction in the presence of various cobalt complexes (see the Supporting Information) as the catalyst. Among them, CoI_2 /dppe, CoI_2 /dppp (dppp = 1,3-bis(diphenylphosphino)propane), CoI_2 / $\text{P}(\text{OMe})_3$, and CoI_2 / $\text{P}(\text{OPh})_3$ were active, but provided **3a** in low yield (27–38%). We also found that CoI_2 (5.0 mol %), *rac*-2,2'-bis-1,1'-binaphthyl (*rac*-binap; 5.0 mol %), Zn (1.10 mmol), ZnI_2 (0.20 mmol), in 1,4-dioxane was active, affording **3a** in 93% yield.

To investigate the scope of the cycloaddition reaction, various alkynes **1b–j** were reacted with **2a** using the CoBr_2 /dppe, Mn, ZnCl_2 system (Table 1, entries 2–10). Thus, di-*p*-tolylacetylene (**1b**) and di-*p*-chlorophenylacetylene (**1c**) underwent cycloaddition to give **3b** and **3c** in 96 and 85% yield, respectively (entries 2 and 3). In contrast, di-*p*-bromophenylacetylene (**1d**) afforded product **3d** in low yield (entry 4). Heteroaromatic alkyne, **1e** successfully partici-

[*] Dr. C.-H. Wei, Dr. S. Mannathan, Prof. Dr. C.-H. Cheng
Department of Chemistry, National Tsing Hua University
Hsinchu 30013 (Taiwan)
E-mail: chcheng@mx.nthu.edu.tw
Homepage: <http://mx.nthu.edu.tw/%7Echcheng>

[**] We thank the National Science Council of the Republic of China (NSC 98-2119-M-007-002-MY3) for support.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201205115>.

Table 1: Reductive [3+2] cycloaddition of alkynes with cyclic enones.^[a]

$R^1 \equiv R^2$ + Cycloenone		$\xrightarrow[\text{CH}_3\text{CN, 40}^\circ\text{C, 24 h}]{\text{CoBr}_2, \text{dppe, Mn, ZnCl}_2}$		Product 3		Yield [%] ^[b,c]
1a-k	2a-d					
2a: $n = 2, R^3, R^4 = H$	2b: $n = 1, R^3, R^4 = H$					
2c: $n = 1, R^3, R^4 = Me$	2d: $n = 2, R^3, R^4 = Me$					
Entry	1	2	Product 3			
1	1a	2a	3a: $R^1, R^2 = Ph$			86 (93)
2	1b	2a	3b: $R^1, R^2 = 4-MeC_6H_4$			96 (97)
3	1c	2a	3c: $R^1, R^2 = 4-ClC_6H_4$			85 (87)
4	1d	2a	3d: $R^1, R^2 = 4-BrC_6H_4$			21 (75)
5	1e	2a	3e: $R^1, R^2 = 2\text{-thienyl}$			74 (76)
6	1f	2a	3f: $R^1, R^2 = n\text{-propyl}$			85 (83)
7	1g	2a	3g: $R^2 = Ph$			93 (88)
8	1h	2a	3h: $R^2 = 4\text{-biphenyl}$			92 (91)
9	1i	2a	3i: $R^2 = naphthyl$			90 (93)
10	1j	2a	3j: $R^2 = 3\text{-MeOC}_6\text{H}_5$			81 (0)
11 ^[d,e]	1k	2a	3k (3k')			78 (32)
12	1a	2b	3l: $R^1, R^2 = Ph$			87 (84)
13	1b	2b	3m: $R^1, R^2 = 4-MeC_6H_4$			92 (91)
14	1a	2c	3n: $R^1 = Ph$			83 (77)
15	1g	2c	3o: $R^1 = Me$			86 (81)
16	1a	2d	3p			70 (0)

[a] Unless otherwise mentioned, all reactions were carried out using alkyne **1** (0.40 mmol) and alkene **2** (0.60 mmol) in the presence of CoBr_2 (5.0 mol %), dppe (5.0 mol %), Mn (1.10 mmol), ZnCl_2 (0.20 mmol) in CH_3CN at 40°C for 24 h. [b] Yield of isolated product. [c] Data in parentheses are yields of reactions carried out using CoI_2 (5.0 mol %), rac-binap (5.0 mol %), Zn (1.10 mmol), and ZnI_2 (0.20 mmol) in 1,4-dioxane (0.80 mL) at 40°C for 24 h. [d] The regioisomeric ratio of **3k/3k'** is 75:25. [e] Reaction was carried out at 60°C .

pated in the reaction to provide **3e** in 74 % yield (entry 5) and the aliphatic alkyne 4-octyne (**1f**) gave cycloadduct **3f** in good yield (entry 6). Unsymmetrical alkynes also underwent reductive cycloaddition, providing the corresponding bicyclic compounds with good to high regioselectivity (entries 7–11). Thus, 1-phenyl-1-propyne (**1g**) coupled with **2a** to provide **3g** in 93 % yield (entry 7). Similarly, **1h–j** gave the respective cycloadducts, **3h–j**, in good to excellent yield (entries 8–10). 1-Phenyl-1-pentyne (**1k**) also participated in the reaction, providing bicyclic product **3k/3k'** in a regioisomeric ratio of 75:25 (entry 11). Unfortunately, silyl-substituted alkynes did not furnish any [3+2] cycloadducts with **2a**. The present reductive [3+2] cycloaddition reaction is not suitable for terminal alkynes,^[10f] but instead leads to facile homocyclo-trimerization of alkynes under the optimized reaction conditions.

The present catalytic reaction was successfully extended to various cyclic enones, **2b–d** (Table 1, entries 11–15). Thus, 2-cyclopentenone (**2b**) reacted with **1a** and **1b**, providing **3l** and **3m** in 87 and 92 % yield, respectively (entries 12 and 13). Similarly, 4,4-dimethyl-substituted cyclopentenone (**2c**)

reacted with **1a** and **1g** to afford the respective products, **3n** and **3o**, in good yields (entries 14 and 15). Likewise, **2d** provided **3p** in 70 % yield (entry 16). On the other hand, 2-cycloheptenone and acyclic enones did not afford any [3+2] cycloadducts with **1** under the standard reaction conditions. The catalytic reactions can also be carried out using the $\text{CoI}_2/\text{rac-binap}$, Zn , ZnI_2 system with 1,4-dioxane as the solvent to give similar product yields (see Table 1).

Having successfully established the reductive [3+2] cycloaddition reaction, we then studied the asymmetric version of the reaction (Table 2). After systematic evaluation of various

Table 2: Enantioselective reductive [3+2] cycloaddition of alkynes with cyclic enones.^[a]

Entry	1	2	Product 3	Yield [%] ^[b]	ee [%]
1	1a	2a	3a: $R^1, R^2 = Ph$	71	90 (S,S)
2	1b	2a	3b: $R^1, R^2 = 4-MeC_6H_4$	64	98
3	1c	2a	3c: $R^1, R^2 = 4-ClC_6H_4$	54	98
4	1d	2a	3d: $R^1, R^2 = 4-BrC_6H_4$	61	> 99
5	1e	2a	3e: $R^1, R^2 = 2\text{-thienyl}$	76	77
6	1g	2a	3g: $R^2 = Ph$	63	90
7	1i	2a	3i: $R^2 = naphthyl$	50	92
8 ^[c]	1a	2b	3l	69	78
9	1a	2c	3n: $R^1 = Ph$	54	93
10	1g	2c	3o: $R^1 = Me$	60	64
11	1a	2d	3p	53	> 99

[a] Unless otherwise mentioned, all reactions were carried out using alkyne **1** (0.40 mmol) and alkene **2** (0.60 mmol) in the presence of CoI_2 (5.0 mol %), (*R,R,S,S*)-Duanphos (5.0 mol %), Zn (1.10 mmol), and ZnI_2 (0.20 mmol) in 1,4-dioxane (0.80 mL) at RT for 24 h. [b] Yield of isolated product. [c] Reactions were carried out using CoI_2 (5.0 mol %), (*R*)-binap (5.0 mol %), Zn (1.10 mmol), and ZnCl_2 (0.20 mmol) in 1,4-dioxane (0.80 mL) at RT for 24 h.

chiral ligands using **1a** and **2a** as model substrates in the presence of CoI_2 (5.0 mol %), Zn (1.10 mmol), and ZnI_2 (0.20 mmol) at room temperature with CH_3CN as the solvent, we found that (*R,R,S,S*)-2,2'-Di-*tert*-butyl-2,3,2',3'-tetrahydro-1*H*,1'*H*-(1,1')biisophosphindolyl (Duanphos) was the most effective chiral ligand, affording product **3a** in 71 % yield with 90 % *ee*. Other chiral ligands, such as (*R*)-binap and its derivatives, including (*R*)- H_8 -binap and (*S*)-Tol-binap, were also active, providing **3a** in good yields and moderate *ee*. On the other hand, at a slightly higher reaction temperature (40°C), (*R*)-1-(2-diphenylphosphino-1-naphthyl)isoquinoline (quinap) afforded **3a** with 96 % *ee* and moderate yield. The absolute configuration of **3a** was determined to be *S,S* by single crystal X-ray analysis^[17] of a tosylated derivative of **3a** (see the Supporting Information).

The $\text{CoI}_2/(\text{R,R,S,S})$ -Duanphos system was then applied to the asymmetric reductive [3+2] cycloaddition reaction of symmetrical aromatic alkynes, **1b–d**, with **2a**, which provided

3b–d in good yields and with high enantioselectivities (90–99%), (Table 2, entries 2–4). Similarly, **1e** furnished **3e** in 76% yield with a somewhat low *ee* value (77%; entry 5). Unsymmetrical alkynes **1g** and **1i** also underwent cycloaddition with **2a** to give single regioisomeric products **3g** and **3i** in good yields and with 90 and 92% *ee*, respectively (entries 6 and 7).

The scope of the enantioselective cycloaddition reaction was also examined with various cyclic enones, **2b–d** (Table 2, entries 8–11). In contrast to **2a**, 2-cyclopentenone (**2b**) afforded **3l** only in 78% *ee* (entry 8). In the present catalytic asymmetric reaction, it appears that the ring size and the substituent on the cyclic enones significantly affect the *ee* value of the coupling product.^[16] For example, 4,4-dimethyl-substituted cyclic enones, **2c** and **2d**, underwent cycloaddition with **1a** to afford products **3n** and **3p** in good yields and with excellent *ee* values (entries 9 and 11). Conversely, a moderate *ee* was observed for **3o** (entry 10). In the present reductive [3+2] cycloaddition reaction of **1** and **2**, a minor amount of corresponding reductive coupling product was obtained in both the racemic and asymmetric variants.

A reaction mechanism that accounts for the absolute configuration of product **3a** is proposed in Scheme 2. The reaction is likely initiated by the reduction of Co^{II} ^[16, 18a–c] to Co^{I} species **5** by zinc dust. Then, coordination of **1a** in the

equatorial position and **2a** with its *si* face in the axial position of the Co^{I} center to form **6** followed by oxidative cyclization gives cobaltacyclopentene^[18d,e] intermediate **7**. Selective protonation^[19] at the α -carbon to the keto group of **7** generates **8**. Next, carbonyl insertion into the cobalt-carbon bond forms cobaltalkoxide **9**. Reduction of **9** by Zn dust provides **10** and regenerates the Co^{I} species. Ultimately, the hydrolysis of **10** in air affords **3a**.

The [3+2] cycloaddition is closely related to the corresponding enyne reductive coupling reaction in the mechanistic pathway. Both reactions likely proceed with the same initial steps until the formation of intermediate **8** (Scheme 2). Protonation of **8** at the Co–C bond gives the reductive coupling product **4a**, whereas carbonyl insertion into the Co–C bond of **8** affords product **3a**. As shown in Scheme 2, the absolute configurations of products **3a** and **4a** appear to be determined by the coordination of enone **2a** to the cobalt complex (**6**, *si* face; or **6'**, *re* face). Thus, we anticipate that both reactions using the same chiral ligand should give products with the same absolute configuration and may be the same *ee* values. However, the reaction of **1a** with **2a** in the presence of 2 equivalents of water and a $\text{CoI}_2/(\text{R,R,S,S})$ -Duanphos complex as the catalyst afforded a mixture of reductive coupling product (*S,S*)-**4**^[16] and [3+2] cycloadduct (*S,S*)-**3a** in 23 and 41% yield with 54% and 90% *ee*, respectively; whereas, in the absence of extra water (*S,S*)-**4** and (*S,S*)-**3a** were obtained in 7 and 71% yield, with 52% and 90% *ee*, respectively. These results show that the cycloadduct **3a** and the reductive coupling product **4a** have the same absolute configurations, but their *ee* values are quite different. It is noteworthy that intermediate **8** also contains two diastereomers. The ratio of these two diastereomers and their relative reactivity will decide the *ee* values of the products. A possible cause for the *ee* difference is the reactivity difference of the two diastereomers of **8**.


In conclusion, we have demonstrated an atom-economic cobalt-catalyzed reductive [3+2] cycloaddition of alkynes and cyclic enones that affords bicyclic tertiary alcohols with high regioselectivity. Besides, we have also described an asymmetric variant for the synthesis of tertiary alcohols with high *ee* values (up to 99% *ee*). This catalytic reaction proceeds with high regio- and stereoselectivity. In the reaction, an air-stable, relatively inexpensive cobalt catalyst, a mild reducing agent, Zn, and one of the simplest hydrogen sources, water, were used.

Received: June 29, 2012

Revised: August 13, 2012

Published online: September 13, 2012

Keywords: alkynes · cobalt · cobaltacyclopentene · cycloaddition · tertiary alcohols



Scheme 2. Proposed mechanism for enantioselective reductive [3+2] cycloaddition of **1a** and **2a**.

[1] For reviews, see: a) M. Juhl, D. Tanner, *Chem. Soc. Rev.* **2009**, 38, 2983; b) K. C. Nicolaou, S. A. Snyder, T. Montagnon, G. Vassilikogiannakis, *Angew. Chem.* **2002**, 114, 1742; *Angew. Chem. Int. Ed.* **2002**, 41, 1668.

[2] For reviews, see: a) K. P. C. Vollhardt, *Angew. Chem.* **1984**, 96, 525; *Angew. Chem. Int. Ed. Engl.* **1984**, 23, 539; b) S. Saito, Y.

- Yamamoto, *Chem. Rev.* **2000**, *100*, 2901; c) L. Yet, *Chem. Rev.* **2000**, *100*, 2963; d) J. A. Varela, C. Saá, *Chem. Rev.* **2003**, *103*, 3787; e) M. P. Croatt, P. A. Wender, *Eur. J. Org. Chem.* **2010**, 19; f) I. Omae, *Appl. Organomet. Chem.* **2008**, *22*, 149.
- [3] For selected reviews of transition-metal-catalyzed cycloaddition reactions, see: a) M. Lautens, W. Klute, W. Tam, *Chem. Rev.* **1996**, *96*, 49; b) H.-U. Reissig, R. Zimmer, *Chem. Rev.* **2003**, *103*, 1151; c) M. Rubin, M. Rubina, V. Gevorgyan, *Chem. Rev.* **2007**, *107*, 3117; d) Z.-X. Yu, Y. Wang, Y. Wang, *Chem. Asian J.* **2010**, *5*, 1072.
- [4] a) K. Suzuki, H. Urabe, F. Sato, *J. Am. Chem. Soc.* **1996**, *118*, 8729; b) H. Urabe, K. Suzuki, F. Sato, *J. Am. Chem. Soc.* **1997**, *119*, 10014.
- [5] a) S. K. Chowdhury, K. K. D. Amarasinghe, M. J. Heeg, J. Montgomery, *J. Am. Chem. Soc.* **2000**, *122*, 6775; b) G. M. Mahandru, A. R. L. Skauge, S. K. Chowdhury, K. K. D. Amarasinghe, M. J. Heeg, J. Montgomery, *J. Am. Chem. Soc.* **2003**, *125*, 13481.
- [6] a) A. Herath, J. Montgomery, *J. Am. Chem. Soc.* **2006**, *128*, 14030; b) A. D. Jenkins, A. Hearth, M. Song, J. Montgomery, *J. Am. Chem. Soc.* **2011**, *133*, 14460; for related nickel-catalyzed reductive coupling of alkynes and enones, see: c) A. Herath, B. B. Thompson, J. Montgomery, *J. Am. Chem. Soc.* **2007**, *129*, 8712; d) W. Li, A. Herath, J. Montgomery, *J. Am. Chem. Soc.* **2009**, *131*, 17024; e) J. Montgomery, K. K. D. Amarasinghe, S. K. Chowdhury, E. Oblinger, J. Seo, A. V. Savchenko, *Pure Appl. Chem.* **2002**, *74*, 129; f) A. Herath, J. Montgomery, *J. Am. Chem. Soc.* **2008**, *130*, 8132; g) A. Herath, W. Li, J. Montgomery, *J. Am. Chem. Soc.* **2008**, *130*, 469.
- [7] a) H.-T. Chang, T. T. Jayanth, C.-H. Cheng, *J. Am. Chem. Soc.* **2007**, *129*, 4166; b) K. C. Chao, D. K. Rayabarapu, C.-C. Wang, C.-H. Cheng, *J. Org. Chem.* **2001**, *66*, 8804.
- [8] T. Hata, N. Hirone, S. Sujaku, K. Nakano, H. Urabe, *Org. Lett.* **2008**, *10*, 5031.
- [9] M. Ohoshi, T. Taniguchi, S. Ogoshi, *J. Am. Chem. Soc.* **2011**, *133*, 14900.
- [10] a) C.-C. Wang, P.-S. Lin, C.-H. Cheng, *J. Am. Chem. Soc.* **2002**, *124*, 9696; b) H.-T. Chang, T. T. Jayanth, C.-C. Wang, C.-H. Cheng, *J. Am. Chem. Soc.* **2007**, *129*, 12032; c) D. K. Rayabarapu, C.-H. Cheng, *Chem. Eur. J.* **2003**, *9*, 3164; d) C.-H. Yeh, R. P. Kori, C.-H. Cheng, *Angew. Chem.* **2008**, *120*, 4970; *Angew. Chem. Int. Ed.* **2008**, *47*, 4892; e) Y.-C. Wong, K. Parthasarathy, C.-H. Cheng, *J. Am. Chem. Soc.* **2009**, *131*, 18252; f) S. Mannathan, C.-H. Cheng, *Chem. Eur. J.* **2012**, *18*, 11771.
- [11] a) T. Sambaiah, L.-P. Li, D.-J. Huang, C.-H. Lin, D. K. Rayabarapu, C.-H. Cheng, *J. Org. Chem.* **1999**, *64*, 3663; b) M. Shanmugasundaram, M.-S. Wu, C.-H. Cheng, *Org. Lett.* **2001**, *3*, 4233.
- [12] M. Movassaghi, B. Chen, *Angew. Chem.* **2007**, *119*, 571; *Angew. Chem. Int. Ed.* **2007**, *46*, 565.
- [13] J. Karthikeyan, M. Jeganmohan, C.-H. Cheng, *Chem. Eur. J.* **2010**, *16*, 8989.
- [14] a) B. M. Trost, *Science* **1991**, *254*, 1471; b) B. M. Trost, *Angew. Chem.* **1995**, *107*, 285; *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 259; c) B. M. Trost, *Acc. Chem. Res.* **2002**, *35*, 695.
- [15] a) W. Zi, S. Yu, D. Ma, *Angew. Chem.* **2010**, *122*, 6023; *Angew. Chem. Int. Ed.* **2010**, *49*, 5887; b) S. Xing, Y. Li, Z. Li, C. Liu, J. Ren, Z. Wang, *Angew. Chem.* **2011**, *123*, 12813; *Angew. Chem. Int. Ed.* **2011**, *50*, 12605; c) U. Rinner, C. Lentsch, C. Aichinger, *Synthesis* **2010**, 3763.
- [16] C.-H. Wei, S. Mannathan, C.-H. Cheng, *J. Am. Chem. Soc.* **2011**, *133*, 6942.
- [17] CCDC 885584 (an (*S,S*)-tosylated derivative of **3a**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [18] a) G. Hilt, J. Treutwein, *Angew. Chem.* **2007**, *119*, 8653; *Angew. Chem. Int. Ed.* **2007**, *46*, 8500; b) J. Treutwein, G. Hilt, *Angew. Chem.* **2008**, *120*, 6916; *Angew. Chem. Int. Ed.* **2008**, *47*, 6811; c) S. Mannathan, C.-H. Cheng, *Chem. Commun.* **2010**, *46*, 1923; d) B. M. Trost, A. B. Pinkerton, D. J. Kremzow, *J. Am. Chem. Soc.* **2000**, *122*, 12007; e) B. M. Trost, A. McClory, *Org. Lett.* **2006**, *8*, 3627.
- [19] We believe that a trace amount of water present in the solvent may lead to the protonation of intermediate **7**.