


Tandem Brønsted Acid Promoted and Nazarov Carbocyclizations of Enyne Acetals to Hydroazulenones**

Luz Escalante, Carlos González-Rodríguez, Jesús A. Varela, and Carlos Saá*

The presence of hydroazulen(on)e skeletons in many bioactive natural products, such as guanacastepene^[1] or pleocarpenone,^[2] has resulted in sustained interest in their synthesis. A wide variety of approaches have been described.^[3] The most common approaches, other than the use of rearrangement reactions,^[3,4] have started from the five-membered ring, on which the seven-membered ring has been assembled by ring enlargement, reductive cyclization, metathesis, or aldol condensation.^[3,5] Alternative methods have involved the construction of the five-membered ring on the seven-membered ring and have employed metathesis, aldol condensation, ring expansion, or Nazarov cyclization.^[3,6]

Highly efficient approaches in which both rings are created in one pot have also been developed, most of them involve metal-catalyzed cycloaddition reactions (Scheme 1).^[7–14] Thus Wender et al. have used both intramolecular Rh-catalyzed [5+2] cycloaddition (Scheme 1, path a)^[7a] and intermolecular tandem Rh-catalyzed [5+2]/Nazarov cyclization;^[7b] Trost et al. have explored [5+2] cyclization protocols using Ru catalysts;^[8] Mascareñas and co-workers have employed Pd- and Pt-catalyzed [4+3] cycloaddition^[9] (Scheme 1, path b);^[10] and Ahmar et al.,^[11a] Brummond et al.^[11b] and Mukai et al.^[11c] have reported various approaches employing allenic reactions of the Pauson–Khand type (Scheme 1, path c).^[12] Strategies based on the cyclization of acyclic dienynes, mediated^[13a] or catalyzed^[13b] by metal carbenes, have also been reported (Scheme 1, paths d and e).^[14]

As a contribution to the development of metal-free, environmentally less hazardous synthetic methods,^[15] we recently described an efficient intramolecular cyclization of alkynals promoted by Brønsted acids.^[16,17] Herein we report its use in tandem with a Nazarov cyclization^[7b,18] to construct hydroazulenone skeletons **2** from enyne acetals **1**^[19] (Scheme 1).^[20]



Scheme 1. Metal-catalyzed and tandem enyne acetal-Nazarov Brønsted acid promoted carbocyclizations to hydroazulen(on)es.

Initially, we subjected enyne acetal **1a** (Table 1), to the optimized reaction conditions identified in our previous work (heating in DCE in the presence of excess trifluoroacetic acid).^[16] Cyclization proceeded smoothly to give an almost equimolar mixture of enones **2a** and **3a** in excellent combined yield (Table 1, entry 1). When the reaction was carried out at room temperature a similar yield was obtained but there was a higher proportion of **2a** (Table 1, entry 2). The major regioisomer was the single-bond-fused bicycle **2a**, which is the thermodynamically less-stable isomer.^[21] This bicycle can potentially be further functionalized in the seven-membered ring for future applications;^[6f,22] therefore we proceeded to seek reaction conditions to optimize this regioselectivity.


The use of HBF₄ instead of TFA as acid resulted in an increase in the **2a/3a** ratio to 3:1 (Table 1, entries 3 and 4). When the reaction temperature was lowered to –15°C the regioselectivity increased to 4.5:1, but a lower yield was obtained (Table 1, entry 5). No reaction occurred if the amount of HBF₄ was reduced from 3 to 1 equivalents (Table 1, entry 6). With BF₃·OEt₂ or H₂SO₄ as acid, yields were low to moderate and **3a** was slightly favored over **2a** (Table 1, entries 7 and 8). Formation of enone **3a** was also favored when triflimide was employed, and it was the exclusive product when triflic acid was used,^[20c] although in both cases yields were low (Table 1, entries 9 and 10).^[23] When the aldehyde corresponding to acetal **1a** was used as the starting compound and HBF₄ as the acid, both the yield and the **2a/3a** ratio decreased (compare Table 1, entries 4 and 11), undoubtedly because the reaction intermediate was less

[*] L. Escalante, Dr. C. González-Rodríguez, Dr. J. A. Varela, Prof. C. Saá Departamento de Química Orgánica y Centro Singular de Investigación en Química Biológica y Materiales Moleculares (CIQUS) Universidad de Santiago de Compostela 15782 Santiago de Compostela (Spain) E-mail: carlos.saa@usc.es Homepage: <http://www.usc.es/gi1603/saa>

[**] We thank the MICINN [Projects CTQ2011-28258, Consolider Ingenio 2010 (CSD2007-00006)] and Xunta de Galicia (2007/XA084 and CN2011/054) for financial support. L.E. thanks Fundayacucho (Venezuela) and Xunta de Galicia for predoctoral grants and C. G.-R. thanks the MICINN for a Juan de la Cierva Contract (JCI-2011-09946).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/ange.201205823>.

Table 1: Brønsted acid promoted carbocyclization of enyne acetal **1a** to hydroazulenones **2a** and **3a**.^[a]



Entry	Brønsted acid	T [°C]	t [h]	2a/3a ^[b]	Yield [%] ^[c]
1	TFA (20 equiv)	90 ^[d]	0.5	1.2:1	93
2	TFA (20 equiv)	20 ^[d]	3	1.8:1	96
3	HB F_4 (3 equiv)	20	0.5	2.5:1	84
4	HB F_4 (3 equiv)	0	1	3:1	89
5	HB F_4 (3 equiv)	−15	1.3	4.5:1	63
6	HB F_4 (1 equiv)	20	72	—	—
7	BF $_3$ ·OEt $_2$ (3 equiv)	20	1.5	1:1.6	39
8	H $_2$ SO $_4$ (3 equiv)	20	0.7	1:1.6	60
9	TfOH (3 equiv)	20	0.4	0:1	39
10	Tf $_2$ NH (3 equiv)	0	0.4	1:2.6	47
11 ^[e]	HB F_4 (3 equiv)	0	0.6	2.3:1	52

[a] Reaction conditions: **1a** (0.3 mmol), CH $_2$ Cl $_2$ (3 mL). [b] Regioisomers **2a** and **3a** are easily separated by chromatography. [c] Combined yields of the isolated products. [d] DCE as solvent. [e] With the aldehyde corresponding to acetal **1a** as starting compound (0.3 mmol). DCE = 1,2-dichloroethane, Tf = trifluoromethanesulfonyl, TFA = trifluoroacetic acid.


electrophilic.^[24] In view of these results, in most subsequent experiments we used the reaction conditions of entry 4 (referred to below as optimized conditions A) or entry 2 (optimized conditions B).

The relative configuration of **2a** corresponds to the conrotatory ring closure of an *E* olefin, thus indicating that, under the reaction conditions, the divinyl ketone intermediate might undergo a *Z*-to-*E* isomerization prior to the Nazarov cyclization (Scheme 1).^[25]

Once the viability of the tandem diastereoselective process had been established we explored its scope, initially by applying it to the β -substituted enyne acetals **1b–e** (see Table 2), in which, as in **1a**, the tether between the acetal and the triple bond includes a quaternary carbon with two COOMe substituents. Surprisingly, cyclization of **1b**, the *trans* isomer of **1a**, under optimized conditions A gave a lower yield and **2a/3a** ratio than that of **1a** (1.5:1; Table 2, entry 2); this result indicates that a more complex equilibrium of intermediates could affect the regioselectivity of the reaction.^[26] Not unexpectedly, the styrene-containing enyne acetal **1c** was one of the least reactive substrates (66%; Table 2, entry 3), whereas the reaction of the β,β -disubstituted **1e** was one of the more higher yielding reactions (84%, Table 2, entry 5). The results of the reaction of **1d** were better than we expected; the presence of the bulky *tert*-butyl β substituent resulted in an increased **2/3** ratio of 1:0 (Table 2, entry 4).

We next investigated the effect of modifying the tether. Satisfyingly, moderate to good yields and complete regioselectivity were achieved with the appropriate substituents; the reaction of the substrate containing geminal methyl groups adjacent to the alkyne fragment gave only **3f** (Table 2,

Table 2: Hydroazulenones and azahydroazulenones prepared by tandem carbocyclizations of β -substituted enyne acetals **1a–j**.^[a,b]



Entry	Enyne acetal	Hydroazulenone ^[c]	2/3	Yield ^[d] [%]
1			3:1	89
2			1.5:1 (1:1.5)	45 (62) ^[e]
3			2:1	66
4			1:0	62
5			2:1	84
6			0:1	77
7			1:0	63
8			1 ^[f] :1	73
9			1:3	76
10			2:1	80

Table 2: (Continued)

Entry	Enyne acetal	Hydroazulenone ^[c]	2/3	Yield ^[d] [%]
11			4.5:1	72


[a] Conditions A: **1** (0.3 mmol), HBF₄ (0.9 mmol), CH₂Cl₂ (3 mL), 0 °C, 20–60 min. [b] X = C(CO₂Me)₂. [c] Only the major regioisomer is shown. [d] Combined yields of the isolated products. [e] Conditions B: **4** (0.3 mmol), TFA (6.0 mmol), DCE (3 mL), RT, 30 min. [f] Compound **2h** was obtained as a 2:1 mixture of *trans* and *cis* diastereomers as a result of the partial isomerization of the divinylketone intermediate (only the major diastereomer is shown).^[25] Ts = *p*-toluenesulfonyl.

entry 6) and that of methyl ketal **1g** gave only **2g** (Table 2, entry 7). Interestingly, when the substrate not containing the CO₂Me groups was used, thus eliminating the Thorpe–Ingold effect,^[27] the cyclization still occurred, but with slightly reduced yield and no regioselectivity (Table 2, entry 8).^[28] The opposite regioselectivity was obtained (that is **3** was favored) when the substrate with the C(CO₂Me)₂ unit placed closer to the acetal fragment (Table 2 entry 9); this result probably due to the change of the original conformation of the seven-membered ring. Gratifyingly, this tandem reaction could be used to prepare azabicycles, as confirmed by the successful reactions of the tertiary amines (*Z*)-**1j** and (*E*)-**1j**, which afforded cyclopenta[*c*]azepinones^[29] with good yields and a 2/3 ratio of up to 4.5:1 in the case of (*E*)-**1j** (Table 2, entries 10 and 11).

The tandem reaction was equally successful when applied to α -substituted substrates **4** (Table 3), although to our initial surprise the major product afforded by enyne acetal **4a** was hydroazulenone **6a**, a bicyclic enone with both bridgeheads saturated and *cis* stereochemistry (Table 3, entry 1). When the α substituent was *tert*-butyl the reaction was totally regioselective (Table 3, entry 2). In contrast, when the α substituent was TMS, which was lost during the reaction, the other regioisomer was formed with high selectivity (Table 3, entry 3). Cyclization of the α,β -unsubstituted enyne acetal **4d** gave a similar result but required a longer reaction time (14 h vs. 5 h, Table 3, entry 4), thus suggesting that desilylation most likely occurred after the Nazarov cyclization^[30,31] Finally, the toluenesulfonamide **4e** afforded cyclopenta[*c*]azepinones **5e** and **6e** with a moderate combined yield (Table 3, entry 5).


We believe the mechanism of the tandem carbocyclizations of enyne acetals **1** and **4** most likely starts with the Brønsted acid induced formation of the electrophilic oxonium species **A** (Scheme 2).^[32] This cationic species would undergo standard heteroalkyne metathesis to the oxete intermediate **B** (through [2+2] cycloaddition),^[17a] which upon ring opening would give the divinyl ketone **C**. Subsequently, the Brønsted acid would induce a stereospecific Nazarov reaction (4π -electrocyclization)^[18] of this unpolarized^[25d] dienone **C** to give one of two possible oxyallyl cations, **D** and **E**, depending whether the starting compound had been β or α substituted. Proton elimination and enol-to-ketone tautomerization

Table 3: Hydroazulenones prepared by tandem carbocyclization of α -substituted enyne acetals **4a–e**.^[a,b]



Entry	Enyne acetal	Hydroazulenone ^[c]	5/6	Yield ^[d] [%]
1 ^[e]			1:2	72
2			0:1	58
3 ^[f]			1:0	85 ^[g] (5 h)
4 ^[f]			1:0	79 ^[g] (14 h)
5			2:1	52

[a] Conditions A: **4** (0.3 mmol), HBF₄ (0.9 mmol), CH₂Cl₂ (3 mL), 0 °C, 20–60 min. [b] X = C(CO₂Me)₂. [c] Only the major regioisomer is shown. [d] Combined yields of the isolated products. [e] Conditions B: **4** (0.3 mmol), TFA (6.0 mmol), DCE (3 mL), RT, 30 min. [f] Conditions B but at 90 °C. [g] A low yield of the cyclic enone **2k** (15 %, 24 %) was also obtained in the tandem carbocyclization of **4c** and **4d**, respectively. TMS = trimethylsilyl.




would finally afford the observed bicyclo[5,3,0]decenones **2**, **3**, **5**, and **6**.


As expected, the regioselectivity of the reaction seems to depend to a large extent on the position of the alkene substituent, α or β , and on the nature of the tether, through its influence on the conformation of the oxyallyl cation intermediate.^[6f,18b–e,33] In most cases, it is the single-bond-fused product that is favored.

Finally, we investigated the effect of linked α and β substituents by replacing the terminal alkene group with thiophene systems (Scheme 3).^[34] To our delight, the attractive tricyclic systems **8** and **10** were obtained in good yields from the 2- and 3-thiophenyl species **7** and **9**, respectively.

In summary, we have developed a new, simple, and versatile method for the synthesis of hydroazulenones from



Scheme 2. Mechanistic rationale for tandem Brønsted acid promoted carbocyclizations from enyne acetals to hydroazulenones.



Scheme 3. Preparation of hydroazulenothiophenones **8** and **10** by tandem carbocyclization of enyne acetals **7** and **9**, respectively.

linear enyne acetals. This metal-free procedure may be described as taking place through tandem Brønsted acid promoted carbocyclizations, a regioselective *exo*-carbocyclization previously applied to alkynals,^[16] and a stereospecific Nazarov cyclization. The new approach complements previously described methods based on metal-catalyzed cyclizations. Its full scope is currently being further investigated by application to a number of challenging cyclizations.

Received: July 23, 2012

Published online: November 4, 2012

Keywords: Brønsted acids · carbocyclizations · enynes · hydroazulenone · Nazarov cyclizations

- [1] a) S. F. Brady, M. P. Singh, J. E. Janso, J. Clardy, *J. Am. Chem. Soc.* **2000**, *122*, 2116–2117; b) S. F. Brady, S. M. Bondi, J. Clardy, *J. Am. Chem. Soc.* **2001**, *123*, 9900–9901; c) M. P. Singh, J. E. Janso, S. W. Luckman, S. F. Brady, J. Clardy, M. Greenstein, W. M. Maiese, *J. Antibiot.* **2000**, *53*, 256–261.
- [2] M. J. Williams, H. L. Deak, M. L. Snapper, *J. Am. Chem. Soc.* **2007**, *129*, 486–487.
- [3] D. A. Foley, A. R. Maguire, *Tetrahedron* **2010**, *66*, 1131–1175, and references therein.

- [4] a) P. M. Wovkulich in *Comprehensive Organic Synthesis*, Vol. 1 (Eds.: B. M. Trost, I. Fleming), Pergamon, New York, **1991**, chap. 3.3, pp. 733–775.
- [5] For ring enlargement, see: a) B. B. Snider, N. A. Hawryluk, *Org. Lett.* **2001**, *3*, 569–572; b) W. D. Shipe, E. J. Sorensen, *Org. Lett.* **2002**, *4*, 2063–2066; c) G. L. Mislin, M. Miesch, *J. Org. Chem.* **2003**, *68*, 433–441; for reductive cyclization, see: d) G. B. Dudley, S. J. Danishefsky, *Org. Lett.* **2001**, *3*, 2399–2402; for metathesis, see: e) M. F. Schneider, H. Junga, S. Blechert, *Tetrahedron* **1995**, *51*, 13003–13014; f) L. A. Paquette, F. Gallou, Z. Zhao, D. G. Young, J. Liu, J. Yang, D. Friedrich, *J. Am. Chem. Soc.* **2000**, *122*, 9610–9620; g) G. Mehta, J. D. Umarye, *Org. Lett.* **2002**, *4*, 1063–1066; h) A. Nakazaki, U. Sharma, M. A. Tius, *Org. Lett.* **2002**, *4*, 3363–3366; for aldol condensation, see: i) W. Tochtermann, S. Bruhn, M. Meints, C. Wolff, *Tetrahedron* **1994**, *50*, 9657–9670; for other synthetic approaches to hydroazulenones starting from the five-membered ring, see: j) J. Lee, H. Kim, J. K. Cha, *J. Am. Chem. Soc.* **1995**, *117*, 9919–9920; k) D. Schinzer, K. Ringe, *Tetrahedron* **1996**, *52*, 7475–7485; l) C. C. Hughes, J. J. Kennedy-Smith, D. Trauner, *Org. Lett.* **2003**, *5*, 4113–4115.

- [6] For metathesis, see: a) T. J. Brocksom, U. Brocksom, D. Frederico, *Tetrahedron Lett.* **2004**, *45*, 9289–9291; for aldol condensation, see: b) B. Föhlisch, R. Flogaus, G. H. Henle, S. Sendelbach, S. Henkel, *Eur. J. Org. Chem.* **2006**, 2160–2173; for ring expansion, see: c) S. Carret, J.-P. Depres, *Angew. Chem.* **2007**, *119*, 6994–6997; *Angew. Chem. Int. Ed.* **2007**, *46*, 6870–6873; for Nazarov cyclizations, see: d) E. A. Braude, W. F. Forbes, *Nature* **1951**, *168*, 874–875; e) S. E. Denmark, T. K. Jones, *J. Am. Chem. Soc.* **1982**, *104*, 2642–2645; f) P. Chiu, S. Li, *Org. Lett.* **2004**, *6*, 613–616; for other synthetic approaches to hydroazulenones starting from the seven-membered ring, see: g) A. M. Montana, K. M. Nicholas, *J. Org. Chem.* **1990**, *55*, 1569–1578.
- [7] a) P. A. Wender, H. Takahashi, B. Witulski, *J. Am. Chem. Soc.* **1995**, *117*, 4720–4721; b) P. A. Wender, R. T. Stemmler, L. E. Sirois, *J. Am. Chem. Soc.* **2010**, *132*, 2532–2533.
- [8] a) B. M. Trost, F. D. Toste, H. Shen, *J. Am. Chem. Soc.* **2000**, *122*, 2379–2380; b) B. M. Trost, J. Waser, A. Meyer, *J. Am. Chem. Soc.* **2007**, *129*, 14556–14557.
- [9] For an early report of an intramolecular allyl cationic [4+3] cycloaddition in the presence of TiF_2O , see: R. J. Giguere, S. M. Duncan, J. M. Bean, L. Purvis, *Tetrahedron Lett.* **1988**, *29*, 6071–6074.
- [10] a) M. Gulías, J. Duran, F. López, L. Castedo, J. L. Mascareñas, *J. Am. Chem. Soc.* **2007**, *129*, 11026–11027; b) B. Trillo, F. López, M. Gulías, L. Castedo, J. L. Mascareñas, *Angew. Chem.* **2008**, *120*, 965–968; *Angew. Chem. Int. Ed.* **2008**, *47*, 951–954.
- [11] a) M. Ahmar, C. Locatelli, D. Colombier, B. Cazes, *Tetrahedron Lett.* **1997**, *38*, 5281–5284; b) K. M. Brummond, H. Chen, K. D. Fisher, A. D. Kerekes, B. Rickards, P. C. Sill, S. J. Geib, *Org. Lett.* **2002**, *4*, 1931–1934; c) C. Mukai, I. Nomura, K. Yamanishi, M. Hanaoka, *Org. Lett.* **2002**, *4*, 1755–1758.
- [12] For other examples of cycloaddition procedures, see: a) B. M. Trost, R. I. Higuchi, *J. Am. Chem. Soc.* **1996**, *118*, 10094–10105; b) Y. Ni, J. Montgomery, *J. Am. Chem. Soc.* **2004**, *126*, 11162–11163.
- [13] For an Mo/carbene-mediated cyclization, see: a) D. F. Harvey, K. P. Lund, *J. Am. Chem. Soc.* **1991**, *113*, 5066–5068; for examples of domino metathesis reactions, see: b) S.-H. Kim, N. Bowden, R. H. Grubbs, *J. Am. Chem. Soc.* **1994**, *116*, 10801–10802.

- [14] For a synthesis of a bicyclo[5.3.0]decane core by a one-pot reaction involving aldol condensations, see: M. Mischne, *Tetrahedron Lett.* **2003**, *44*, 5823–5826.
- [15] For recent reviews on Brønsted acid catalysis, see: a) T. Akiyama, *Chem. Rev.* **2007**, *107*, 5744–5758; b) D. Kampen, C. M. Reisinger, B. List, *Top. Curr. Chem.* **2010**, *291*, 395–456; c) C. H. Cheon, H. Yamamoto, *Chem. Commun.* **2011**, *47*, 3043–3056.
- [16] C. González-Rodríguez, L. Escalante, J. A. Varela, L. Castedo, C. Saá, *Org. Lett.* **2009**, *11*, 1531–1533.
- [17] For Brønsted acid and Lewis acid catalyzed carbocyclizations of acetylenic aldehydes or ketones, see: a) C. E. Harding, S. L. King, *J. Org. Chem.* **1992**, *57*, 883–886; b) A. Balog, S. J. Geib, D. P. Curran, *J. Org. Chem.* **1995**, *60*, 345–352; c) M. F. Wempe, J. R. Grunwell, *J. Org. Chem.* **1995**, *60*, 2714–2720; d) J. U. Rhee, M. J. Krische, *Org. Lett.* **2005**, *7*, 2493–2495; e) T. Jin, Y. Yamamoto, *Org. Lett.* **2007**, *9*, 5259–5262; f) L.-P. Liu, D. Malhotra, Z. Jin, R. S. Paton, K. N. Houk, G. B. Hammond, *Chem. Eur. J.* **2011**, *17*, 10690–10699; g) K. Bera, S. Sarkar, S. Biswas, S. Maiti, U. Jana, *J. Org. Chem.* **2011**, *76*, 3539–3544; for a review on alkyne activation with Brønsted acids, see: h) Y. Yamamoto, I. D. Gridnev, N. T. Patil, T. Jin, *Chem. Commun.* **2009**, 5075–5087.
- [18] a) I. N. Nazarov, I. I. Zaretskaya, *Izv. Akad. Nauk SSSR Ser. Khim.* **1941**, 211–224; for recent reviews, see: b) M. A. Tius, *Eur. J. Org. Chem.* **2005**, 2193–2206; c) H. Pellissier, *Tetrahedron* **2005**, *61*, 6479–6517; d) A. J. Frontier, C. Collison, *Tetrahedron* **2005**, *61*, 7577–7606; e) N. Shimada, C. Stewart, M. A. Tius, *Tetrahedron* **2011**, *67*, 5851–5870; for recent examples of Nazarov cyclization in tandem processes, see: f) L. Zhang, S. Wang, *J. Am. Chem. Soc.* **2006**, *128*, 1442–1443; g) M. Janka, W. He, I. E. Haedicke, F. R. Fronczek, A. J. Frontier, R. Eisenberg, *J. Am. Chem. Soc.* **2006**, *128*, 5312–5313; h) W. Yin, Y. Ma, J. Xu, Y. Zhao, *J. Org. Chem.* **2006**, *71*, 4312–4315; i) H.-F. Cui, K.-Y. Dong, G.-W. Zhang, L. Wang, J.-A. Ma, *Chem. Commun.* **2007**, 2284–2286; j) G. Lemièrre, V. Gandon, K. Cariou, T. Fukuyama, A.-L. Dhimané, L. Fensterbank, M. Malacria, *Org. Lett.* **2007**, *9*, 2207–2209; k) P. Cao, X.-L. Sun, B.-H. Zhu, Q. Shen, Z. Xie, Y. Tang, *Org. Lett.* **2009**, *11*, 3048–3051; l) L. Liu, L. Wei, Y. Lu, J. Zhang, *Chem. Eur. J.* **2010**, *16*, 11813–11817; m) M. E. Krafft, D. V. Vidhani, J. W. Cran, M. Manoharan, *Chem. Commun.* **2011**, *47*, 6707–6709.
- [19] Starting acetals are more readily available than the corresponding aldehydes and give cleaner reactions. For Lewis acid catalyzed carbocyclizations of acetylenic acetals, see: a) T. Xu, Z. Yu, L. Wang, *Org. Lett.* **2009**, *11*, 2113–2116; b) T. Xu, Q. Yang, D. Li, J. Dong, Z. Yu, Y. Li, *Chem. Eur. J.* **2010**, *16*, 9264–9272.
- [20] For tandem combinations of heteroenyne metathesis and Nazarov cyclization, see: a) A. Saito, M. Umakoshi, N. Yagyu, Y. Hanzawa, *Org. Lett.* **2008**, *10*, 1783–1785; b) T. Jin, Y. Yamamoto, *Org. Lett.* **2008**, *10*, 3137–3139; c) T. Jin, F. Yang, C. Liu, Y. Yamamoto, *Chem. Commun.* **2009**, 3533–3535.
- [21] After treatment of enone **2a** with HBF₄ (3 equiv) at 0 °C or 20 °C for 24 h the starting material was intact, with total recovery. A similar result was found starting with enone **3a**.
- [22] a) T. Tsunoda, M. Amaike, U. S. Tambunan, Y. Fujise, S. Ito, *Tetrahedron Lett.* **1987**, *28*, 2537–2540; b) I. G. Collado, F. A. Macias, G. M. Massanet, F. R. Luis, *J. Chem. Soc. Perkin Trans. I* **1987**, 1641–1644; c) W. Kübler, O. Petrov, E. Winterfeldt, *Tetrahedron* **1988**, *44*, 4271–4388.
- [23] The regioselectivity of the reaction seems to be affected by the basicity of the Brønsted acid counterion, with a more basic counterion the kinetic product **2a** is favored. For counterion effects in Nazarov cyclization, see: J. Huang, D. Leboeuf, A. J. Frontier, *J. Am. Chem. Soc.* **2011**, *133*, 6307–6317.
- [24] For a related effect, see: K. M. McQuaid, D. Sames, *J. Am. Chem. Soc.* **2009**, *131*, 402–403.
- [25] For related cases, see: a) S. Giese, F. G. West, *Tetrahedron* **2000**, *56*, 10221–10228; b) Ref [18f]; c) O. Nieto Faza, C. S. Lopez, R. Alvarez, A. R. de Lera, *Chem. Eur. J.* **2004**, *10*, 4324–4333; d) W. He, I. R. Herrick, T. A. Atesin, P. A. Caruana, C. A. Kellenberger, A. J. Frontier, *J. Am. Chem. Soc.* **2008**, *130*, 1003–1011.
- [26] a) In fact, when the reaction was performed under optimized conditions B (TFA, 20 equiv; 20 °C), the reverse regiosomeric ratio was found (**2a/3a**, 1:1.5, 62 % combined yield). b) Careful monitoring by ¹H NMR spectroscopy (controlled amounts of HBF₄ added in CD₂Cl₂) showed that a faster but more-complex reaction occurred for the *trans* isomer (**1b**) than for the *cis* isomer (**1a**) thus most probably indicating, a less controlled reaction.
- [27] a) R. M. Beesley, C. K. Ingold, J. F. Thorpe, *J. Chem. Soc.* **1915**, *107*, 1080–1106; b) G. Hammond in *Steric Effects in Organic Chemistry* (Ed.: M. S. Newman), Wiley, New York, **1956**, pp. 462–470.
- [28] However, when (*E*)-**1h** was employed (not shown), a 3:1 ratio was obtained, with moderately good yield. See the Supporting Information for details.
- [29] For the thermal rearrangement of annelated azepines, see: a) L. A. Paquette, D. E. Kuhla, J. H. Barrett, *J. Org. Chem.* **1969**, *34*, 2879–2884; for a radical approach to azepines, see: b) A. J. Blake, G. H. Hollingworth, G. Pattenden, *Synlett* **1996**, 643–644; for an Rh-catalyzed [3+2] cycloaddition to annelated azepines, see: c) Q. Li, G.-J. Jiang, L. Jiao, Z.-X. Yu, *Org. Lett.* **2010**, *12*, 1332–1335.
- [30] a) M. E. Krafft, L. V. R. Boñaga, A. S. Felts, C. Hirose, S. Kerrigan, *J. Org. Chem.* **2003**, *68*, 6039–6042; b) A. de Meijere, H. Becker, A. Stolle, S. I. Kozhushkov, M. T. Bes, J. Salaün, M. Notemeyer, *Chem. Eur. J.* **2005**, *11*, 2471–2482.
- [31] When there was a TMS group in the β position, this substrate afforded a complex mixture instead of undergoing the regio- and stereocontrolled silyl-directed Nazarov cyclization described in Ref. [6e]. See the Supporting Information for details.
- [32] For the alternative route from **1** to divinyl ketone **C** that is shown in Scheme 2 (direct addition of the activated alkyne to the cationic carbon of the carbonyl group), see Refs [16] and [20c].
- [33] a) J. Ichikawa, S. Miyazaki, M. Fujiwara, T. Minami, *J. Org. Chem.* **1995**, *60*, 2320–2321; b) M. Harmata, *Chemtracts: Org. Chem.* **2004**, *17*, 416–435.
- [34] For an example of a Nazarov cyclization employing heteroaromatic compounds, see: J. A. Malona, J. M. Colbourne, A. J. Frontier, *Org. Lett.* **2006**, *8*, 5661–5664.