


Preferential Formation of Cyclic Trimers by Palladium-Catalyzed Oxidative Coupling Reactions of 2,18-Diethynylporphyrins**

Sumito Tokuji, Hideki Yorimitsu,* and Atsuhiko Osuka*

Inspired by the gigantic cyclic arrangements of bacteriochlorophylls and chlorophylls in photosynthetic light-harvesting antenna, considerable efforts have been devoted to the synthesis of cyclic porphyrin oligomers (CPOs).^[1] Among these, directly 1,3-butadiyne-bridged conjugated CPOs have a special place owing to their intriguing structural, electronic, and optical properties. Usually, effective macrocyclizations require synthetic tricks. Sugiura et al. used L-shaped 5,10-diaryl-15,20-diethynylporphyrin as a key corner part for the synthesis of square *meso*-to-*meso* butadiyne-bridged planar CPOs.^[2] Anderson and co-workers developed an elegant template-directed synthesis of *meso*-to-*meso* butadiyne-bridged circular CPOs using linear 5,15-diaryl-10,20-diethynylporphyrin precursors,^[3] in which the structure of the template–substrate complex dictates the major CPO product. This approach has evolved into a conceptually new Vernier-template synthesis.^[3d,f] We have also reported Cu-mediated oxidative dimerization reaction of 5,10,15-triaryl-2,18-diethynylporphyrins **1M** that provided β -to- β 1,3-butadiyne-bridged cyclic porphyrin dimers **2M** in good yields.^[4]


It has been recognized that Pd-catalyzed oxidative coupling of acetylene can produce different products from the usual Cu-mediated reaction. As a representative example, Haley and co-workers have reported metal-dependent selective synthesis of two different dehydrobenzoannulenes by the intramolecular coupling reactions of single precursors, which is owing to the difference of their reaction mechanisms.^[5] While still under debate, the mechanism of Cu-mediated oxidative butadiyne synthesis is proposed to involve a dimeric Cu acetylide arranged in a *pseudo-trans* configuration prior to reductive elimination (Scheme 1).^[6] On the other hand, in Pd-catalyzed reactions, two ethynyl groups are proposed to be arranged in a *cis* geometry for reductive elimination.^[7] In contrast to Haley and co-workers intramolecular reactions, metal-catalyzed intermolecular oxidative butadiyne formation is more difficult to control.^[8,9] Herein, we report



Scheme 1. Proposed metal acetylide intermediates in Cu-mediated (left) and Pd-catalyzed (right) reactions.

preferential formation of trimers **3M** by Pd-catalyzed coupling reactions of **1M**.

A solution of **1Ni** in toluene and diisopropylamine was stirred at room temperature in the presence of catalytic amounts of $\text{PdCl}_2(\text{PPh}_3)_2$ and CuI and two equivalents of 1,4-benzoquinone as the oxidant. Separation by preparative gel permeation chromatography provided **2Ni**^[4] and **3Ni** and in 9.3 and 72 % yields, respectively (Scheme 2).^[10] The MALDI-TOF mass spectrum of **3Ni** displayed a parent ion peak at 2933.39 (calcd for $\text{C}_{198}\text{H}_{210}\text{N}_{12}\text{Ni}_3$, m/z 2933.49 $[M]^+$). The



Scheme 2. Chemical formulas of **1M**, **2M**, **3M**, and **4Ni**. Ar = 3,5-di-*tert*-butylphenyl. M = Ni, Zn, 2H.

[*] S. Tokuji, Prof. Dr. H. Yorimitsu, Prof. Dr. A. Osuka
Department of Chemistry, Graduate School of Science,
Kyoto University
Sakyo-ku, Kyoto 606-8502 (Japan)
E-mail: yori@kuchem.kyoto-u.ac.jp
osuka@kuchem.kyoto-u.ac.jp

[**] This work was supported by MEXT through Grants-in-Aid (Nos. 22245006 (A), 20108001 “ π -Space”, and 24106721 “Reaction Integration”). S.T. acknowledges a JSPS Fellowship for Young Scientists. We thank Prof. Dr. H. Maeda and Y. Bando of Ritsumeikan University for MALDI-TOF MS measurement.

Supporting information for this article is (experimental details) available on the WWW under <http://dx.doi.org/10.1002/anie.201207763>.

^1H NMR spectrum of **3Ni** in CDCl_3 revealed a singlet signal at 10.35 ppm owing to the *meso* proton, reflecting its highly symmetric structure. Under similar concentrations, treatment of **1Ni** with $\text{Cu}(\text{OAc})_2$ in pyridine/tetrahydrofuran (THF) provided **2Ni** in 82 % yield with a trace amount of **3Ni**.^[4] Similar results were obtained for a Zn^{II} porphyrin substrate **1Zn**; the Pd-catalyzed reaction provided **2Zn** in 24 % yield and **3Zn** in 60 % yield, while the Cu-mediated reaction yielded **2Zn** in 78 % yield with a trace amount of **3Zn**. Demetalation of **3Zn** with hydrochloric acid provided **3H** in 54 % yield.

The triangular structure of **3Ni** has been confirmed by X-ray crystallographic analysis (Figure 1).^[11] The average deviation from the mean planes composed of twenty-four atoms of porphyrin cores is 0.21 Å. The averaged bond angles of $\text{C}(\text{sp}^2)-\text{C}_\alpha=\text{C}_\beta$ and $\text{C}_\alpha=\text{C}_\beta-\text{C}_\gamma$ (θ_1 and θ_2) of six non-equivalent $\text{C}\equiv\text{C}$ units are 170.0° and 172.5° , respectively, hence showing a distinct deviation from the linear structure, and the average $\text{C}_\alpha-\text{C}_\alpha$ and $\text{C}_\beta-\text{C}_\beta$ distances (d_α and d_β) are 5.50 and 6.31 Å, respectively. These data indicate that the structural distortion is concentrated at the 1,3-butadiyne bridges in **3Ni**. We have also determined the X-ray crystal structures of **1Ni** and **2Ni**.^[11] The distances d_α and d_β in **2Ni** are 5.29 Å and 5.37 Å, respectively, which are close to those in **1Ni** ($d_\alpha=5.21$ Å and $d_\beta=5.34$ Å). The averaged bond angles (θ_1 and θ_2) are 173.7° and 176.8° in **2Ni**, respectively, indicating there is less structural distortion.

Interestingly, trimer **3Ni** can be used to prepare the 2,5-thienylene-bridged triporphyrin **4Ni**. Simple heating of **3Ni** in the presence of $\text{Na}_2\text{S}\cdot 9\text{H}_2\text{O}$ in *p*-xylene/2-methoxyethanol gave **4Ni** in 77 % yield as the sole product (Scheme 2).^[12] Notably, the free base trimer **4H** was prepared by Suzuki–Miyaura coupling of β,β' -diborylporphyrin with 2,5-dibromothiophene, in 10 % yield along with many other porphyrin macrocycles.^[13,14]

Figure 2 shows the UV/Vis absorption and fluorescence spectra of **1Zn–3Zn** measured in CH_2Cl_2 . **3Zn** displays a highly perturbed absorption spectrum, which is similar to the case of **2Zn**. The Soret band of **3Zn** is split into two peaks at 452 and 487 nm and Q-bands are observed at 534, 566, 580 and 603 nm with a shoulder around 620 nm. This shoulder peak of **3Zn** is red-shifted as compared with that of **1Zn** (around 600 nm) but blue-shifted from that of **2Zn** (around 645 nm). The fluorescence spectrum of **3Zn** shows peak maxima at 630 and 689 nm, which are blue shifted by 27 and 43 nm, respectively, as compared with those of **2Zn** (Figure 2b). Interestingly, the fluorescence quantum yield Φ_F of **3Zn** was determined by a sphere photon-counting apparatus to be 0.056, larger than those of **1Zn** ($\Phi_F=0.025$) and **2Zn** ($\Phi_F=0.009$).^[4] The blue shifts in the absorption and fluorescent spectra of **3Zn** may be ascribed to its less effective π conjugation as compared with **2Zn**, probably due to its distorted 1,3-butadiyne structure.

These Cu-mediated and Pd-catalyzed reactions are believed to proceed in a stepwise manner via an acyclic intermediate **2M_AC** or its related species (Scheme 3).^[15] Intramolecular cyclization of **2M_AC** should be smooth in the Cu-mediated reactions through an easily available pseudo-*trans* geometry, but might be slow in the Pd-catalyzed




Figure 1. X-ray crystal structures. a) Top view and b) side view of **3Ni**, c) top view and d) side view of **2Ni**, e) top view and f) side view of **1Ni**. Thermal ellipsoids are shown at the 30 % probability level. Solvent molecules and 3,5-di-*tert*-butylphenyl groups are omitted for clarity. Light gray spheres = Ni atoms.

reactions owing to difficulty in forming the requisite *cis* geometry, which would drive the intermolecular coupling reaction of **2M_AC** with another **1M**, to give the intermediate **3M_AC**. Then, the macrocyclic ring closure of **3M_AC** can proceed better, probably because **3M_AC** allows formation of the *cis* geometry that is key for the reductive elimination.^[16] To confirm the stepwise mechanism, **2Ni_AC** was prepared separately (see Supporting Information) and its reactions were examined. While the Cu-mediated oxidative coupling of **2Ni_AC** provided **2Ni** in 74 %, the yield of **2Ni**






Figure 2. a) UV/Vis absorption spectra and b) fluorescence spectra of 1Zn, 2Zn, and 3Zn in CH₂Cl₂. ε = molar extinction coefficient.



Scheme 3. Plausible reaction mechanisms of the Cu-mediated and Pd-catalyzed reactions. Ar = 3,5-di-*tert*-butylphenyl. M = Ni, Zn.

dropped to 44% with various polymeric products in the Pd-catalyzed reaction. The Pd-catalyzed reaction of 1Ni with an equimolar amount of 2Ni_{AC} gave 2Ni and 3Ni in 20% and 45% yield, respectively.^[17] These results are consistent with the stepwise mechanism.

Finally, this strategy was applied to a *meso-meso* directly linked diporphyrin 5Zn. The Cu-mediated reaction of 5Zn produced predominantly 6Zn in 86% yield, while the Pd-catalyzed reaction of 5Zn provided the porphyrin hexamer 7Zn preferentially in 54% yield along with 6Zn in 19% yield (Scheme 4). These contrasting results indicate the generality of this synthetic method. The hexamer 7Zn is attractive because of its functional similarity to the natural light-



Scheme 4. Chemical formulas of 5Zn, 6Zn, and 7Zn. Ar = 3,5-di-*tert*-butylphenyl.

harvesting antenna in terms of possible excitation energy transfer from the peripheral porphyrins to the trimeric porphyrin core. The absorption spectra of 6Zn and 7Zn are roughly similar to those of 2Zn and 3Zn with additional contributions of the excitonically coupled *meso*-linked peripheral porphyrins (Figure 3a). The fluorescence spec-




Figure 3. a) UV/Vis absorption spectra and b) fluorescence spectra of 5Zn, 6Zn, and 7Zn in CH₂Cl₂.

trum of **7Zn** exhibits bands at 643 and 702 nm (Figure 3b), which are distinctly blue-shifted as compared with those of **6Zn** (674 and 747 nm), a similar trend as was observed for **2Zn** and **3Zn**. The fluorescence quantum yields are 0.038, 0.036, and 0.06 for **5Zn**, **6Zn**, and **7Zn**, respectively, again indicating that the trimeric coupling product is the most fluorescent. Collectively, these data indicate that the electronic conjugation along the 1,3-butadiyne bridges is stronger than the exciton coupling in *meso-meso*-linked diporphyrin segments.

In summary, the Pd-catalyzed oxidative coupling of 2,18-diethynylporphyrins **1M** led to the preferential formation of 1,3-butadiyne-bridged porphyrin trimers **3M**, which is complementary to Cu-mediated macrocyclization. The application of this synthetic strategy to **5Zn** provided hexaporphyrin **7Zn** as a novel light-harvesting antenna model. This strategy may be applied to other diethynylated compounds, which are actively being studied in our laboratory.

Received: September 26, 2012

Published online: November 9, 2012

Keywords: alkyne coupling · butadiyne bridge · palladium · porphyrinoids · triporphyrin synthesis

- [1] a) S. Anderson, H. L. Anderson, J. K. M. Sanders, *Acc. Chem. Res.* **1993**, 26, 469; b) J. Li, A. Ambroise, S. I. Yang, J. R. Diers, J. Seth, C. R. Wack, D. F. Bocian, D. Holten, J. S. Lindsey, *J. Am. Chem. Soc.* **1999**, 121, 8927; c) Y. Kuramochi, A. Satake, Y. Kobuke, *J. Am. Chem. Soc.* **2004**, 126, 8668; d) S. Rucareanu, A. Schuwey, A. Gossauer, *J. Am. Chem. Soc.* **2006**, 128, 3396; e) Y. Nakamura, N. Aratani, A. Osuka, *Chem. Soc. Rev.* **2007**, 36, 831; f) N. Aratani, D. Kim, A. Osuka, *Acc. Chem. Res.* **2009**, 42, 1922; g) N. Aratani, A. Osuka, in *Handbook of Porphyrin Science, Vol. 1* (Eds.: K. M. Kadish, K. M. Smith, R. Guilard), World Scientific Publishing, Singapore, **2010**, pp. 1.
- [2] a) K. Sugiura, Y. Fujimoto, Y. Sakata, *Chem. Commun.* **2000**, 1105; b) K. Sugiura, *Top. Curr. Chem.* **2003**, 228, 65; c) A. Kato, K. Sugiura, H. Miyasaka, H. Tanaka, T. Kawai, M. Sugimoto, M. Yamashita, *Chem. Lett.* **2004**, 33, 578.
- [3] a) M. Hoffmann, C. J. Wilson, B. Odell, H. L. Anderson, *Angew. Chem.* **2007**, 119, 3183; *Angew. Chem. Int. Ed.* **2007**, 46, 3122; b) M. Hoffmann, J. Kärnbratt, M.-H. Chang, L. M. Herz, B. Albinsson, H. L. Anderson, *Angew. Chem.* **2008**, 120, 5071; *Angew. Chem. Int. Ed.* **2008**, 47, 4993; c) J. K. Sprafke, D. V. Kondratuk, M. Wykes, A. L. Thompson, M. Hoffmann, R. Drevinskas, W. H. Chen, C. K. Yong, J. Kärnbratt, J. E. Bullock, M. Malfois, M. R. Wasielewski, B. Albinsson, L. M. Herz, D. Zigmantas, D. Beljonne, H. L. Anderson, *J. Am. Chem. Soc.* **2011**, 133, 17262; d) M. C. O'Sullivan, J. K. Sprafke, D. V. Kondratuk, C. Rinfrey, T. D. W. Claridge, A. Saywell, M. O. Blunt, J. N. O'Shea, P. H. Beton, M. Malfois, H. L. Anderson, *Nature* **2011**, 469, 72; e) J. K. Sprafke, B. Odell, T. D. W. Claridge, H. L. Anderson, *Angew. Chem.* **2011**, 123, 5687; *Angew. Chem. Int. Ed.* **2011**, 50, 5572; f) D. V. Kondratuk, L. M. A. Perdigo, M. C. O'Sullivan, S. Svatek, G. Smith, J. N. O'Shea, P. H. Beton, H. L. Anderson, *Angew. Chem.* **2012**, 124, 6800; *Angew. Chem. Int. Ed.* **2012**, 51, 6696.
- [4] I. Hisaki, S. Hiroto, K. S. Kim, S. B. Noh, D. Kim, H. Shinokubo, A. Osuka, *Angew. Chem.* **2007**, 119, 5217; *Angew. Chem. Int. Ed.* **2007**, 46, 5125.
- [5] a) J. A. Marsden, J. J. Miller, M. M. Haley, *Angew. Chem.* **2004**, 116, 1726; *Angew. Chem. Int. Ed.* **2004**, 43, 1694; b) J. A. Marsden, J. J. Miller, L. D. Shirtcliff, M. M. Haley, *J. Am. Chem. Soc.* **2005**, 127, 2464; c) J. A. Marsden, M. M. Haley, *J. Org. Chem.* **2005**, 70, 10213; d) C. A. Johnson II, M. M. Haley, *Organometallics* **2005**, 24, 1161; e) E. L. Spitler, J. M. Monson, M. M. Haley, *J. Org. Chem.* **2008**, 73, 2211; f) T. Takeda, A. G. Fix, M. M. Haley, *Org. Lett.* **2010**, 12, 3824.
- [6] a) F. Bohlmann, H. H. Schönowsky, E. Inhoffen, G. Grau, *Chem. Ber.* **1964**, 97, 794; b) P. Siemsen, R. C. Livingston, F. Diederich, *Angew. Chem.* **2000**, 112, 2740; *Angew. Chem. Int. Ed.* **2000**, 39, 2632; c) L. Fomina, B. Vazquez, E. Tkatchouk, S. Fomine, *Tetrahedron* **2002**, 58, 6741.
- [7] a) A. Gillie, J. K. Stille, *J. Am. Chem. Soc.* **1980**, 102, 4933; b) F. Ozawa, T. Ito, Y. Nakamura, A. Yamamoto, *Bull. Chem. Soc. Jpn.* **1981**, 54, 1868; c) J. J. Low, W. A. Goddard III, *J. Am. Chem. Soc.* **1986**, 108, 6115; d) J. M. Brown, N. A. Cooley, *Chem. Rev.* **1988**, 88, 1031.
- [8] S. S. Jester, N. Shabelina, S. M. L. Blanc, S. Höger, *Angew. Chem.* **2010**, 122, 6237; *Angew. Chem. Int. Ed.* **2010**, 49, 6101.
- [9] In the case of dibenz[*a,j*]anthracene-based macrocycle formations, dimers were the main products; J. M. W. Chan, J. R. Tischler, S. E. Kooi, V. Bulovic, T. M. Swager, *J. Am. Chem. Soc.* **2009**, 131, 5659.
- [10] Use of a catalyst bearing a *cis*-bidentate ligand PdCl₂(dppf) led to decreased yields of **3Ni** (14%) and **2Ni** (18%) along with polymerized products (see Supporting Information).
- [11] Crystallographic data for **3Ni**: C₁₉₈H₂₁₀N₁₂Ni₃, *M_w* = 2933.91, monoclinic, space group *C2/c* (No. 15), *a* = 65.51(3), *b* = 24.309(7), *c* = 31.96(2) Å, β = 113.67(5)°, *V* = 46614(43) Å³, *Z* = 8, *D_{calc}* = 0.836 g cm⁻³, *T* = 93(2) K, *R₁* = 0.1121 (*I* > 2.0σ(*I*)), *R_w* = 0.2532 (all data), GOF = 1.151 (*I* > 2.0σ(*I*)). Some unassigned electron density owing to severely disordered solvent was removed by using the utility SQUEEZE in the PLATON software package;^[18] Crystallographic data for **2Ni**: C₁₃₂H₁₄₀N₈Ni₂, 6(C₆H₄Cl₂), *M_w* = 1418.95, triclinic, space group *P*-1 (No. 2), *a* = 11.9002(2), *b* = 16.0714(3), *c* = 20.3575(4) Å, α = 88.0309(10), β = 74.0268(10), γ = 79.8791(16)°, *V* = 3684.46(12) Å³, *Z* = 2, *D_{calc}* = 1.279 g cm⁻³, *T* = 93(2) K, *R₁* = 0.0949 (*I* > 2.0σ(*I*)), *R_w* = 0.2843 (all data), GOF = 1.073 (*I* > 2.0σ(*I*)); Crystallographic data for **1Ni**: C₆₆H₇₂N₄Ni, 2.5(C₂H₄Cl₂), *M_w* = 1227.37, triclinic, space group *P*-1 (No. 2), *a* = 10.6798(4), *b* = 17.1731(6), *c* = 18.9668(6) Å, α = 70.036(2), β = 81.909(2), γ = 87.581(2)°, *V* = 3236.97(19) Å³, *Z* = 2, *D_{calc}* = 1.259 g cm⁻³, *T* = 93(2) K, *R₁* = 0.0954 (*I* > 2.0σ(*I*)), *R_w* = 0.2443 (all data), GOF = 1.037 (*I* > 2.0σ(*I*)). CCDC 899883 (**3Ni**), 899882 (**2Ni**) and 899881 (**1Ni**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [12] For transformation of 1,3-butadiynes to thiophenes, see a) J. Krömer, I. Rios-Carreras, G. Fuhrmann, C. Musch, M. Wunderlin, T. Debaerdemaeker, E. Mena-Osteritz, P. Bäuerle, *Angew. Chem.* **2000**, 112, 3623; *Angew. Chem. Int. Ed.* **2000**, 39, 3481; b) J. Lagan, S. K. Arora, *J. Org. Chem.* **1983**, 48, 4317; c) N. Sumi, H. Nakanishi, S. Ueno, K. Takimiya, Y. Aso, T. Otsubo, *Bull. Chem. Soc. Jpn.* **2001**, 74, 979; d) M. J. O'Connor, M. M. Haley, *Org. Lett.* **2008**, 10, 3973; e) C. Maeda, T. Yoneda, N. Aratani, M.-C. Yoon, J. M. Lim, D. Kim, N. Yoshioka, A. Osuka, *Angew. Chem.* **2011**, 123, 5809; *Angew. Chem. Int. Ed.* **2011**, 50, 5691.
- [13] J. Song, S. Y. Jang, S. Yamaguchi, J. Sankar, S. Hiroto, N. Aratani, J.-Y. Shin, S. Easwaramoorthi, K. S. Kim, D. Kim, H. Shinokubo, A. Osuka, *Angew. Chem.* **2008**, 120, 6093; *Angew. Chem. Int. Ed.* **2008**, 47, 6004.
- [14] Suzuki-Miyaura coupling of β,β'-diboryl Ni^{II} porphyrin with bromo(*tert*-butyldimethylsilyl)acetylene and subsequent desilylation with tetrabutylammonium fluoride (TBAF) provided **1Ni** in 72% yield (Supporting Information).

- [15] These Pd-catalyzed or Cu-mediated reactions are different from the Bäuerle's case, in which pre-organization of Pt complexes enabled macrocycle formations. See Ref. [12a].
- [16] For ring-closure processes, Pd-catalyzed acetylene coupling reactions were sometimes adopted. See Refs [3,5] and a recent review; M. Iyoda, J. Yamakawa, M. J. Rahman, *Angew. Chem.* **2011**, *123*, 10708; *Angew. Chem. Int. Ed.* **2011**, *50*, 10522.
- [17] The yields were calculated on the basis of the number of porphyrin subunits.
- [18] a) A. L. Spek, *PLATON, A Multipurpose Crystallographic Tool*, Utrecht University, Utrecht, The Netherlands, **2005**; b) P. van der Sluis, A. L. Spek, *Acta Crystallogr. Sect. A* **1990**, *46*, 194.
-