

Robust Structure and Reactivity of Aqueous Arsenous Acid–Platinum(II) Anticancer Complexes**

Đenana U. Miodragović, Jeremy A. Quentzel, Josh W. Kurutz, Charlotte L. Stern, Richard W. Ahn, Irawati Kandela, Andrew Mazar, and Thomas V. O'Halloran*

Two inorganic drugs, the widely used *cis*-diamminedichloroplatinum(II)^[1] and antileukemia agent arsenic trioxide, are highly successful agents for treatment of cancer. Cisplatin is used in combination chemotherapy to treat ovarian, testicular, head, neck, and bladder cancers.^[2] Unfortunately, these and other cancers frequently develop resistance to this drug and there are intensive efforts to develop new agents that overcome cisplatin resistance.^[3] As₂O₃, discovered as a traditional Chinese medicine, is now an FDA-approved front line treatment for acute promyelocytic leukemia^[4] and has also shown preliminary efficacy in the treatment of blood cancers such as multiple myeloma and myelodysplastic syndromes.^[4a] Both compounds induce apoptotic cell death, but through different pathways: cisplatin reacts with DNA and causes intra- and interstrand DNA cross-links,^[2a,d] whereas at low concentrations arsenous acid, the principal component of aqueous solutions of As₂O₃ at pH 7, can react with and trigger degradation of key zinc-dependent regulatory proteins and also inhibit angiogenesis, migration, and invasion. At higher concentrations it triggers apoptosis^[5,6] through pathways that involve elevated levels of reactive oxygen species (ROS) in mitochondria.^[4a,5,7] Synergistic activity of these drugs has been reported^[8] supporting the idea

that compounds combining both species may have advantages as anticancer therapeutics. The only example of a platinum adduct with arsenous acid in the literature emerged in efforts to develop efficient systems for loading As₂O₃ into liposomes with aquated forms of cisplatin: extended X-ray absorption fine structure (EXAFS) spectroscopy suggested that a new type of Pt^{II}–As^{III} center was stabilized in the nanocrystalline formulation.^[9] Given the absence of structural precedents in the literature, it was not clear that such complexes could exist or would be stable in aqueous solution. Herein we report synthetic routes to a novel family of small-molecule complexes of the aqueous form of As₂O₃ bound directly to Pt^{II} as an As(OH)₂ moiety and demonstrate that their robust molecular structure involves an As^{III} center that acts simultaneously as a Lewis acid and a Lewis base. These arsenoplatins




Figure 1. Thermal ellipsoid plots of a) complex **1a** (**1** crystallizes in two crystal systems) and b) complex **2**. Solvent molecules have been omitted for clarity. The plots are drawn at 50% probability level.^[10]

[*] Dr. Đ. U. Miodragović, J. A. Quentzel, Dr. J. W. Kurutz, C. L. Stern, Dr. R. W. Ahn, Dr. I. Kandela, Prof. Dr. A. Mazar, Prof. Dr. T. V. O'Halloran
Chemistry of Life Processes Institute and Department of Chemistry
Northwestern University
2145 Sheridan Road, Evanston, IL 60208 (USA)
E-mail: t-ohalloran@northwestern.edu
Homepage: <http://www.chemistry.northwestern.edu/faculty/thomas-ohalloran.html>

[**] We thank the NIH (NCI Cancer Nanotechnology Platform Partnerships Grant U01CA151461 and PSOC grant U54CA143869, Center of Cancer Nanotechnology Excellence Grants U54CA119341, P50CA090386, and Core Grant P30CA060553 to the Robert H. Lurie Comprehensive Cancer Center) and the CDMRP Breast Cancer Research Program BC076723 for funding. R. W. Ahn thanks the CDMRP Breast Cancer Research Program Predoctoral Fellowship (BC073413). We thank Prof. M. Djuran, Dr. E. Que, and E. Swindell for valuable discussions.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201303251>.

are stable in solution and exhibit chemical bonding, ligand substitution chemistry, and biological activities that are distinct from the parent compounds and show promising activity in drug-resistant cancer cell lines.

Arsenoplatin **1** (Figure 1) is synthesized by heating cisplatin with As₂O₃ in an acetonitrile/water mixture (9:1, v/v) at 90 °C for three days. The yield increases from 23 % to 75 % when the starting material is K₂[PtCl₄] (see the Supporting Information for syntheses and characterizations of **1–3**).

Variations on this Pt–As core complex are accessible by varying the substituent on the nitrile. For instance, arsenoplatin **2** is obtained from the reaction of K₂[PtCl₄] with As₂O₃ in the presence of propionitrile (Figure 1). Conditions for the synthesis of **2** were different from **1** owing to the different miscibility of propionitrile in water (1:9, v/v). Complex **2** is obtained at room temperature after 4 days, whereas **1** is

obtained at elevated temperatures. In **1** and **2**, the Pt^{II} center adopts a square-planar geometry, with arsenic, chloride, and two nitrogen donors in a *trans* configuration. The nitrogen donors are derived from acetamide (propanamide) formed through Pt-assisted acetonitrile (propionitrile) hydrolysis in situ.^[11] The Pt–N bond lengths in **1a** (Pt1–N1 = 2.000(3) Å and Pt1–N2 = 2.004(3) Å) are consistent with the Pt–N bond lengths obtained in other Pt^{II} complexes with the deprotonated form of acetamide.^[11,12] The N1–C1 and O1–C1 bond lengths (both 1.302(4) Å) and N2–C3 and O2–C3 (1.289(6) and 1.297(6) Å) in **1a** are indicative of a high degree of delocalization^[11] present in the chelate rings formed by bridging N,O acetylamido ligands.

The closest precedents for these arsenous acid/platinum complexes are found in heterometallic clusters where the Ni^{II}^[13] and Pd^{II}^[14] centers bind directly to arsenous acid. In Pd^{II} and Ni^{II} complexes, As(OH)₃ is bound to the metal as a Lewis base with arsenic in a distorted tetrahedral and pyramidal environment, respectively. In arsenoplatins, the geometry at the As^{III} is best described as trigonal bipyramidal (TBP), with the Pt^{II} and two hydroxides binding in the trigonal plane. In **1** and **2**, arsenic retains a formal oxidation state of three, and displays distorted TBP geometry with a PtO₄ coordination sphere. The TBP geometry around As^{III} in metal complexes is uncommon, however it has been observed in complexes of an organoarsenic ligand with Pt^{II}^[15] and in complexes of Ag^I^[16] and Fe^{II}.^[17] While As^{III} can act as either a Lewis base or a Lewis acid,^[18] in **1** and **2** As^{III} acts simultaneously as a Lewis base (As→Pt) and as a Lewis acid (O→As). These multiple intramolecular interactions explain in part the strong Pt–As interaction in arsenoplatins: 2.2687(4) Å in **2**, 2.2729(2) Å in **1b**, and 2.2732(3) Å in **1a** (the shortest Pt–As bond length found in the CSD is 2.267(2) Å in one organoarsenic compound).^[19]

Heteronuclear NMR spectroscopy reveals that the strong Pt–As interaction observed in the solid state persists in aqueous solution (Figures S2–S7 in the Supporting Information). Owing to symmetry, the ¹H NMR spectrum of **1** contains only one NH signal at 8.16 ppm and one OH signal at 8.92 ppm; these are the first chemical shifts reported for an M–As–OH moiety. The ¹⁹⁵Pt chemical shift for **1** (–3589 ppm) lies between the signals for Pt^{II} diamines (e.g., cisplatin at –2097 ppm)^[20] and those of mixed Pt^{II} arsine halides (e.g., [Pt{o-C₆H₄(AsMe₂)₂]Cl₂] at –4556 ppm),^[21] consistent with a Pt^{II}N₂ClAs coordination sphere.

This Pt–As core is also stable to ligand substitution reactions. In general, hydrolysis of Pt–Cl bonds is slow (*t*_{1/2} = 2 h at 37 °C and 4 mM Cl[–]),^[22] and rapid substitution usually requires addition of reagents such as AgNO₃. The substitution of the Cl[–] ligand in **1** with SCN[–] in water occurs immediately at room temperature, likely driven by the *trans* effect of the arsenic moiety.^[23] NMR spectroscopy in solution and X-ray crystallography confirm that the Pt–As bond remained intact. Crystals suitable for a single-crystal X-ray analysis were obtained when this complex (**3**) was synthesized in a 1:1 water/methanol mixture (Figure 2). NMR spectroscopy reveals facile linkage isomerization of **3** in solution at room temperature (Figures S8–S12 in the Supporting Information). Specifically, upon dissolving **3** in [D₆]DMSO solution, an




Figure 2. Thermal ellipsoid plot of **3**. The plot is drawn at 50% probability level.

equilibrium mixture of (64 ± 1.2)% of S isomer and (36 ± 1.5)% of N isomer is quickly established, i.e., the ¹H NMR spectrum obtained after 5 min upon dissolution of **3** does not change over time. Assignments of chemical shifts for the S- and N-bound isomers are based on two-dimensional ¹⁹⁵Pt (Figure 3) and ¹⁵N NMR spectroscopy on a sample of **3**, which was synthesized using thiocyanate enriched in ¹³C and ¹⁵N at 99% (see the Supporting Information).

Initial formation of **3** with S-bound thiocyanate can be kinetically or thermodynamically controlled, but both isomers [Eq. (1)] are sufficiently stable in [D₆]DMSO solution to be





Figure 3. a) ¹H–¹³C HSQC NMR spectrum of **3** (with ¹³C/¹⁵N) in [D₆]DMSO, acquired at 25 °C at 600 MHz ¹H with high resolution in the indirect (¹³C) dimension to distinguish the methyl resonances of the N and S isomers; b) ¹⁹⁵Pt NMR spectrum of **3** (with ¹³C/¹⁵N) in [D₆]DMSO, referenced indirectly to ¹H TMS such that Na₂¹⁹⁵PtCl₆ resonates at 0.0 ppm. The 456 Hz splitting of the ¹⁹⁵Pt peak at –3724 ppm arises from scalar coupling to the SCN[–] ¹⁵N.

observed using NMR spectroscopy. Analysis of variable-temperature NMR experiments reveals the thermodynamics of this facile linkage isomer equilibrium. A Van't Hoff plot of the temperature-dependent NMR spectra (Figure S13 in the Supporting Information) reveals an equilibrium constant K_{eq} for isomerization of 0.563, $\Delta H^\circ = -15.7 \text{ kJ mol}^{-1}$, $\Delta S^\circ = -57.5 \text{ J mol}^{-1} \text{ K}^{-1}$, and $\Delta G^\circ = 1.42 \text{ kJ mol}^{-1}$ ($[\text{D}_6]\text{DMSO}$ solution, 25°C). These parameters indicate a low barrier to substitution at the Pt^{II} site *trans* to the As^{III} ligand, consistent with a very strong *trans* effect of the $\text{As}^{\text{III}}(\text{OH})_2$ moiety. On the basis of our thermodynamic data, the N isomer is enthalpically favored in solution by 15.7 kJ mol^{-1} . Interestingly, only the S-linked complex could be isolated in the solid state, which may be the result of both rapid equilibration and a lower solubility for the S isomer.



Complex **1** demonstrates significant anticancer activity in a panel of human cancer cell lines (Table S10 in the Supporting Information) and also overcomes one of the most significant limitations of platinum drugs, namely tumor-based drug resistance mechanisms. The ovarian cisplatin-resistant A2780^{CP} cancer cell line is of special interest, since it encompasses all of the known major mechanisms of resistance to cisplatin (reduced uptake, increased level of glutathione, increased DNA repair, and tolerance to Pt^{II} -induced lesions).^[24] The results show that **1** exhibited more than twice the cytotoxicity of cisplatin against the cisplatin-resistant cell line A2780^{CP} [$\text{IC}_{50} = (21.4 \pm 1.8) \mu\text{M}$ versus $(47.3 \pm 2.1) \mu\text{M}$, Figure 4]. The ability of **1** to circumvent cisplatin-acquired resistance was determined from the resistance factor (RF), and an RF value of < 2 denotes no cross-resistance.^[25] In the case of ovarian A2780 and A2780^{CP} cell lines all approved platinum drugs have RFs between 6.1 and 16.0.^[24,25] The RF of 1.1 for **1** indicates that it is far more

effective at killing this cisplatin-resistant cancer cell line and may be able to bypass drug resistance mechanism(s) that lower cisplatin cytotoxicity.

Complex **1** showed better cytotoxic activity than either cisplatin or As_2O_3 in colon HCT-116 [$\text{IC}_{50} = (1.6 \pm 0.4) \mu\text{M}$ versus $(5.5 \pm 1.3) \mu\text{M}$ and $(9.4 \pm 0.9) \mu\text{M}$] and glioblastoma U-87 [$\text{IC}_{50} = (0.37 \pm 0.11) \mu\text{M}$ versus $(9.6 \pm 0.8) \mu\text{M}$ and $(1.6 \pm 2.9) \mu\text{M}$] cancer cell lines (Figure 4). Additionally, **1** showed twice the cytotoxicity of cisplatin against MDA-MB-231-mCherry cells [$\text{IC}_{50} = (9.5 \pm 0.1) \mu\text{M}$ versus $(22.3 \pm 2.8) \mu\text{M}$], as well as improved cytotoxicity compared with As_2O_3 in RPMI 8226 multiple myeloma cells [$\text{IC}_{50} = (4.5 \pm 1.0) \mu\text{M}$ versus $(7.1 \pm 0.2) \mu\text{M}$].

trans-Platinum compounds in comparison with *cis*-compounds display different patterns of ligand substitution, which contributes to the potency of *trans*-platinum compounds in cisplatin-resistant cell lines.^[26] Although we do not have evidence that arsenoplatin compounds target DNA, the distinct biological activity of **1** *in vitro* may be the result of the strong *trans* effect of the $\text{As}(\text{OH})_2$ moiety combined with the *trans* stereochemistry of the N atoms at the platinum center.

In conclusion, the first compounds containing a $\text{Pt-As}(\text{OH})_2$ core (arsenoplatins **1** and **2**) have been synthesized and characterized as robust complexes that are stable in aqueous solution. Single-crystal X-ray structure characterization reveals that these unprecedented compounds contain a very short Pt-As bond with the square-planar coordination expected for Pt^{II} but an atypical five coordinate As^{III} geometry. Intriguingly, the arsenic atom in these complexes exhibits both Lewis acid and Lewis base behavior upon binding to the platinum acetylaminato moiety. The Pt-As core in these complexes readily undergoes ligand exchange reactions at the Pt^{II} center with retention of core bonding and stereochemistry. Both, the rapid substitution of chloride in **1** and isomerism in **3** demonstrate a strong *trans* effect of the arsenic moiety. Complex **1** has significant biological activity in several cancer cell lines and preliminary data are consistent with the ability of arsenoplatins to overcome drug resistance mechanisms. These results are promising and future work with mouse xenograft models should help shed more light on the real potential of this unique class of compounds.




Figure 4. Dose-response curves for a) ovarian cisplatin-resistant A2780^{CP}, b) colon HCT-116, and c) glioblastoma U-87 cancer cell lines after exposure to **1** (red ■), cisplatin (blue ●), and As_2O_3 (green ▲).

Received: April 17, 2013
Revised: July 4, 2013
Published online: August 22, 2013

Keywords: antitumor agents · arsenic · drug design · platinum · X-ray diffraction

- [1] B. Rosenberg, L. Vancamp, J. E. Trosko, V. H. Mansour, *Nature* **1969**, 222, 385–386.
- [2] a) Z. H. Siddik, *Oncogene* **2003**, 22, 7265–7279; b) L. Kelland, *Nat. Rev. Cancer* **2007**, 7, 573–584; c) N. J. Wheate, S. Walker, G. E. Craig, R. Oun, *Dalton Trans.* **2010**, 39, 8113–8127; d) D. Wang, S. J. Lippard, *Nat. Rev. Drug Discovery* **2005**, 4, 307–320.
- [3] a) N. Graf, S. J. Lippard, *Adv. Drug Delivery Rev.* **2012**, 64, 993–1004; b) A. Casini, J. Reedijk, *Chem. Sci.* **2012**, 3, 3135–3144.
- [4] a) A. Emadi, S. D. Gore, *Blood Rev.* **2010**, 24, 191–199; b) X.-W. Zhang, X.-J. Yan, Z.-R. Zhou, F.-F. Yang, Z.-Y. Wu, H.-B. Sun, W.-X. Liang, A.-X. Song, V. Lallemand-Breitenbach, M. Jeanne, Q.-Y. Zhang, H.-Y. Yang, Q.-H. Huang, G.-B. Zhou, J.-H. Tong, Y. Zhang, J.-H. Wu, H.-Y. Hu, H. de The, S.-J. Chen, Z. Chen, *Science* **2010**, 328, 240–243.
- [5] L. C. Platanias, *J. Biol. Chem.* **2009**, 284, 18583–18587.
- [6] a) S. J. Ralph, *Metal-Based Drugs*, **2008**, 260146; b) M. R. Litzow, *Expert Opin. Pharmacother.* **2008**, 9, 1773–1785.
- [7] S. Fulda, L. Galluzzi, G. Kroemer, *Nat. Rev. Drug Discovery* **2010**, 9, 447–464.
- [8] a) N. Zhang, Z.-M. Wu, E. McGowan, J. Shi, Z.-B. Hong, C.-W. Ding, P. Xia, W. Di, *Cancer Sci.* **2009**, 100, 2459–2464; b) H. Li, X. Zhu, Y. Zhang, J. Xiang, H. Chen, *J. Exp. Clin. Cancer Res.* **2009**, 28, 110.
- [9] H. Chen, S. Pazicni, N. L. Krett, R. W. Ahn, J. E. Penner-Hahn, S. T. Rosen, T. V. O'Halloran, *Angew. Chem.* **2009**, 121, 9459–9463; *Angew. Chem. Int. Ed.* **2009**, 48, 9295–9299.
- [10] CCDC 831785 (**1a**), CCDC 831786 (**1b**), CCDC 831788 (**2**) and CCDC 831787 (**3**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [11] A. Erxleben, I. Mutikainen, B. Lippert, *J. Chem. Soc. Dalton Trans.* **1994**, 3667–3675.
- [12] E. J. Ziolkowski, P. Turner, L. M. Rendina, *Inorg. Chem. Commun.* **2006**, 9, 53–56.
- [13] R. Hernández-Molina, M. N. Sokolov, M. Clausen, W. Clegg, *Inorg. Chem.* **2006**, 45, 10567–10575.
- [14] M. N. Sokolov, A. V. Virovets, D. N. Dybtsev, E. V. Chubarova, V. P. Fedin, D. Fenske, *Inorg. Chem.* **2001**, 40, 4816–4817.
- [15] I. M. Müller, W. S. Sheldrick, *Eur. J. Inorg. Chem.* **1998**, 1999–2003.
- [16] R. C. Bott, G. Smith, D. S. Sagatys, D. E. Lynch, C. H. L. Kennard, *Aust. J. Chem.* **2000**, 53, 917–924.
- [17] Y. Yamamoto, K. Toyota, Y. Wakisaka, K.-y. Akiba, *Heteroat. Chem.* **2000**, 11, 42–47.
- [18] a) T. M. Klapötke, H. Nöth, T. Schütt, M. Suter, *Eur. J. Inorg. Chem.* **2002**, 2511–2517; b) T. G. Carter, W. J. Vickaryous, V. M. Cangelosi, D. W. Johnson, *Comments Inorg. Chem.* **2007**, 28, 97–122; c) N. J. Hill, W. Levason, G. Reid, *J. Chem. Soc. Dalton Trans.* **2002**, 1188–1192.
- [19] M. K. Cooper, P. J. Guernsey, H. J. Goodwin, M. McPartlin, *J. Chem. Soc. Dalton Trans.* **1982**, 757–764.
- [20] B. M. Still, P. G. A. Kumar, J. R. Aldrich-Wright, W. S. Price, *Chem. Soc. Rev.* **2007**, 36, 665–686.
- [21] E. G. Hope, W. Levason, N. A. Powel, *Inorg. Chim. Acta* **1986**, 115, 187–192.
- [22] a) S. J. B.-P. a. T. G. Appleton, *Platinum-Based Drugs in Cancer Therapy*, Humana, Totowa, **2000**; b) S. van Zutphen, E. Pantoja, R. Soriano, C. Soro, D. Tooke, A. Spek, H. Dulk, J. Brouwer, J. Reedijk, *Dalton Trans.* **2006**, 1020–1023.
- [23] N. Kuznik, O. F. Wendt, *J. Chem. Soc. Dalton Trans.* **2002**, 3074–3078.
- [24] H. B. J. Helleman, I. H. L. Hamelers, A. W. M. Boersma, A. I. P. M. de Kroon, G. Stoter, K. Nooter, *Cancer Biol. Ther.* **2006**, 5, 943.
- [25] F. P. Intini, R. Z. Pellicani, A. Boccarelli, R. Sasanelli, M. Coluccia, G. Natile, *Eur. J. Inorg. Chem.* **2008**, 4555–4561.
- [26] a) U. Kalinowska-Lis, J. Ochocki, K. Matlawska-Wasowska, *Coord. Chem. Rev.* **2008**, 252, 1328–1345; b) J. Reedijk, *Eur. J. Inorg. Chem.* **2009**, 1303–1312; c) S. Chen, D. Xu, H. Jiang, Z. Xi, P. Zhu, Y. Liu, *Angew. Chem.* **2012**, 124, 12424–12428; *Angew. Chem. Int. Ed.* **2012**, 51, 12258–12262.