

An Approach to Benzophosphole Oxides through Silver- or Manganese-Mediated Dehydrogenative Annulation Involving C–C and C–P Bond Formation**

Yuto Unoh, Koji Hirano, Tetsuya Satoh,* and Masahiro Miura*

Benzophosphole derivatives have recently attracted much attention in the field of materials chemistry because of their unique optical and electronic properties.^[1] However, synthetic approaches to such promising frameworks are so far limited.^[2] Most of the currently available methods rely on the cyclization of phenylphosphorus compounds with alkynyl groups preinstalled in the *ortho* position; these precursors are usually prepared through complicated multistep reactions. Moreover, the substituent tolerance is low because the cyclizations are performed under strongly basic conditions. To expand the application of benzophosphole species, the development of new, effective processes for synthesizing them from relatively simple starting materials is much needed.

As one promising route from readily available substrates to fused (hetero)aromatic compounds, the rhodium-catalyzed dehydrogenative annulation of monosubstituted benzenes with alkynes through directed C–H bond cleavage has been significantly developed in recent years (Scheme 1).^[3,4] For example, both we^[5] and Fagnou and co-workers^[3c] have reported that indenone imines and indoles can be constructed


through the coupling of benzyldeneanilines and *N*-acetylanilines, respectively, with internal alkynes through C–H and N–H bond cleavages (–LH = –CH=NAr and –NHAc in Scheme 1).

In light of these results, we undertook to develop a direct method for the construction of benzophospholes by using rhodium catalysis. As an initial attempt, we examined the coupling of diphenylphosphine oxide (**1a**) with diphenylacetylene (**2a**) in the presence of a cationic Cp*Rh^{III} complex and AgOAc as catalyst and oxidant, respectively (–LH = –PH(=O)Ph, R¹ = R² = Ph in Scheme 1 and entry 1 in Table 1),^[6,7] and succeeded in obtaining the expected benzophosphole derivative **3aa** in a good yield. It was somewhat surprising to

Table 1: Reaction of diphenylphosphine oxide (**1a**) with diphenylacetylene (**2a**).^[a]

Entry	Oxidant (mmol)	Solvent	T [°C]	Yield of 3aa [%] ^[b]
1 ^[c]	AgOAc (1)	DMF	100	93
2	AgOAc (1)	DMF	100	96 (96)
3 ^[d]	AgOAc (1)	DMF	100	80
4	AgOAc (0.5)	DMF	100	6
5	AgNO ₃ (0.05) + K ₂ S ₂ O ₈ (1)	DMF	100	19
6	AgOAc (1)	DMF	80	13
7	AgOAc (1)	AcOH	100	81
8 ^[e]	AgOAc (12)	DMF	100	(82)
9	Mn(OAc) ₃ ·2H ₂ O (1)	DMF	100	50
10	Mn(OAc) ₃ ·2H ₂ O (1)	DMF	RT	70
11	Mn(OAc) ₃ ·2H ₂ O (1)	AcOH	RT	78
12	Mn(OAc) ₃ ·2H ₂ O (1)	CH ₂ Cl ₂	RT	58

[a] Reaction conditions: **1a** (0.5 mmol), **2a** (0.25 mmol), and oxidant in solvent (3 mL) under N₂ for 4 h. [b] Yield based on the amount of **2a** used as determined by GC. Values in parentheses indicate yield after purification. [c] With [Cp*Rh(MeCN)₃][SbF₆]₂ (0.01 mmol). Cp* = pentamethylcyclopentadienyl. [d] With **1a** (0.25 mmol). [e] With **1a** (6 mmol) and **2a** (3 mmol) in DMF (36 mL).



Scheme 1. Rhodium-catalyzed dehydrogenative annulation. Shown are examples of products from previous reports, and the benzophosphole oxides described in this work.

[*] Y. Unoh, Dr. K. Hirano, Prof. Dr. T. Satoh, Prof. Dr. M. Miura
Department of Applied Chemistry, Faculty of Engineering
Osaka University
Suita, Osaka 565-0871 (Japan)
E-mail: satoh@chem.eng.osaka-u.ac.jp
miura@chem.eng.osaka-u.ac.jp
Homepage: <http://www.chem.eng.osaka-u.ac.jp/~miura-lab/index-Eng.htm>


Prof. Dr. T. Satoh
JST, ACT-C
4-1-8 Honcho, Kawaguchi, Saitama 332-0012 (Japan)

[**] This work was partly supported by Grants-in-Aid from MEXT, JSPS, and JST (Japan).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201307211>.

observe that the reaction proceeded efficiently even without any rhodium catalyst (see below).

Based on this intriguing observation, we postulated an annulation mechanism with the silver promoter alone as illustrated in Scheme 2. In the presence of the Ag^I salt, a P-centered radical appears to be formed from **1a** through homolytic P–H bond cleavage, as proposed previously.^[8] Subsequently, the addition of the radical across the triple



Scheme 2. Proposed mechanism for silver-mediated annulation of **1a** with **2**.

bond of an alkyne **2** followed by cyclization on the phenyl moiety of **1a** may occur to form the benzophosphole oxide **3**. In contrast to well-established reactions involving the addition of P-centered radicals to alkenes, an approach that has been widely employed for synthesizing organophosphorus compounds, intermolecular addition to alkynes has been little explored.^[9] In particular, examples of cascade processes involving successive reactions of the resulting alkenyl radicals, aside from simple reduction, are so far limited.^[10] Consequently, we investigated the annulation reaction in detail to provide an unprecedented direct route to benzophosphole derivatives. Furthermore, it was found that these radical processes can also be promoted by a Mn^{III} salt^[10a,11] as well as Ag^I.

In a typical experiment, diphenylphosphine oxide (**1a**; 0.5 mmol) was treated with diphenylacetylene (**2a**; 0.25 mmol) in the presence of AgOAc (1 mmol) in DMF (3 mL) at 100 °C for 4 h under N₂. The oxidative annulation product 1,2,3-triphenyl-1*H*-phosphindole-1-oxide (**3aa**) was obtained exclusively, in 96 % yield (entry 2 in Table 1). While the reaction using the substrates in a 1:1 ratio gave **3aa** in 80 % yield (entry 3 in Table 1), decreasing the amount of AgOAc (0.5 mmol) substantially reduced the product yield (entry 4). The combination of AgNO₃ (0.05 mmol) and K₂S₂O₈ (1 mmol), which has been employed to generate P-centered radicals,^[8b,c] was not effective for the present reaction (entry 5 in Table 1). At 80 °C, the reaction was sluggish (entry 6 in Table 1). In AcOH, the reaction also proceeded smoothly, but the product yield was somewhat low (entry 7 in Table 1). Note that the present reaction could be readily scaled up to a gram scale. Thus, from **1a** (6 mmol) and **2a** (3 mmol), **3aa** was obtained in 82 % yield (0.93 g, entry 8 in Table 1).


As shown in Scheme 2, the present reaction seems to proceed through a radical process promoted by Ag^I. To obtain additional support for this mechanism, we conducted the reaction of **1a** with **2a** in the presence of radical inhibitors (see the Supporting Information). As expected, adding 2,2,6,6-tetramethylpiperidine-1-oxyl (TEMPO) or 2,6-di-*tert*-butyl- α -(3,5-di-*tert*-butyl-4-oxo-2,5-cyclohexadiene-1-ylidene)-*p*-tolyl-oxyl (Galvinoxyl) severely retarded or completely suppressed the reaction.

It has been reported that P-centered radicals can be formed in the presence of Mn(OAc)₃ through homolytic P–H bond cleavage.^[10a] Therefore, we employed Mn(OAc)₃·2H₂O (1 mmol) in the place of AgOAc for the reaction of **1a** with

2a, and **3aa** was formed in 50 % yield (entry 9 in Table 1). Interestingly, the reaction could be conducted even at room temperature (entries 10–12 in Table 1). Comparable results were obtained in DMF and AcOH, whereas the product yield decreased slightly in CH₂Cl₂.

The diarylacetylenes **2b–f** underwent coupling with **1a** when using AgOAc as the oxidant to selectively produce the corresponding 2,3-diaryl-1-phenyl-1*H*-phosphindole-1-ones **3ab–3af** (entries 1–5 in Table 2). 4-Octyne (**2g**) could also be employed in this reaction (entry 6 in Table 2). Interestingly, the asymmetrical phenylacetylenes **2h–p** reacted with **1a** in a regioselective manner to produce the 2-substituted 1,3-diphenyl-1*H*-phosphindole-1-ones **3ah–3ap** in moderate to good yields (entries 7–16 in Table 2). Notably, no other isomers were detected in any of these reactions. One possible reason for the regioselectivity seems to be the facile formation of phenyl-stabilized alkenyl radicals at the addition step (Scheme 2).^[9] In the reactions with silylacetylenes **2j** and **2k**, better results were obtained by using Mn(OAc)₃·2H₂O as the oxidant at room temperature (entries 10 and 11 in Table 2). Under the standard conditions using AgOAc at 100 °C, small amounts (ca. 10 %) of desilylated coupling products were detected by GC–MS (entry 9 in Table 2). As expected, **1a** reacted with 1,4-di(prop-1-yn-1-yl)benzene (**2q**) in a 2:1 manner to form a bis(benzophosphole-3-yl)benzene framework (entry 17 in Table 2).^[2a,12] This type of structure is of interest for application in organic light-emitting diodes and thin-film photovoltaics.^[1c,e,g]

Next, the reactions of variously substituted phenylphosphine oxides with **2a** were examined. The reaction of bis(4-methylphenyl)phosphine oxide (**1b**) in the presence of AgOAc gave both the normal coupling product **3ba** and the unexpected isomer **3ba'** as a 1:1 mixture (entry 1 in Table 3). This may imply that an unanticipated pathway in the annulation process intervenes. The formation of **3ba'** may be rationalized by assuming an attack by alkenyl radical **A** on the *ipso*-carbon atom of one of the 4-methylphenyl rings (Scheme 3a).^[10h,13] The resulting spirocyclohexadienyl radical



Scheme 3. Possible pathways to **3ba** and **3ba'**.

Table 2: Reaction of diphenylphosphine oxide (**1a**) with alkynes **2**.^[a]

Entry	2	Product, yield [%] ^[b]
1		3ab : R = Me, 80
2		3ac : R = OMe, 77
3		3ad : R = NMe ₂ , 35
4		3ae : R = Cl, 89
5		3af : R = Br, 60
6		3ag , 63
7		3ah : R = Me, 86
8		3ai : R = Bu, 73
9		3aj : R = TMS, 0
10 ^[c]		3aj : R = TMS, 54
11 ^[c]		3ak : R = SiMe ₂ Ph, 57
12		3al : R = CO ₂ Et, 57
13		3am : R = Ac, 62
14		3an : R = C(OH)Me ₂ , 61
15		3ao : R = CH ₂ OTBS, 58
16		3ap : R = PO(OEt) ₂ , 41
17 ^[d]		3aq , 38

[a] Reaction conditions: **1a** (0.5 mmol), **2** (0.25 mmol), and AgOAc (1 mmol) in DMF (3 mL) at 100 °C under N₂ for 4 h. TMS = trimethylsilyl, TBS = *tert*-butyldimethylsilyl. [b] Yield of isolated product. [c] Using Mn(OAc)₃·2H₂O (1 mmol) at RT in place of AgOAc. [d] Using **1a** (1 mmol) and AgOAc (2 mmol).

B may then undergo ring expansion to form **C**, thus leading to **3ba'**.

The reaction of 4-methoxy-substituted substrate **1c** also gave a 1:1 mixture of **3ca** and **3ca'** (entry 2 in Table 3). When using Mn(OAc)₃·2H₂O as the oxidant at room temperature, the relative product ratio of **3ca** increased (entry 3 in

Table 3: Reaction of arylphosphine oxides **1** with **2a**.^[a]

Entry	1	Product(s), yield [%] ^[b]
1		3ba + 3ba' : R = Me, 79 (1:1)
2		3ca + 3ca' : R = OMe, 50 (1:1)
3 ^[c]		3ca + 3ca' : R = OMe, 56 (1.9:1)
4		3da + 3da' , 84 (1:2)
5		3ea : R = Bu, 59
6		3fa : R = <i>t</i> Bu, 66
7		3ga : R = OEt, 53
8		3ha , 0
9 ^[c]		3ha , 50

[a] Reaction conditions: **1** (0.5 mmol), **2a** (0.25 mmol), and AgOAc (1 mmol) in DMF (3 mL) at 100 °C under N₂ for 4 h. [b] Yield of isolated product. [c] Using Mn(OAc)₃·2H₂O (1 mmol) at RT in place of AgOAc.

Table 3). As expected, the reaction of bis(3-methylphenyl)-phosphine oxide with **2a** gave a complex mixture of product isomers. By contrast, in the reaction of sterically hindered bis(2-methylphenyl)phosphine oxide (**1d**), **3da'** was formed as the predominant product, along with **3da** (entry 4 in Table 3). It is possible that the 5-*exo-trig* cyclization is retarded by steric repulsion between the methyl group and the aryl moiety on the phosphorus atom (Scheme 3b).^[14]

Alkyl(phenyl)phosphine oxides such as **1e** and **1f** also underwent coupling with **2a** to produce the corresponding *P*-alkylbenzophosphine derivatives **3ea** and **3fa** (entries 5 and 6 in Table 3). As well as phosphine oxides, phenylphosphinate **1g** could also be used in the annulation (entry 7 in Table 3). The reaction of diphenylphosphine sulfide (**1h**) did not proceed at all under the standard conditions (entry 8 in Table 3). This is likely due to the thiophilicity of silver. Fortunately, the reaction of **1h** could be carried out by using

Mn(OAc)₃·2H₂O as the oxidant at room temperature to give **3ha** (entry 9 in Table 3).

Most of the benzophosphole oxides **3** obtained showed solid-state fluorescence as expected. The quantum efficiencies (Φ) of the solid-state fluorescence of **3aa**, **3ac**, and **3ad** were measured as absolute values of 0.54, 0.63, and 0.53, respectively (see the Supporting Information).

In summary, we have demonstrated that the silver- or manganese-mediated dehydrogenative coupling of phenylphosphine oxides with alkynes proceeds efficiently to produce benzophosphole oxide derivatives. Given that a wide range of substrates can be utilized for annulative coupling, this simple protocol may provide a general, one step approach to benzophosphole oxide frameworks of importance in materials chemistry.

Experimental Section

General procedure for the reaction of phenylphosphine oxides **1** with alkynes **2**: A mixture of phenylphosphine oxide (**1**, 0.5 mmol), alkyne (**2**, 0.25 mmol), AgOAc (1.0 mmol, 167 mg), and dibenzyl (ca. 40 mg) as an internal standard was stirred in DMF (3.0 mL) under N₂ at 100 °C for 4 h. GC and GC–MS analyses of the mixtures confirmed consumption of **2**. After cooling, the reaction mixture was diluted with ethyl acetate (40 mL) and insoluble solids were removed on a Celite plug. The filtrate was washed with brine (20 mL × 3). The organic layer was then dried over Na₂SO₄ and concentrated in vacuo. The desired product **3** was isolated by column chromatography on silica gel using dichloromethane/ethyl acetate (1:1 v/v) as eluent. Characterization data of the products are summarized in the Supporting Information.

Received: August 16, 2013

Published online: October 11, 2013

Keywords: annulation · C–C coupling · dehydrogenation · phosphorus heterocycles · radical reactions

- [1] a) A. Fukazawa, E. Yamaguchi, E. Ito, H. Yamada, J. Wang, S. Irle, S. Yamaguchi, *Organometallics* **2011**, *30*, 3870; b) A. Fukazawa, T. Murai, L. Li, Y. Chen, S. Yamaguchi, *C. R. Chim.* **2010**, *13*, 1082; c) H. Tsuji, K. Sato, Y. Sato, E. Nakamura, *Chem. Asian J.* **2010**, *5*, 1294; d) A. Fukazawa, Y. Ichihashi, Y. Kosaka, S. Yamaguchi, *Chem. Asian J.* **2009**, *4*, 1729; e) H. Tsuji, K. Sato, Y. Sato, E. Nakamura, *J. Mater. Chem.* **2009**, *19*, 3364; f) A. Fukazawa, H. Yamada, S. Yamaguchi, *Angew. Chem.* **2008**, *120*, 5664; *Angew. Chem. Int. Ed.* **2008**, *47*, 5582; g) H. Tsuji, K. Sato, L. Ilies, Y. Itoh, Y. Sato, E. Nakamura, *Org. Lett.* **2008**, *10*, 2263; h) A. Fukazawa, M. Hara, T. Okamoto, E. C. Son, C. Xu, K. Tamao, S. Yamaguchi, *Org. Lett.* **2008**, *10*, 913.
- [2] a) T. Sanji, K. Shiraishi, T. Kashiwabara, M. Tanaka, *Org. Lett.* **2008**, *10*, 2689; b) J. G. Cordaro, D. Stein, H. Grützmacher, *J. Am. Chem. Soc.* **2006**, *128*, 14962; c) G. Märkl, G. Y. Jin, K. P. Berr, *Tetrahedron Lett.* **1993**, *34*, 3103; d) T. Butters, W. Winter, *Chem. Ber.* **1984**, *117*, 990; e) T. M. Balthazor, *J. Org. Chem.* **1980**, *45*, 2519; f) W. Winter, *Tetrahedron Lett.* **1975**, *16*, 3913; g) M. D. Rausch, L. P. Klemann, *J. Am. Chem. Soc.* **1967**, *89*, 5732. For relevant synthesis of dibenzophosphole oxides, see: h) Y. Kuninobu, T. Yoshida, K. Takai, *J. Org. Chem.* **2011**, *76*, 7370; i) O. Berger, C. Petit, E. L. Deal, J.-L. Montchamp, *Adv. Synth. Catal.* **2013**, *355*, 1361.
- [3] For earlier examples, see: a) K. Ueura, T. Satoh, M. Miura, *Org. Lett.* **2007**, *9*, 1407; b) L. Li, W. W. Brennessel, W. D. Jones, *J. Am. Chem. Soc.* **2008**, *130*, 12414; c) D. R. Stuart, M. Bertrand-Laperle, K. M. N. Burgess, K. Fagnou, *J. Am. Chem. Soc.* **2008**, *130*, 16474; d) D. A. Colby, R. G. Bergman, J. A. Ellman, *J. Am. Chem. Soc.* **2008**, *130*, 3645; e) S. Rakshit, F. W. Patureau, F. Glorius, *J. Am. Chem. Soc.* **2010**, *132*, 9585; f) T. K. Hyster, T. Rovis, *J. Am. Chem. Soc.* **2010**, *132*, 10565; g) G. Song, D. Chen, C.-L. Pan, R. H. Crabtree, X. Li, *J. Org. Chem.* **2010**, *75*, 7487.
- [4] Selected reviews for transition-metal-catalyzed C–H functionalization: a) D. A. Colby, A. S. Tsai, R. G. Bergman, J. A. Ellman, *Acc. Chem. Res.* **2012**, *45*, 814; b) K. M. Engle, T.-S. Mei, M. Wasa, J.-Q. Yu, *Acc. Chem. Res.* **2012**, *45*, 788; c) E. A. Mitchell, A. Pesciulli, N. Lefevre, L. Meerpoel, B. U. W. Maes, *Chem. Eur. J.* **2012**, *18*, 10092; d) G. Song, F. Wang, X. Li, *Chem. Soc. Rev.* **2012**, *41*, 3651; e) F. W. Patureau, J. Wencel-Delord, F. Glorius, *Aldrichimica Acta* **2012**, *45*, 31; f) S. H. Cho, J. Y. Kim, J. Kwak, S. Chang, *Chem. Soc. Rev.* **2011**, *40*, 5068; g) J. Wencel-Delord, T. Droge, F. Liu, F. Glorius, *Chem. Soc. Rev.* **2011**, *40*, 4740; h) Y. Kuninobu, K. Takai, *Chem. Rev.* **2011**, *111*, 1938; i) C. Liu, H. Zhang, W. Shi, A. Lei, *Chem. Rev.* **2011**, *111*, 1780; j) L. Ackermann, *Chem. Rev.* **2011**, *111*, 1315; k) J. Le Bras, J. Muzart, *Chem. Rev.* **2011**, *111*, 1170; l) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147; m) D. A. Colby, R. G. Bergman, J. A. Ellman, *Chem. Rev.* **2010**, *110*, 624; n) C.-L. Sun, B.-J. Li, Z.-J. Shi, *Chem. Commun.* **2010**, *46*, 677; o) T. Satoh, M. Miura, *Chem. Eur. J.* **2010**, *16*, 11212; p) T. Satoh, M. Miura, *Synthesis* **2010**, 3395; q) B. Karimi, H. Behzadnia, D. Elhamifar, P. F. Akhavan, F. K. Esfahani, A. Zamani, *Synthesis* **2010**, 1399; r) X. Chen, K. M. Engle, D.-H. Wang, J.-Q. Yu, *Angew. Chem.* **2009**, *121*, 5196; *Angew. Chem. Int. Ed.* **2009**, *48*, 5094; s) O. Daugulis, H.-Q. Do, D. Shabashov, *Acc. Chem. Res.* **2009**, *42*, 1074; t) See Ref. [4s]; u) C.-J. Li, *Acc. Chem. Res.* **2009**, *42*, 335; v) F. Kakiuchi, T. Kochi, *Synthesis* **2008**, 3013; w) Y. J. Park, J.-W. Park, C.-H. Jun, *Acc. Chem. Res.* **2008**, *41*, 222; x) I. V. Seregin, V. Gevorgyan, *Chem. Soc. Rev.* **2007**, *36*, 1173; y) E. M. Beccalli, G. Brogini, M. Martinelli, S. Sottocornola, *Chem. Rev.* **2007**, *107*, 5318; z) D. Alberico, M. E. Scott, M. Lautens, *Chem. Rev.* **2007**, *107*, 174; aa) K. Godula, D. Sames, *Science* **2006**, *312*, 67.
- [5] T. Fukutani, N. Umeda, K. Hirano, T. Satoh, M. Miura, *Chem. Commun.* **2009**, 5141.
- [6] For six-membered ring construction by annulation of diarylphosphinic acids under rhodium catalysis, see: Y. Unoh, Y. Hashimoto, D. Takeda, K. Hirano, T. Satoh, M. Miura, *Org. Lett.* **2013**, *15*, 3258.
- [7] Recently, related oxidative coupling reactions involving P–OH directed C–H bond cleavage have been reported: a) J. Seo, Y. Park, I. Jeon, T. Ryu, S. Park, P. H. Lee, *Org. Lett.* **2013**, *15*, 3358; b) X. Meng, S. Kim, *Org. Lett.* **2013**, *15*, 1910; c) L. Y. Chan, S. Kim, T. Ryu, P. H. Lee, *Chem. Commun.* **2013**, *49*, 4682.
- [8] a) Y.-M. Li, M. Sun, H.-L. Wang, Q.-P. Tian, S.-D. Yang, *Angew. Chem.* **2013**, *125*, 4064; *Angew. Chem. Int. Ed.* **2013**, *52*, 3972; b) X. Mao, X. Ma, S. Zhang, H. Hu, C. Zhu, Y. Cheng, *Eur. J. Org. Chem.* **2013**, 4245; c) C.-B. Xiang, Y.-J. Bian, X.-R. Mao, Z.-Z. Huang, *J. Org. Chem.* **2012**, *77*, 7706; d) F. Effenberger, H. Kottmann, *Tetrahedron* **1985**, *41*, 4171. For related Ag-mediated C–H/C–H dehydrogenative coupling, see: e) C. He, S. Guo, J. Ke, J. Hao, H. Xu, H. Chen, A. Lei, *J. Am. Chem. Soc.* **2012**, *134*, 5766.
- [9] U. Wille, *Chem. Rev.* **2013**, *113*, 813.
- [10] a) J. Zhou, G.-L. Zhang, J.-P. Zou, W. Zhang, *Eur. J. Org. Chem.* **2011**, 3412; b) F. Beaufils, F. Dénès, P. Renaud, *Angew. Chem.* **2005**, *117*, 5407; *Angew. Chem. Int. Ed.* **2005**, *44*, 5273; c) A. Sato, H. Yorimitsu, K. Oshima, *Angew. Chem.* **2005**, *117*, 1722; *Angew. Chem. Int. Ed.* **2005**, *44*, 1694; d) A. Kondoh, H. Yorimitsu, K. Oshima, *Chem. Asian J.* **2010**, *5*, 398. For related addition of S- or C-radical/cyclization, see: e) K. Liu, F. Jia, H. Xi, Y. Li, X. Zheng, Q. Guo, B. Shen, Z. Li, *Org. Lett.* **2013**, *15*, 2026; f) D. P. Hari, T. Hering, B. König, *Org. Lett.* **2012**, *14*, 5334; g) M. Tobisu,

- K. Koh, T. Furukawa, N. Chatani, *Angew. Chem.* **2012**, *124*, 11525; *Angew. Chem. Int. Ed.* **2012**, *51*, 11363; h) L. Benati, G. Calestani, R. Leardini, M. Minozzi, D. Nanni, P. Spagnolo, S. Strazzari, G. Zanardi, *J. Org. Chem.* **2003**, *68*, 3454.
- [11] a) X.-Q. Pan, J.-P. Zou, G.-L. Zhang, W. Zhang, *Chem. Commun.* **2010**, *46*, 1721; b) O. Tayama, A. Nakano, T. Iwahama, S. Sakaguchi, Y. Ishii, *J. Org. Chem.* **2004**, *69*, 5494; c) T. Nagayama, A. Nakano, S. Sakaguchi, Y. Ishii, *Org. Lett.* **2006**, *8*, 407. For a review, see: d) B. B. Snider, *Chem. Rev.* **1996**, *96*, 339.
- [12] Compound **3aq** seems to be formed as a mixture of diastereomers. However, the ratio could not be determined owing to overlap in the NMR chemical shifts.
- [13] a) B. S. Bhakuni, A. Kumar, S. J. Balkrishna, J. A. Sheikh, S. Konar, S. Kumar, *Org. Lett.* **2012**, *14*, 2838; b) D. S. Roman, Y. Takahashi, A. B. Charette, *Org. Lett.* **2011**, *13*, 3242.
- [14] It is also possible that the 4-*exo-trig* cyclization is promoted by the methyl group as a result of its stabilization of the corresponding spirocyclohexadienyl radical through hyperconjugation.