


Synthesis and Characterization of a Gold Vinylidene Complex Lacking π -Conjugated Heteroatoms**

Robert J. Harris and Ross A. Widenhoefer*

Abstract: Hydride abstraction from the gold (disilyl)ethynyl complex $[(P)Au\{\eta^1-C\equiv CSi(Me)_2CH_2CH_2Si(Me)_2H\}]$ ($P = P(tBu)_2o$ -biphenyl) with triphenylcarbenium tetrakis(pentafluorophenyl)borate at $-20^\circ C$ formed the cationic gold $(\beta,\beta$ -disilyl)vinylidene complex $[(P)Au=C=CSi(Me)_2CH_2CH_2Si(Me)_2]^+ B(C_6F_5)_4^-$ with $\geq 90\%$ selectivity. ^{29}Si NMR analysis of this complex pointed to delocalization of positive charge onto both the β -silyl groups and the $(P)Au$ fragment. The C1 and C2 carbon atoms of the vinylidene complex underwent facile interconversion ($\Delta G^\ddagger = 9.7$ kcal mol $^{-1}$), presumably via the gold π -disilacyclohexyne intermediate $[(P)Au\{\eta^2-C\equiv CSi(Me)_2CH_2CH_2Si(Me)_2\}]^+ B(C_6F_5)_4^-$.

Cationic gold carbene complexes are widely invoked as intermediates in a range of gold-catalyzed transformations including enyne^[1] and propargyl carboxylate cycloaddition,^[2] alkyne oxidation,^[3] and alkene cyclopropanation.^[4,5] Although the intermediacy of gold carbenes in these transformations is generally accepted, there remains considerable debate regarding the electronic structure of the Au–C bond,^[6–9] due in large part to the dearth of suitable model complexes for structural and/or spectroscopic interrogation. It is therefore significant that within the past year well-defined gold carbene complexes that 1) lack π -conjugated heteroatoms (**A**, **B**),^[7,10] 2) display alkylidene transfer behavior (**C**),^[8] or 3) contain a cyclopropyl group at C1 (**D**)^[9] have been reported (Scheme 1), which provide new insight into the electronic structure and behavior of gold carbene complexes.^[11,12]

Gold vinylidene complexes of the form $[(L)Au=C=CR_2]^+$ constitute a particularly intriguing class of gold carbene complexes that have attracted considerable recent interest. These species have been invoked as intermediates in a number of transformations,^[13] most notably the gold-catalyzed polycyclization of *o*-dialkynyl arenes.^[14] Although the existence of gold vinylidene complexes is supported by both indirect experimental evidence and computation,^[14] direct experimental evidence for these complexes is absent.^[15] The only relevant model complexes are the Fischer-type gold allenylidene complexes **E** ($L = IPr$ or PCy_3 ; $IPr = 1,3$ -bis(2,6-diisopropylphenyl)imidazol-2-ylidene,




Scheme 1. Recent examples of gold carbenoid complexes ($L = N$ -heterocyclic carbene or tertiary phosphine).^[5–9]

Cy = cyclohexyl), which are significantly biased toward the oxonium/iminium contributor **E'** owing to the presence of two π -conjugated heteroatoms (Scheme 1).^[16] Herein we report the synthesis and characterization of a cationic gold vinylidene complex that lacks π -conjugated heteroatoms.

Toward the generation of a cationic gold vinylidene complex lacking π -conjugated heteroatoms, we targeted gold $(\beta,\beta$ -disilyl)vinylidene complexes generated by addition of a pendant silylium ion to the $C\equiv C$ bond of a gold acetylide complex. This approach was inspired by studies from the research groups of Lambert,^[17] Siehl,^[18,19] and Müller,^[19–23] who have both documented and exploited the stabilization of alkyl and vinyl carbenium ions by β -SiC hyperconjugation.^[24] To this end, treatment of disilylethynylacetylene **1** with *n*BuLi followed by auration of the resulting lithium acetylide with $[(P)AuCl]$ ($P = P(tBu)_2o$ -biphenyl) led to isolation of the gold (disilylethynyl)acetylide complex $[(P)Au\{\eta^1-C\equiv CSi(Me)_2CH_2CH_2Si(Me)_2H\}]$ (**2**) in 36% yield (Scheme 2). Treatment of **2** with triphenylcarbenium tetrakis(pentafluorophenyl)borate in CD_2Cl_2 at $-20^\circ C$ for 10 min led to hydride abstraction and cyclization, presumably via the transient silylium ion species **I**, to form the $(\beta,\beta$ -disilyl)vinylidene complex **3** with $\geq 90\%$ selectivity, as determined by ^{31}P NMR analysis.^[25]

Although **3** was stable in solution for short periods (≤ 5 min) at room temperature, concentration or attempted precipitation at $-20^\circ C$ led to rapid decomposition, and **3** was therefore characterized in solution by spectroscopy. The low-temperature ($-80^\circ C$) ^{31}P and ^{29}Si NMR spectra of **3** displayed single resonances at $\delta = 58.3$ and $\delta = 33.5$, respectively, and




Scheme 2. Synthesis of gold $(\beta,\beta$ -disilyl)vinylidene complex **3**.

[*] R. J. Harris, Prof. R. A. Widenhoefer
French Family Science Center, Duke University
Durham, NC 27708-0346 (USA)
E-mail: rwidenho@chem.duke.edu

[**] We acknowledge the NSF (CHE-1213957) for support of this research.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201501474>.



Scheme 3. Resonance contributors to gold vinylidene complex **3** along with vinyl cations **4** and **5**.

the ^1H NMR spectrum of **3** at -80°C displayed a 9:2:6 ratio of resonances at $\delta = 1.31$ (d, $J_{\text{PH}} = 15.5$ Hz), 1.14 (s), and 0.47 (s) corresponding to the *tert*-butyl, Si-CH₂, and Si-CH₃ protons, respectively. Together, these spectroscopic features established both the cyclization of **2** and the high symmetry (static or time-averaged) of gold vinylidene complex **3**. The significant downfield shift of the ^{29}Si NMR resonance of **3** relative to the C(sp)-Si resonance of neutral acetylide precursor **2** ($\delta = -20.6$; $\Delta\delta = +54$) points to accumulation of positive charge on the β -silicon atoms of **3**, as represented by resonance structures **3c** and **3d** (Scheme 3). A number of studies have shown that β -SiC hyperconjugation in β -silyl-substituted carbenium ions increases with the increasing electron demand of the electron-deficient carbon atom.^[26] Müller et al. extended and quantified this relationship by establishing a correlation between the downfield shift of the ^{29}Si NMR chemical shift ($\Delta\delta$) of the α -aryl- β , β -disilylvinyl cations **4** relative to their neutral arylacetylene precursors versus the Hammett σ^+ parameter.^[22] It is therefore worth noting that the $\Delta\delta$ value for **3** is significantly less than that of the analogous α -*tert*-butyl- β , β -disilylvinyl cation **5** ($\Delta\delta = 86$),^[23,27] which suggests that the electron-donor ability of the (P)Au fragment in **3** significantly exceeds that of a *tert*-butyl group, in accord with our previous analysis of Au $\rightarrow$ C electron donation in the cyclopropyl carbene complex **D**.^[9] The IR spectrum of **3** displayed an absorbance at 1955 cm^{-1} assigned to the vinylidene C=C stretch, which is similar to those observed for cations **4**.^[21]

Although the ^{13}C NMR spectrum of **3** at -90°C was likewise consistent with a highly symmetric structure, the key C1 and C2 vinylidene resonances could not be detected, even upon extended acquisition. However, ^{13}C NMR analysis of the doubly labeled isotopomer $[1,2-^{13}\text{C}_2]\text{-3}$ at -90°C displayed broad multiplets at $\delta = 206$ (dd, $J_{\text{CP}} = 110$ Hz, $J_{\text{CC}} = 60$ Hz) and $\delta = 112$ (d, $J_{\text{CC}} = 60$ Hz) corresponding to the vinylidene C1 and C2 atoms, respectively (Scheme 4). As the temperature was raised, these resonances broadened further and disappeared into the baseline between -70 and -50°C . Although these resonances did not coalesce at 25°C , they reappeared unchanged when the solution was again cooled to -90°C . ^{13}C -Spin saturation transfer analysis of $[1,2-^{13}\text{C}_2]\text{-3}$ at -90°C confirmed interconversion of the C1 and C2 vinylidene carbon atoms and established an energy barrier of $\Delta G^\ddagger_{183\text{K}} = 9.7\text{ kcal mol}^{-1}$ for this process. The simplest mechanism that accounts for interconversion of the C1 and C2



Scheme 4. Fluxional behavior of gold vinylidene isotopomer $[1,2-^{13}\text{C}_2]\text{-3}$.

vinylidene carbon atoms of **3** involves 1,2-silyl migration to generate the unobserved gold η^2 -disilacyclohexyne intermediate $[(\text{P})\text{Au}\{\eta^2\text{-C}\equiv\text{CSi}(\text{Me})_2\text{CH}_2\text{CH}_2\text{Si}(\text{Me})_2\}]^+ \text{B}(\text{C}_6\text{F}_5)_4^-$ (**II**; Scheme 4).^[28] Similar fluxional behavior has been documented for β , β -disilylvinyl cations possessing an α -SiR₃ or α -GeR₃ group.^[23]

In summary, we have synthesized the gold (β , β -disilyl)vinylidene complex **3** by hydride abstraction/cyclization of neutral gold acetylide complex **2**. Complex **3** represents the first example of a gold vinylidene complex and a rare example of a gold carbene complex that lacks π -conjugated heteroatoms.^[7,10] The C1 and C2 vinylidene atoms of **3** undergo facile interconversion ($\Delta G^\ddagger_{183\text{K}} = 9.7\text{ kcal mol}^{-1}$), presumably via the gold π -disilacyclohexyne intermediate **II**. ^{29}Si NMR analysis of **3** points to delocalization of positive charge onto both the β -silyl groups and the (P)Au fragment. Ongoing efforts are directed toward further quantifying the extent of Au $\rightarrow$ C electron donation in these complexes.

Keywords: bonding modes · carbene ligands · carbenoids · gold · vinylidenes

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 6867–6869
Angew. Chem. **2015**, *127*, 6971–6973

- [1] a) C. Obradors, A. M. Echavarren, *Acc. Chem. Res.* **2014**, *47*, 902–912; b) A. Fürstner, *Acc. Chem. Res.* **2014**, *47*, 925–938; c) P. Y. Toullec, V. Michelet, *Top. Curr. Chem.* **2011**, *302*, 31–80; d) A. M. Echavarren, E. Jiménez-Núñez, *Top. Catal.* **2010**, *53*, 924–930; e) A. Fürstner, *Chem. Soc. Rev.* **2009**, *38*, 3208–3221; f) E. Jiménez-Núñez, A. M. Echavarren, *Chem. Rev.* **2008**, *108*, 3326–3350; g) V. Michelet, P. Y. Toullec, J. P. Genêt, *Angew. Chem. Int. Ed.* **2008**, *47*, 4268–4315; *Angew. Chem.* **2008**, *120*, 4338–4386.
- [2] A. Correa, N. Marion, L. Fensterbank, M. Malacria, S. P. Nolan, L. Cavallo, *Angew. Chem. Int. Ed.* **2008**, *47*, 718–721; *Angew. Chem.* **2008**, *120*, 730–733.
- [3] D. Qian, J. Zhang, *Chem. Soc. Rev.* **2015**, *44*, 677–698.
- [4] a) L. Zhang, *Acc. Chem. Res.* **2014**, *47*, 877–888; b) H.-S. Yeom, S. Shin, *Acc. Chem. Res.* **2014**, *47*, 966–977.
- [5] For general reviews of gold catalysis, see a) M. Bandini, *Chem. Soc. Rev.* **2011**, *40*, 1358–1367; b) A. Corma, A. Leyva-Pérez, M. J. Sabater, *Chem. Rev.* **2011**, *111*, 1657–1712; c) M. Rudolph, A. S. K. Hashmi, *Chem. Commun.* **2011**, *47*, 6536–6544; d) N. Krause, C. Winter, *Chem. Rev.* **2011**, *111*, 1994–2009; e) T. C. Boorman, I. Larrosa, *Chem. Soc. Rev.* **2011**, *40*, 1910–1925; f) A. Pradal, P. Y. Toullec, V. Michelet, *Synthesis* **2011**, 1501–1514.
- [6] a) A. S. K. Hashmi, *Angew. Chem. Int. Ed.* **2008**, *47*, 6754–6756; *Angew. Chem.* **2008**, *120*, 6856–6858; b) A. Fürstner, L. Morency, *Angew. Chem. Int. Ed.* **2008**, *47*, 5030–5033; *Angew. Chem.* **2008**, *120*, 5108–5111; c) A. M. Echavarren, *Nat. Chem.* **2009**, *1*, 431–433; d) G. Seidel, R. Mynott, A. Fürstner, *Angew. Chem. Int. Ed.* **2009**, *48*, 2510–2513; *Angew. Chem.* **2009**, *121*, 2548–2551; e) D. Benitez, N. D. Shapiro, E. Tkatchouk, Y. Wang, W. A. Goddard, F. D. Toste, *Nat. Chem.* **2009**, *1*, 482–486;

- f) G. Seidel, B. Gabor, R. Goddard, B. Heggen, W. Thiel, A. Fürstner, *Angew. Chem. Int. Ed.* **2014**, *53*, 879–882; *Angew. Chem.* **2014**, *126*, 898–901; g) G. Ciancaleoni, L. Biasioli, G. Bistoni, A. Macchioni, F. Tarantelli, D. Zuccaccia, L. Belpassi, *Chem. Eur. J.* **2015**, *21*, 2467–2473.
- [7] M. W. Hussong, F. Rominger, P. Krämer, B. F. Straub, *Angew. Chem. Int. Ed.* **2014**, *53*, 9372–9375; *Angew. Chem.* **2014**, *126*, 9526–9529.
- [8] G. Seidel, A. Fürstner, *Angew. Chem. Int. Ed.* **2014**, *53*, 4807–4811; *Angew. Chem.* **2014**, *126*, 4907–4911.
- [9] R. E. M. Brooner, R. A. Widenhoefer, *Chem. Commun.* **2014**, *50*, 2420–2423.
- [10] R. J. Harris, R. A. Widenhoefer, *Angew. Chem. Int. Ed.* **2014**, *53*, 9369–9371; *Angew. Chem.* **2014**, *126*, 9523–9525.
- [11] For gas-phase studies of gold carbene complexes, see a) A. Fedorov, M.-E. Moret, P. Chen, *J. Am. Chem. Soc.* **2008**, *130*, 8880–8881; b) A. Fedorov, P. Chen, *Organometallics* **2009**, *28*, 1278–1281; c) L. Batiste, A. Fedorov, P. Chen, *Chem. Commun.* **2010**, *46*, 3899–3901; d) A. Fedorov, L. Batiste, A. Bach, D. M. Birney, P. Chen, *J. Am. Chem. Soc.* **2011**, *133*, 12162–12171; e) D. H. Ringger, P. Chen, *Angew. Chem. Int. Ed.* **2013**, *52*, 4686–4689; *Angew. Chem.* **2013**, *125*, 4784–4787; f) D. H. Ringger, I. J. Kobylanskii, D. Serra, P. Chen, *Chem. Eur. J.* **2014**, *20*, 14270–14281; g) L. Batiste, P. Chen, *J. Am. Chem. Soc.* **2014**, *136*, 9296–9307.
- [12] For additional examples of heteroatom-stabilized gold carbenes, see a) L.-P. Liu, B. Xu, M. S. Mashuta, G. B. Hammond, *J. Am. Chem. Soc.* **2008**, *130*, 17642–17643; b) M. N. Fañanás-Mastral, F. Aznar, *Organometallics* **2009**, *28*, 666–668; c) U. Schubert, K. Ackermann, R. Aumann, *Cryst. Struct. Commun.* **1982**, *11*, 591–594; d) G. Ung, G. Bertrand, *Angew. Chem. Int. Ed.* **2013**, *52*, 11388–11391; *Angew. Chem.* **2013**, *125*, 11599–11602.
- [13] a) Y. Xia, A. S. Dudnik, Y. Li, V. Gevorgyan, *Org. Lett.* **2010**, *12*, 5538–5541; b) V. Lavallio, G. D. Frey, S. Kousar, B. Donnadiou, G. Bertrand, *Proc. Natl. Acad. Sci. USA* **2007**, *104*, 13569–13573; c) H. Rabaâ, B. Engels, T. Hupp, A. S. K. Hashmi, *Int. J. Quantum Chem.* **2007**, *107*, 359–365; d) V. Mamane, P. Hannen, A. Fürstner, *Chem. Eur. J.* **2004**, *10*, 4556–4575; e) E. Soriano, J. Marco-Contelles, *Organometallics* **2006**, *25*, 4542–4553; f) I. V. Seregin, V. Gevorgyan, *J. Am. Chem. Soc.* **2006**, *128*, 12050–12051.
- [14] a) A. Gómez-Suárez, S. P. Nolan, *Angew. Chem. Int. Ed.* **2012**, *51*, 8156–8159; *Angew. Chem.* **2012**, *124*, 8278–8281; b) L. Ye, Y. Wang, D. H. Aue, L. Zhang, *J. Am. Chem. Soc.* **2012**, *134*, 31–34; c) A. S. K. Hashmi, M. Wietek, I. Braun, M. Rudolph, F. Rominger, *Angew. Chem. Int. Ed.* **2012**, *51*, 10633–10637; *Angew. Chem.* **2012**, *124*, 10785–10789; d) A. S. K. Hashmi, M. Wietek, I. Braun, P. Nösel, L. Jongbloed, M. Rudolph, F. Rominger, *Adv. Synth. Catal.* **2012**, *354*, 555–562; e) M. M. Hansmann, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Angew. Chem. Int. Ed.* **2013**, *52*, 2593–2598; *Angew. Chem.* **2013**, *125*, 2653–2659; f) I. Braun, A. M. Asiri, A. S. K. Hashmi, *ACS Catal.* **2013**, *3*, 1902–1907; g) P. Nösel, T. Lauterbach, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Chem. Eur. J.* **2013**, *19*, 8634–8641; h) M. M. Hansmann, S. Tsupova, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Chem. Eur. J.* **2014**, *20*, 2215–2223; i) A. S. K. Hashmi, *Acc. Chem. Res.* **2014**, *47*, 864–876; j) J. Bucher, T. Wurm, K. S. Nalivela, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Angew. Chem. Int. Ed.* **2014**, *53*, 3854–3858; *Angew. Chem.* **2014**, *126*, 3934–3939; k) H. M. Vilhelmsen, A. S. K. Hashmi, *Chem. Eur. J.* **2014**, *20*, 1901–1908; l) J. Xie, C. Pan, A. Abdukadara, C. Zhu, *Chem. Soc. Rev.* **2014**, *43*, 5245–5256; m) O. J. S. Pickup, I. Khazal, E. J. Smith, A. C. Whitwood, J. M. Lynam, K. Bolaky, T. C. King, B. W. Rawe, N. Fey, *Organometallics* **2014**, *33*, 1751–1761; n) Y. Tokimizu, M. Wietek, M. Rudolph, S. Oishi, N. Fujii, A. S. K. Hashmi, H. Ohno, *Org. Lett.* **2015**, *17*, 604–607; o) P. Nösel, V. Müller, S. Mader, S. Moghimi, M. Rudolph, I. Braun, F. Rominger, A. S. K. Hashmi, *Adv. Synth. Catal.* **2015**, *357*, 500–506; p) J. Bucher, T. Stöber, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Angew. Chem. Int. Ed.* **2015**, *54*, 1666–1670; *Angew. Chem.* **2015**, *127*, 1686–1690; q) P. Morán-Poladura, E. Rubio, J. M. González, *Angew. Chem. Int. Ed.* **2015**, *54*, 3052–3055; *Angew. Chem.* **2015**, *127*, 3095–3098.
- [15] For recent reviews of transition-metal vinylidene complexes, see a) J. M. Lynam, *Chem. Eur. J.* **2010**, *16*, 8238–8247; b) B. M. Trost, A. McClory, *Chem. Asian J.* **2008**, *3*, 164–194.
- [16] a) M. M. Hansmann, F. Rominger, A. S. K. Hashmi, *Chem. Sci.* **2013**, *4*, 1552–1559; b) X.-S. Xiao, W.-L. Kwong, X. Guan, C. Yang, W. Lu, C.-M. Che, *Chem. Eur. J.* **2013**, *19*, 9457–9462.
- [17] a) J. B. Lambert, Y. Zhao, H. Wu, *J. Org. Chem.* **1999**, *64*, 2729–2736; b) J. B. Lambert, Y. Zhao, *J. Am. Chem. Soc.* **1996**, *118*, 7867–7868.
- [18] a) H.-U. Siehl, F.-P. Kaufmann, Y. Apeloig, V. Braude, D. Danovich, A. Berndt, N. Stamatis, *Angew. Chem. Int. Ed. Engl.* **1991**, *30*, 1479–1482; *Angew. Chem.* **1991**, *103*, 1546–1549; b) H.-U. Siehl, F.-P. Kaufmann, *J. Am. Chem. Soc.* **1992**, *114*, 4937–4939; c) H.-U. Siehl, F.-P. Kaufmann, K. Hori, *J. Am. Chem. Soc.* **1992**, *114*, 9343–9349; d) H.-U. Siehl, *Pure Appl. Chem.* **1995**, *67*, 769–775; e) H.-U. Siehl, S. Brixner, C. Coletti, N. Reb, B. Chiavarinoc, M. E. Crestonic, A. De Petris, S. Fornarini, *Int. J. Mass Spectrom.* **2013**, *334*, 58–66.
- [19] T. Müller, R. Meyer, D. Lennartz, H.-U. Siehl, *Angew. Chem. Int. Ed.* **2000**, *39*, 3074–3077; *Angew. Chem.* **2000**, *112*, 3203–3206.
- [20] a) H.-U. Steinberger, T. Müller, N. Auner, C. Maerker, P. von R. Schleyer, *Angew. Chem. Int. Ed. Engl.* **1997**, *36*, 626–628; *Angew. Chem.* **1997**, *109*, 667–669; b) A. Klaer, W. Saak, D. Haase, T. Müller, *J. Am. Chem. Soc.* **2008**, *130*, 14956–14957; c) T. Müller, C. Bauch, M. Ostermeier, M. Bolte, N. Auner, *J. Am. Chem. Soc.* **2003**, *125*, 2158–2168; d) T. Müller, M. Juhász, C. A. Reed, *Angew. Chem. Int. Ed.* **2004**, *43*, 1543–1546; *Angew. Chem.* **2004**, *116*, 1569–1572.
- [21] A. Klaer, T. Müller, *J. Phys. Org. Chem.* **2010**, *23*, 1043–1048.
- [22] T. Müller, D. Margraf, Y. Syha, *J. Am. Chem. Soc.* **2005**, *127*, 10852–10860.
- [23] A. Klaer, Y. Syha, H. R. Nasiri, T. Müller, *Chem. Eur. J.* **2009**, *15*, 8414–8423.
- [24] J. B. Lambert, *Tetrahedron* **1990**, *46*, 2677–2689.
- [25] The length of the alkyl chain connecting the two silicon atoms was critical to successful formation of a stable gold vinylidene complex. For example, hydride abstraction from the (disilylmethyl)acetylide complex $[(P)Au\{\eta^1-C\equiv Si(Me)_2CH_2SiMe_2H\}]$ formed no detectable vinylidene complex, whereas preliminary investigation of hydride abstraction from the corresponding (disilylpropyl)acetylide complex $[(P)Au\{\eta^1-C\equiv Si(Me)_2(CH_2)_3SiMe_2H\}]$ suggested formation of a ground-state π -disilacycloheptyne complex.^[28]
- [26] a) W. Zhang, J. A. Stone, M. A. Brook, G. A. McGibbon, *J. Am. Chem. Soc.* **1996**, *118*, 5764–5771; b) X. Li, J. A. Stone, *J. Am. Chem. Soc.* **1989**, *111*, 5586–5592; c) H.-U. Siehl in *Dicoordinated Carbocations* (Eds.: Z. Rappoport, P. J. Stang), Wiley, New York, **1997**, p. 189.
- [27] Unfortunately, no correlation between the $\Delta\delta$ and σ^+ values has been established for α -aryl- β -disilylvinyl cations in which the β -silicon atoms are part of a five-membered ring, such as in **3** and **5**. As such, no definitive assignment of the electron-donor ability of the (P)Au fragment of **3** can be determined from these data.
- [28] For an example of a neutral gold π -cycloheptyne complex and a cationic π -bis(*tert*-butyldimethylsilyl)acetylene complex, see a) P. Schulte, U. Behrens, *Chem. Commun.* **1998**, 1633–1634; b) D. Zhu, X. Cao, B. Yu, *Org. Chem. Front.* **2015**, *2*, 369–365.

Received: February 14, 2015

Published online: April 23, 2015