

A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie


International Edition



www.angewandte.org

2008–47/15

Metal–Oxygen Isopolyhedra Fullerenes



Metal–oxygen cage clusters ...

... with fullerene topologies can be assembled from uranyl hexagonal bipyramids under certain conditions, as described by P. C. Burns and co-workers in their Communication on page 2824 ff. A cluster containing 50 polyhedra has the same topology as the fullerene $C_{50}Cl_{10}$. Synthesis of a cluster containing 40 uranyl hexagonal bipyramids demonstrates that the inclusion of topological squares in addition to pentagons and hexagons gives access to topological isomers that are not possible for carbon fullerenes.

 WILEY-VCH

Inside Cover

Tori Z. Forbes, J. Gregory McAlpin, Rachel Murphy, and Peter C. Burns*

Metal–oxygen cage clusters with fullerene topologies can be assembled from uranyl hexagonal bipyramids under certain conditions, as described by P. C. Burns and co-workers in their Communication on page 2824 ff. A cluster containing 50 polyhedra has the same topology as the fullerene $C_{50}Cl_{10}$. Synthesis of a cluster containing 40 uranyl hexagonal bipyramids demonstrates that the inclusion of topological squares in addition to pentagons and hexagons gives access to topological isomers that are not possible for carbon fullerenes.

