

A Journal of the Gesellschaft Deutscher Chemiker


# Angewandte Chemie

International Edition



www.angewandte.org

2009–48/18



## The bisoxazoline ligand ...

... enhances the  $\pi$  philicity of palladium(II). The direct conversion of a terminal alkyne to  $\beta$ -methoxyacrylate with the aid of the bisoxazoline (box) ligand is reported by K. Kato and co-workers in their Communication on page 3326 ff. Density functional calculations on the complex [(box)Pd(TFA)<sub>2</sub>] reveal that ligand  $\pi$  orbitals delocalized over the double bonds and the periphery are energetically close to the LUMO. The reaction is also applied to a one-pot synthesis of Kawain.

 WILEY-VCH

## Inside Cover

**Keisuke Kato,\* Satoshi Motodate, Tomoyuki Mochida,  
Takuya Kobayashi, and Hiroyuki Akita\***

**The bisoxazoline ligand** enhances the  $\pi$  philicity of palladium(II). The direct conversion of a terminal alkyne to  $\beta$ -methoxyacrylate with the aid of the bisoxazoline (box) ligand is reported by K. Kato and co-workers in their Communication on page 3326 ff. Density functional calculations on the complex  $[(\text{box})\text{Pd}(\text{TFA})_2]$  reveal that ligand  $\pi$  orbitals delocalized over the double bonds and the periphery are energetically close to the LUMO. The reaction is also applied to a one-pot synthesis of Kawain.

