

A Journal of the Gesellschaft Deutscher Chemiker


Angewandte Chemie

International Edition



www.angewandte.org

2009–48/26



Bidirectional chemo-switching ...

... of magnetism occurs in a microporous coordination polymer containing spin-cross-over subunits, as described by M. Ohba, J. A. Real, S. Kitagawa, and co-workers in their Communication on page 4767 ff. In situ magnetic measurements reveal that most guest molecules transform the framework spin state from diamagnetic low spin (red) to paramagnetic high spin (yellow), whereas the guest CS_2 stabilizes the low-spin state. These induced spin states are retained as a memory effect after the release of the guest.

 WILEY-VCH

Inside Cover

Srijanani Bhaskar, Jonathon Hitt, Sei-Won Laura Chang, and Joerg Lahann*

Bidirectional chemo-switching of magnetism occurs in a microporous coordination polymer containing spin-crossover subunits, as described by M. Ohba, J. A. Real, S. Kitagawa, and co-workers in their Communication on page 4767 ff. In situ magnetic measurements reveal that most guest molecules transform the framework spin state from diamagnetic low spin (red) to paramagnetic high spin (yellow), whereas the guest CS₂ stabilizes the low-spin state. These induced spin states are retained as a memory effect after the release of the guest.

