

Bis-Terminal Hydroxy Polyethers as All-Purpose, Multifunctional Organic Promoters: A Mechanistic Investigation and Applications**

Ji Woong Lee, Hailong Yan, Hyeong Bin Jang, Hong Ki Kim, Sung-Woo Park, Sungyul Lee,*
Dae Yoon Chi,* and Choong Eui Song*

Catalysis is one of the central themes in chemical science. The promotion of physicochemical and biological processes, as demonstrated by enzymes in living cells, is an extremely important subject to be elucidated in all branches of science. Therefore, the design and application of new and efficient types of promoters for organic transformations are of fundamental importance.

A range of important classes of organic reactions can be conducted with fluoride anions as a catalyst or nucleophilic reagent.^[1] Although alkali metal fluorides are a readily available source of the fluoride ion, their applications are limited because of their low solubility in organic solvents. To increase the solubility of alkali metal salts, crown ethers are frequently employed to generate a “naked” fluoride ion (Figure 1 I).^[2] This simple but innovative concept of crown ethers helped lay the foundation for the important fields of ion and molecular recognition and has therefore had a profound impact on science.^[3] However, their application to chemical reactions is often hampered by the strong basicity of fluoride, which causes it to induce the formation of by-products. During the course of our efforts to develop general catalyst (promoter) systems employing alkali metal salts as nucleophilic sources, it was observed that bulky protic solvents (e.g. *t*BuOH) can be suitable to generate a “flexible” fluoride ion from CsF through controlled hydrogen bonding to reduce the basicity of the nucleophile, fluoride (Figure 1 II).^[4] However, owing to




Figure 1. Bis-terminal hydroxy polyethers (III) as multifunctional promoters.

the strong Coulombic influence of K^+ on F^- , KF was observed to be inactive toward S_N2 reactions. Thus, the combination of these two concepts—“naked” and flexible systems together in one molecule (Figure 1 I and II)—became the focus of our study to design a much more versatile promoter system for a wide range of organic transformations. Herein, we describe the phenomenal efficiency of bis-terminal hydroxy polyethers as a new type of all-purpose promoter system (Figure 1 III), which act as extremely efficient catalytic reaction media for nucleophilic fluorination with alkali metal fluorides. Moreover, by using chiral bis-terminal hydroxy polyethers as catalysts, the catalytic desilylative kinetic resolution of the silyl ethers of racemic secondary alcohols with KF was successfully achieved for the first time.

Cooperative catalysis,^[5] the simultaneous binding and activation of reacting partners resulting in both preorganization of the substrates and stabilization of the transition state structures, is a fundamental principle in modern catalysis. Recently, much efforts in this field have focused on the development of efficient multifunctional organocatalysts.^[6] We anticipated that bis-terminal hydroxy polyethers might serve as a new type of multifunctional organic promoters in various organic reactions through a cooperative activation mechanism as depicted in Figure 1 III: The ether groups act as a Lewis base toward K^+ , “freeing” the nucleophile, as well as “enhancing” the solubility of the potassium salts. On the other hand, one of the two OH groups forms controlled hydrogen bonding with the fluoride anion, thus decreasing the basicity of the nucleophile, whereas the other OH group may be able to simultaneously activate the electrophile by hydrogen bonding, thereby stabilizing the transition state (Figure 1 III). More importantly, this type of promoter can easily be modified to produce “chiral variants”.

To prove the validity of this novel concept, we first examined bimolecular nucleophilic fluorination, which is considered to be one of the most difficult S_N2 reactions. We

[*] S.-W. Park, Prof. S. Lee
Department of Applied Chemistry, Kyunghee University
Yongin, Gyeonggi, 446-701 (Korea)
E-mail: syllee@khu.ac.kr

Prof. D. Y. Chi
Department of Chemistry, Sogang University
1 Shinsudong, Mapogu, Seoul 121-742 (Korea)
E-mail: dychi@sogang.ac.kr

J. W. Lee, H. Yan, H. B. Jang, H. K. Kim, Prof. C. E. Song
Department of Chemistry, Sungkyunkwan University
300 Cheoncheon, Jangsan, Suwon, Gyeonggi, 440-746 (Korea)
E-mail: sl673@skku.edu


Prof. C. E. Song
Department of Energy Science, Sungkyunkwan University
300 Cheoncheon, Jangsan, Suwon, Gyeonggi, 440-746 (Korea)

[**] This work was supported by the MOEHRD (grant no. KRF-2008-005J00701), the SRC program of MEST/KOSEF (grant no. R11-2005-008-00000-0), the KOSEF basic research program (grant no. 2009-0070597), and the WCU program (grant no. R31-2008-000-10029-0).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.200903903>.

conducted the nucleophilic fluorination on the model substrate 2-(3-methanesulfonyloxypropoxy)naphthalene (**1**) with 5 equivalents of potassium fluoride in triethylene glycol or tetraethylene glycol as the solvent at 100 °C. The results are summarized in Table 1, together with those obtained in other solvent systems reported previously by us^[4,7] and other


Table 1: Nucleophilic fluorination with alkali metal fluoride in various solvent systems.^[a]



Entry	Solvent	Nucleophile (MF)	t	Yield [%] ^[b]
1 ^[c]	CH ₃ CN	KF	24 h	0
2 ^[c]	[18]crown-6 in CH ₃ CN	KF	24 h	40
3 ^[d]	[18]crown-6 in DMF	KF	1 h	4
4	<i>t</i> BuOH	KF	1.5 h	0
5	<i>tert</i> -amyl alcohol	KF	1.5 h	0
6	triethylene glycol	KF	1.5 h	93
7	tetraethylene glycol	KF	1.5 h	92
8 ^[e]	triethylene glycol	KF	1 min	95
9 ^[e]	tetraethylene glycol	KF	1 min	94
10	PEG 600	KF	1.5 h	70 ^[f]
11	PEG 600	CsF	1.5 h	> 98 ^[f]
12	PEG 1000	KF	1.5 h	39 ^[f]
13	PEG 1000	CsF	1.5 h	> 98 ^[f]

[a] Unless otherwise indicated, all reactions were carried out with **1** (0.1 mmol) and KF (5 equiv) at 100 °C in 0.5 mL of solvent. [b] Yield of isolated product. [c] See Ref. [7]. [d] See Ref. [8]. [e] Reactions were carried out under microwave irradiation (700 W). [f] Yields were determined by ¹H NMR spectroscopy. DMF = *N,N*-dimethylformamide.


group.^[8] Pleasingly, when using tri- or tetraethylene glycol as solvent, the substitution reactions proceeded at an unprecedented fast rate and were completed within 1.5 h to afford the fluorinated product **2** in excellent yields (92–93%) without forming any appreciable amount of by-products (Table 1, entries 6 and 7). Moreover, by applying microwave irradiation (700 W) to the same reaction, the fluorinated product **2** could be obtained in nearly quantitative yields within only 1 minute (Table 1, entries 8 and 9). In contrast to these results, the same fluorination reaction with KF in aprotic polar solvents such as acetonitrile in the presence or absence of [18]crown-6 proceeded sluggishly (Table 1, entries 1–3).^[7,8] Bulky protic solvents such as *t*BuOH and *tert*-amyl alcohol, which are known to be excellent media for fluorination with CsF,^[4] also showed no reactivity with KF (Table 1, entries 4 and 5). Interestingly, PEG 600 or PEG 1000 (containing ca. 13 and 22 ether units, respectively) showed low reactivity for the same reaction (Table 1, entries 10 and 12). Meanwhile the reaction of **1** with CsF in PEG 600 or PEG 1000 proceeded smoothly to give the desired fluorinated product **2** in nearly quantitative yields (Table 1, entries 11 and 13). This result strongly indicates that the effectiveness of alkali metal cation is determined by the chain length of the bis-terminal hydroxy polyethers, as would be expected. Moreover, as shown in Scheme 1, the fluorination of 2-(2-mesyloxyethyl)naphthalene (**3**) proceeded to give predominantly 2-(2-fluoroethyl)naphthalene (**4a**) in 95%



Scheme 1. Nucleophilic fluorination of **3** with KF

yield, along with the alkene **4b** in 5% yield. It is well known that introducing fluorine into haloethyl or alkanesulfonyloxyethyl aromatic compounds is very difficult because of their tendency to undergo elimination reactions to produce styrenes.^[9] All of the aforementioned results are certainly consistent with our hypothesis of a cooperative mechanism depicted in Figure 1 III.

To further verify the importance of the cooperative multifunctionality of the promoters, the fluorination of the mesylate **1** with KF (5 equiv) was carried out in various glycol derivatives as the solvent. As shown in Scheme 2, when triethylene glycol dimethyl ether (in which both terminal




Scheme 2. Nucleophilic fluorination with KF in triethylene glycol derivatives.

hydroxy groups are methylated) was used as the solvent, the reaction did not proceed at all. On the other hand, triethylene glycol monoethyl ether showed about half of the reactivity of triethylene glycol under the same reaction conditions. We also carried out the same reaction in a mixed solvent system consisting of triethylene glycol dimethyl ether and *t*BuOH (1:1, v/v) on the assumption that the ether units of the former can chelate the potassium cation and the latter can stabilize the fluoride anion by controlled hydrogen bonding. However, no conversion was observed. All of these experimental results strongly suggest that, in the case of bis-terminal hydroxy polyethers, the “intramolecular” cooperative activation mechanism is responsible for the rate acceleration in nucleophilic fluorinations.

Support for the proposed cooperative mechanism was obtained from quantum chemical calculations carried out by employing the accurate DFT (MPW1K/6-31++G**) method.^[10] Figure 2 depicts the reactions of C₃H₇OMs with K⁺F[−] in tetraethylene glycol. In Figure 2, five oxygen atoms in the tetraethylene glycol molecule act as a Lewis base toward K⁺, thus drastically lowering its electrostatic effects and thereby “freeing” the nucleophile, F[−]. The ion pair K⁺F[−] resides in the hollow “pocket” formed by the five oxygen atoms in the solvent (very much like an active site in an

the corresponding alcohols in nearly quantitative yields. Notably, the base-sensitive silyl ether (*R*)-**8** could also be selectively deprotected without racemization (Table 2, entry 4). It is well known that tetra-*n*-butylammonium fluoride (TBAF), the most common reagent for the cleavage of silyl ethers, is inappropriate for base sensitive substrates because of the strong basicity of the fluoride anion.^[1a,16]

Finally, we are also in the process of developing asymmetric variants of the catalytic process using our multifunctional promoter. For example, treating the racemic silyl ether **9** with 0.5 equivalents of KF at 15 °C for 24 hours in the presence of the chiral variant (*R*)-**11** (10 mol%) of tetraethylene glycol as a catalyst furnished the resolved alcohol (*S*)-**10** with 73 % *ee* (Scheme 4 and the Supporting Informa-



Scheme 4. Desilylative kinetic resolution of *rac*-**9** with KF. TMS = trimethylsilyl.

tion). To our knowledge, this is the first catalytic example of the desilylative kinetic resolution of the silyl ethers of racemic secondary alcohols.^[17] This process can certainly open up a new avenue for asymmetric reactions catalyzed by chiral fluoride,^[1b,c] which have vast synthetic potential but remain a relatively undeveloped field.

In conclusion, we have demonstrated that multifunctional bis-terminal hydroxy polyether derivatives can strongly promote a range of chemical reactions, in which alkali metal salts are used as nucleophilic sources. Achiral bis-terminal hydroxy polyether derivatives dramatically facilitated the most difficult process of nucleophilic fluorination, even with KF. We also described the first successful desilylative kinetic resolution of the silyl ethers of racemic secondary alcohols with KF by using the chiral variants of bis-terminal hydroxy polyether derivatives as catalysts. Quantum chemical calculations provide detailed insight into the modes of action of this type of organic promoter. On the basis of all these remarkable results, we believe that our new promoter system (bis-terminal hydroxy polyethers and their chiral variants) is a new development that constitutes a significant step forward in many important areas of science, e.g., solvent engineering, enantioselective organocatalysis, and diagnostic medicine. Our current work in the laboratory focuses on 1) expanding the scope of the applicability of bis-terminal hydroxy

polyethers as multifunctional promoters for other organic reactions, 2) developing efficient asymmetric variants, and 3) PET applications.

Received: July 16, 2009

Published online: September 8, 2009

Keywords: cooperative catalysis · density functional calculations · kinetic resolution · nucleophilic fluorination · polyethers

- [1] a) J. H. Clark, *Chem. Rev.* **1980**, *80*, 429–452; b) S. Shirakawa, T. Ooi, K. Maruoka in *Asymmetric Phase Transfer Catalysis* (Ed.: K. Maruoka), Wiley-VCH, Weinheim **2008**, Chapter 9, pp. 189–206; c) T. Ooi, K. Maruoka, *Acc. Chem. Res.* **2004**, *37*, 526–533.
- [2] C. L. Liotta, H. P. Harris, *J. Am. Chem. Soc.* **1974**, *96*, 2250–2252.
- [3] C. J. Pedersen, *Nobel Lecture*, http://nobelprize.org/nobel_prizes/chemistry/laureates/1987/pedersen-lecture.html.
- [4] D. W. Kim, D.-S. Ahn, Y.-H. Oh, S. Lee, H. S. Kil, S. J. Oh, S. J. Lee, J. S. Kim, J. S. Ryu, D. H. Moon, D. Y. Chi, *J. Am. Chem. Soc.* **2006**, *128*, 16394–16397.
- [5] a) J.-A. Ma, D. Cahard, *Angew. Chem.* **2004**, *116*, 4666–4683; *Angew. Chem. Int. Ed.* **2004**, *43*, 4566–4583; b) C. Allemann, R. Gordillo, F. R. Clemente, P. H.-Y. Cheong, K. N. Houk, *Acc. Chem. Res.* **2004**, *37*, 558–569.
- [6] a) B. List, *Chem. Rev.* **2007**, *107*, 5413–5415; b) B. List, J. W. Yang, *Science* **2006**, *313*, 1584–1586; c) D. W. C. MacMillan, *Nature* **2008**, *455*, 304–308.
- [7] D. W. Kim, C. E. Song, D. Y. Chi, *J. Am. Chem. Soc.* **2002**, *124*, 10278–10279.
- [8] D. W. Kim, H.-J. Jeong, S. T. Lim, M.-H. Sohn, *Angew. Chem.* **2008**, *120*, 8532–8534; *Angew. Chem. Int. Ed.* **2008**, *47*, 8404–8406.
- [9] J. M. Gerdes, R. N. Keil, A. T. Shulgin, C. A. Mathis, *J. Fluorine Chem.* **1996**, *78*, 121–129.
- [10] a) C. Adamo, V. Barone, *J. Chem. Phys.* **1998**, *108*, 664–675; b) B. J. Lynch, P. L. Fast, M. Harris, D. G. Truhlar, *J. Phys. Chem. A* **2000**, *104*, 4811–4815.
- [11] Y.-H. Oh, D.-S. Ahn, S.-Y. Chung, J.-H. Jeon, S.-W. Park, S. J. Oh, D. W. Kim, H. S. Kil, D. Y. Chi, S. Lee, *J. Phys. Chem. A* **2007**, *111*, 10152–10161.
- [12] S.-S. Lee, H.-S. Kim, T.-K. Hwang, Y.-H. Oh, S.-W. Park, S. Lee, B. S. Lee, D. Y. Chi, *Org. Lett.* **2008**, *10*, 61–64.
- [13] S. Im, S.-W. Jang, H.-R. Kim, Y.-H. Oh, S.-W. Park, S. Lee, D. Y. Chi, *J. Phys. Chem. A* **2009**, *113*, 3685–3689.
- [14] a) D. L. Zechel, S. P. Reid, O. Nashiru, C. Mayer, D. Stoll, D. L. Jakeman, R. A. J. Warren, S. G. Withers, *J. Am. Chem. Soc.* **2001**, *123*, 4350–4351; b) D. O'Hagan, C. Schaffrath, S. L. Cobb, J. T. G. Hamilton, C. D. Murphy, *Nature* **2002**, *416*, 279–279; c) C. Dong, F. Huang, H. Deng, C. Schaffrath, J. B. Spencer, D. O'Hagan, J. H. Naismith, *Nature* **2004**, *427*, 561–565.
- [15] a) M. E. Phelps, *Proc. Natl. Acad. Sci. USA* **2000**, *97*, 9226–9233; b) V. Marx, *Chem. Eng. News* **2005**, *83*, 25–34.
- [16] E. J. Corey, B. B. Snider, *J. Am. Chem. Soc.* **1972**, *94*, 2549–2550.
- [17] Three examples for catalytic enantioselective silylation of alcohols have been reported; a) Y. Zhao, J. Rodrigo, A. H. Hoveyda, M. L. Snapper, *Nature* **2006**, *443*, 67–70; b) S. Rendler, G. Auer, M. Oestreich, *Angew. Chem.* **2005**, *117*, 7793–7797; *Angew. Chem. Int. Ed.* **2005**, *44*, 7620–7624; c) T. Isobe, K. Fukuda, Y. Araki, T. Ishikawa, *Chem. Commun.* **2001**, 243–244.