

A Journal of the Gesellschaft Deutscher Chemiker


Angewandte Chemie

International Edition



www.angewandte.org

2009–48/43



Snapshots ...

... of the formation of a σ bond in Al_2R_4 isomers are obtained by structural and quantum chemical investigations, as described by H. Schnöckel, W. Kloppe, and co-workers in their Communication on page 8141 ff. By employing $\text{R} = \text{PtBu}_2$, they succeeded in isolating a triplet molecule exhibiting a long Al–Al separation as a first step. A bicyclic butterfly molecule rather than the unbridged R_2AlAlR_2 isomer represents the ground state. The ring-inversion isomers are separated by a “wing beat” and a transition state.

 WILEY-VCH

Inside Cover

Patrick Henke, Tobias Pankewitz, Wim Klopper,* Frank Breher, and Hansgeorg Schnöckel*

Snapshots of the formation of a σ bond in Al_2R_4 isomers are obtained by structural and quantum chemical investigations, as described by H. Schnöckel, W. Klopper, and co-workers in their Communication on page 8141 ff. By employing $\text{R} = \text{PrBu}_2$, they succeeded in isolating a triplet molecule exhibiting a long Al–Al separation as a first step. A bicyclic butterfly molecule rather than the unbridged R_2AlAlR_2 isomer represents the ground state. The ring-inversion isomers are separated by a “wing beat” and a transition state.

