

A Journal of the Gesellschaft Deutscher Chemiker

Angewandte Chemie

50
YEARS

International Edition



www.angewandte.org

2011–50/20



The growth mechanism, ...

... which plays a vital role in the rational design and precise fabrication of heterogeneous nanoparticles, can be unveiled by exploiting single-particle scattering spectra, as shown by T. Kang, J. Yi, and co-workers in their Communication on page 4633 ff. Three patterns of scattering spectra for individual gold/polystyrene dimeric nanoparticles are readily differentiated, and overgrowth mechanisms that correspond to each scattering spectrum are proposed.

 WILEY-VCH

Inside Cover

**Young In Yang, Eunhye Jeong, Inhee Choi, Suseung Lee,
Hyeon Don Song, Kihoon Kim, Yeonho Choi,* Taewook Kang,* and
Jongheop Yi***

The growth mechanism, which plays a vital role in the rational design and precise fabrication of heterogeneous nanoparticles, can be unveiled by exploiting single-particle scattering spectra, as shown by T. Kang, J. Yi, and co-workers in their Communication on page 4633 ff. Three patterns of scattering spectra for individual gold/polystyrene dimeric nanoparticles are readily differentiated, and overgrowth mechanisms that correspond to each scattering spectrum are proposed.

