

## Corrigendum

Primary and Secondary  
Aminophosphines as Novel  
P-Stereogenic Building Blocks for  
Ligand Synthesis

M. Revés, C. Ferrer, T. León, S. Doran,  
P. Etayo, A. Vidal-Ferran, A. Riera,\*  
X. Verdaguer\* **9452–9455**

*Angew. Chem. Int. Ed.* **2010**, 49

DOI 10.1002/anie.201004041

In this Communication, the enantiomeric excess reported for the hydrogenation of *N*-(3,4-dihydronaphthalen-2-yl)acetamide was incorrect. An unnoticed impurity contained in the racemic sample led to the use of an inappropriate HPLC method for the determination of the optical purity. Reanalyzing the sample with a correct HPLC method (Chiralcel OD-H, hexanes/isopropyl alcohol (95:5), 1.0 mL min<sup>-1</sup>, 210 nm, *t*(–) = 23.5 min, *t*(+) = 27.5 min)<sup>[1]</sup> showed that the reduction product was obtained in only 9% *ee*. The authors apologize for this error.

---

J. L. Renaud, P. Dupau, A.-E. Hay, M. Guingouain, P. H. Dixneuf, C. Bruneau, *Adv. Synth. Catal.* **2003**, 345, 230–238.