


Copper-Catalyzed Intramolecular Oxidative 6-*exo*-trig Cyclization of 1,6-Enynes with H₂O and O₂**

Zhi-Qiang Wang, Wen-Wu Zhang, Lu-Bin Gong, Ri-Yuan Tang, Xu-Heng Yang, Yu Liu, and Jin-Heng Li*

Transition-metal-catalyzed cyclization of 1,*n*-enynes has attracted significant interest because of their synthetic utility, and it is now used as a rapid and powerful approach to prepare cyclic derivatives in organic synthesis.^[1–8] In particular, these cyclization transformations are highly efficient and atom economical, and provide opportunities to discover new reactions. Despite considerable progress in the field, the development of mild 1,*n*-enyn cyclization routes using inexpensive catalytic materials to construct new, complex compounds remains a challenge. Copper catalysts are widely used in organic synthesis;^[2] however, examples of copper-catalyzed 1,*n*-enyn cyclizations are rare and all reports to date focus on the use of Cu^I salts.^[3] The key obstacle to Cu^I catalysts is that they have a far stronger affinity for alkynes than alkenes,^[2a, 4a–d] which results in the addition of more nucleophilic reagents (often H₂O, amines, acids, and alcohols) rather than alkenes to alkynes.^[2, 4] Although this obstacle does not favor the classical 1,*n*-enyn cyclization reaction, it may facilitate 1,*n*-enyn cyclizations that include the simultaneous introduction of other more nucleophilic reagents and alkenes to alkynes. After a series of trials, we found that Ce(SO₄)₂ facilitated CuCl₂-catalyzed oxidative 6-*exo*-trig cyclization of enynes with H₂O for preparing 1,4-naphthoquinones using O₂ as an oxidant and a reactant under mild reaction conditions (Scheme 1a).^[5, 6] The method is the first example of using a Cu^{II} catalyst for the oxidative cyclization of enynes, in which 1,4-naphthoquinones are constructed by the incorporation of two oxygen atoms from molecular oxygen^[5c] and water into the organic framework of the product.

Quinones, particularly 1,4-naphthoquinones, are valuable synthetic intermediates, and important structural units found in numerous natural products, pharmaceutical molecules



Scheme 1. Routes to substituted 1,4-naphthoquinones. DMA = dimethylacetamide, LG = leaving group.

(most notably Vitamin K), and functional materials (dyes and pigments).^[7] The traditional route to 1,4-naphthoquinones is the oxidation of substituted naphthols, naphthyl amines, or naphthalenes.^[8] However, this method often furnishes products in low to moderate yields, and has low functional group compatibility. Owing to these drawbacks, several powerful alternative methods have arisen, including (Scheme 1b): 1) the Diels–Alder reaction of substituted 1,3-dienes with preexisting quinones (path a),^[9] 2) the reaction of alkynes with metal carbonyls (path b),^[10] 3) thermo cycloisomerization of cyclobutenones (path c),^[11] and 4) the annulation of Fischer carbene complexes with alkynes (path d).^[12] As shown in Scheme 1, we herein describe an efficient example of achieving 1,4-naphthoquinones by catalytic oxidative 6-*exo*-trig cyclization of enynes with H₂O and O₂ using a simple and inexpensive copper salt as the catalyst. (Scheme 1).

Our investigation began with the reaction of (*E*)-3-phenyl-1-(2-(2-phenylethynyl)phenyl)prop-2-en-1-one (**1a**) with CuCl₂ and O₂ in THF at 80 °C; however, only a trace amount of the target 6-*exo*-trig cyclization product **2a** was observed (Table 1, entry 1).^[13] Substrate **1a** could be successfully cyclized with 10 mol % CuCl₂ and 1 atm O₂ at 80 °C, thus affording the desired product **2a** in 42 % yield, using DMF as the solvent (entry 2). It is noteworthy that the yield is reduced to 38 % using air instead of O₂ (entry 3). The yield was increased to 60 % when the reaction was carried out in DMA

[*] Z.-Q. Wang, L.-B. Gong, X.-H. Yang, Y. Liu, Prof. Dr. J.-H. Li
State Key Laboratory of Chemo/Biosensing and Chemometrics,
College of Chemistry and Chemical Engineering, Hunan University
Changsha 410082 (China)
E-mail: jhli@hunnu.edu.cn

Z.-Q. Wang, L.-B. Gong, X.-H. Yang
Key Laboratory of Chemical Biology & Traditional Chinese Medicine
Research (Ministry of Education), Hunan Normal University
Changsha 410081 (China)

W.-W. Zhang, R.-Y. Tang
College of Chemistry and Materials Engineering
Wenzhou University, Wenzhou, 325035 (China)

[**] We thank the Natural Science Foundation of China (No. 20872112) and the Scientific Research Fund of Hunan Provincial Education Department (No. 08A037) for financial support.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201104082>.

Table 2: (Continued)

Entry	1,4-Naphthoquinone 2	Yield [%] ^[b]	CuCl ₂ [mol %]	T [°C]	t [h]	Entry	1,4-Naphthoquinone 2	Yield [%] ^[b]	CuCl ₂ [mol %]	T [°C]	t [h]
5		2f 65	50	100	24	17		2r 65	30	100	24
6		2g 80	10	80	24	18		2s 61	50	100	24
7		2h 55	10	80	24	19		2t 80	30	100	24
8 ^[c]		2i 45	50	100	48	20		2u 67	50	100	24
9 ^[c]		2j 50	50	100	48	21		2v 39	30	100	24
10 ^[d]		2k 43	50	100	48	22		2w 36			
								4w 26	10	80	24
11		2l 76	10	80	24	23		2x 51	50	100	48
12		2m 69	10	80	24						

[a] Reaction conditions: **1a** (0.2 mmol), CuCl₂ (10 mol %), O₂ (1 atm), Ce(SO₄)₂·4H₂O (150 mol %), and DMA (0.1 % v/v of water; 3 mL) at 80 °C. [b] Yield of isolated product. [c] Substrate was consumed completely, and some unidentified products were observed by GC/MS analysis. [d] A by-product, (*E*)-1-(2-cinnamoylphenyl)-2-cyclohexenylethane-1,2-dione (**3k**), was isolated in 16 % yield.


tion. While substrates bearing an *ortho*- or *meta*-methyl group were treated with CuCl₂, O₂, and Ce(SO₄)₂ to afford the corresponding 6-*exo*-trig cyclization products **2b** and **2c** in low to moderate yields, the substrate with a *para*-methyl group provided the desired product **2d** in a good yield. Other substituents, such as Cl and NO₂ groups, were found to be tolerated under the optimal conditions (**2e–2g**). Notably, the thiophen-2-yl-substituted substrate successfully underwent the annulation reaction with CuCl₂, O₂, and Ce(SO₄)₂ in moderate yield (**2h**). Moderate yields were found to be still achievable from terminal alkynes, aliphatic alkynes, and enynes under the same reaction conditions (**2i–2k**). By using (*E*)-1-(2-ethynylphenyl)-3-phenylprop-2-en-1-one (a terminal alkyne), for instance, the cyclization reaction

proceeded smoothly to form the corresponding 2-benzoyl-naphthalene-1,4-dione (**2i**) in 45 % yield. Surprisingly, treatment of (*E*)-1-(2-(cyclohexenylethynyl)phenyl)-3-phenylprop-2-en-1-one, bearing a cyclohexenyl group, with CuCl₂, O₂, and Ce(SO₄)₂ afforded the cyclization/cyclohexenyl epoxidation product **2k** in 43 % yield together with a by-product, (*E*)-1-(2-cinnamoylphenyl)-2-cyclohexenylethane-1,2-dione (**3k**), in 16 % yield.^[14] Substrates with substituents such as Me, CF₃, and F, on the aromatic ring adjacent to a ketone were also suitable for the reaction (**2l–2n**). For example, an F-substituted substrate furnished the desired product **2n** in 70 % yield. The results disclosed that several functional groups, such as methyl, chloro, nitro, methoxy, iodo, and bromo groups on the aryl ring at the terminal prop-

2-en-1-one were perfectly compatible with the optimal conditions, affording the corresponding products **2o–2u** in moderate to good yields.

Importantly, the chloro, iodo, and bromo substitutions (**2p** and **2r–2u**) can be tolerated in this annulation reaction, thereby facilitating additional modifications at the halogenated positions. Moreover, some heterocycles (**2u**, **2w**, and **2x**) or olefins (**2v**) were introduced into this system, which also makes this methodology more useful for the preparation of pharmaceuticals and natural products.^[15] Interestingly, another 5-*exo*-dig cyclization product **4w** was isolated from the reaction of (*E*)-3-(furan-2-yl)-1-(2-(phenylethynyl)phenyl)prop-2-en-1-one besides the desired product **2w**. However, aliphatic alkyne only furnished the 6-*exo*-trig cyclization product **2x** in 51% yield. The reason may be that the vinylfuran moiety is highly active, thus resulting in the occurrence of a Diels–Alder reaction with arylalkynes in the presence of Lewis acids.^[16]


To elucidate the mechanism, some control experiments were carried out (Scheme 2).^[13] It was surprising to find that the reaction of substrate **1k** with H₂¹⁸O, CuCl₂, an additive (Ce(SO₄)₂ or K₂S₂O₈), and O₂ afforded the diketone product



Scheme 2. Control experiments.

[¹⁸O₂]-**3k** containing two ¹⁸O atoms [Eq. (1)].^[14] Unfortunately, the product [¹⁸O₂]-**3k** could not be converted into the desired cyclization product under standard conditions. It was interesting to disclose that the reaction of substrate **1a** with 10 equivalents of H₂¹⁸O, CuCl₂, and O₂ furnished the ¹⁸O-atom-containing product [¹⁸O]-**2a** as determined by GC/MS and HRMS (ESI) analysis in the presence/absence of additives [Eq. (2)]. The results reveal the source of two new oxygen atoms in product **2**: one oxygen atom is from water through a Wacker-type oxidation reaction,^[14] and the other oxygen atom is from O₂.^[5b] Moreover, the oxygen atom of the 2-benzoyl moiety is from H₂O according to authoritative ¹³C NMR spectra of products **2a** and [¹⁸O]-**2a**.^[17] The reaction between substrates **1a** and ethyl 2-diazoacetate (**5**) was carried out to examine whether the copper carbene intermediate exists:^[18] only product **2a** (45% yield) together with product **6** (93% yield) were obtained [Eq. (3)]. The carbene product **7k** was not observed from the reaction of **1k** with **5** [Eq. (4)]. The cyclization reactions of substrates **8**, either (*E*)-3-phenyl-1-(2-(2-phenylacetyl)phenyl)prop-2-en-1-one or (*E*)-1-phenyl-2-(2-(3-phenylacryloyl)phenyl)ethane-1,2-dione, did not take place under the optimal reaction conditions and only a trace of olefin Wacker oxidation products **9** was observed [Eq. (5)], thus indicating that the reaction does not include the alkyne hydrolysis.^[14,19] Notably, a stoichiometric amount of two radical inhibitors, TEMPO or 1,1-diphenylethane, were also tested with substrate **1a**: greater than 90% of **1a** was recovered together with a trace of target product **2a**, thereby suggesting that the reaction proceeds through a radical process.

Consequently, a possible mechanism as outlined in Scheme 3 is proposed.^[1,2,4,5] Complexation of CuCl₂ with substrate **1a** yields the intermediate **A**, followed by sequential Wacker oxidation of the C–C double bond with H₂O (intermediate **B**)^[14] and intramolecular cyclization to afford intermediate **C**. Oxidation of intermediate **C** with O₂ leads to the peroxycopper(III) intermediate **D** with the aid of the oxidative additive.^[5] Reductive elimination of intermediate **D** gives rise to the formation of the peroxycopper(II) complex **E**.^[5] Finally, the O–O bond cleavage of the peroxycopper(II) complex **E** and oxidation of the alcohol delivers the desired product **2a** and Cu(OH)Cl. The active Cu^{II} species can be regenerated by the reaction of Cu(OH)Cl with HCl.



Scheme 3. Possible mechanism.

There are at least two roles of the oxidative additives in this oxidation cyclization including a) oxidation of intermediate **C** to the Cu^{III} intermediate **D** and b) oxidation of the alcohol to the ketone in intermediate **E**.

In summary, we have described that inexpensive CuCl₂ can catalyze the intramolecular oxidative carbocyclization of (*E*)-3-aryl-1-(2-(arylethynyl)aryl)prop-2-en-1-ones with high substituent compatibility using O₂ as both the terminal oxidant and a reactant. The present method proceeds by a 6-*exo*-trig cyclization process, which provides a new approach that differs from the classical enyne cyclization method (generally by a 5-*exo*-dig cyclization).^[1] Moreover, the mechanism was discussed according to the ¹⁸O-labeling experiments and the results presented. Applications of this copper-catalyzed oxidative transformation in organic synthesis are currently underway in our laboratory.

Experimental Section

Typical experimental procedure for the CuCl₂-catalyzed intramolecular oxidative cyclization of alkynes with alkenes: (*E*)-3-aryl-1-(2-(2-arylethynyl)phenyl)prop-2-en-1-one **1** (0.2 mmol), CuCl₂ (10 mol %), Ce(SO₄)₂·4H₂O (150 mol %), and DMA (3 mL) were added to a Schlenk tube. The tube was then charged with O₂ (1 atm), and the reaction mixture stirred at 80 °C (oil bath temperature) for the indicated time until complete consumption of starting material as monitored by TLC and GC/MS analysis. After the reaction was finished, the reaction mixture was cooled to room temperature, diluted in ethyl acetate, and washed with brine. The aqueous phase was re-extracted with ethyl acetate. The combined organic extracts were dried over Na₂SO₄, concentrated under vacuum, and the resulting residue was purified by silica gel column chromatography (*n*-hexane/ethyl acetate 4:1) to afford the desired product **2**.

Received: June 14, 2011

Published online: August 19, 2011

Keywords: copper · enyne · heterocycles · oxidative cyclization · oxygen

- [1] For selected reviews, see: a) I. Ojima, M. Tzamarioudaki, Z. Li, R. Donovan, *J. Chem. Rev.* **1996**, 96, 635; b) C. Aubert, O. Buisine, M. Malacria, *Chem. Rev.* **2002**, 102, 813; c) C. Bruneau, *Angew. Chem.* **2005**, 117, 2380; *Angew. Chem. Int. Ed.* **2005**, 44, 2328; d) L. Zhang, J. Sun, S. Kozmin, *Adv. Synth. Catal.* **2006**, 348, 2271, and references cited therein; e) Z. Zhang, G. Zhu, X. Tong, F. Wang, X. Xie, J. Wang, L. Jiang, *Curr. Org. Chem.* **2006**, 10, 1457; f) V. Michelet, P. Y. Toullec, J.-P. Genêt, *Angew. Chem.* **2008**, 120, 4338; *Angew. Chem. Int. Ed.* **2008**, 47, 4268; g) E. Jiménez-Núñez, A. M. Echavarren, *Chem. Rev.* **2008**, 108, 3326.
- [2] a) S. R. Chemler, P. H. Fuller, *Chem. Soc. Rev.* **2007**, 36, 1153; b) I. P. Beletskaya, A. V. Cheprakov, *Coord. Chem. Rev.* **2004**, 248, 2337; c) S. V. Ley, A. W. Thomas, *Angew. Chem.* **2003**, 115, 5558; *Angew. Chem. Int. Ed.* **2003**, 42, 5400; d) W. Deng, L. Liu, Q.-X. Guo, *Chin. J. Org. Chem.* **2004**, 24, 150; e) G. Evans, N. Blanchard, M. Toumi, *Chem. Rev.* **2008**, 108, 3054; f) F. Monnier, M. Taillefer, *Angew. Chem.* **2009**, 121, 7088; *Angew. Chem. Int. Ed.* **2009**, 48, 6954.
- [3] a) C. Fehr, I. Farris, H. Sommer, *Org. Lett.* **2006**, 8, 1839; b) C. Fehr, J. Galindo, *Angew. Chem.* **2006**, 118, 2967; *Angew. Chem. Int. Ed.* **2006**, 45, 2901; c) N. T. Patil, H. Wu, Y. Yamamoto, *J. Org. Chem.* **2005**, 70, 4531.
- [4] a) T. Suzuki, R. D. Noble, C. A. Koval, *Inorg. Chem.* **1997**, 36, 136; b) P. Brandt, M. J. Södergren, P. G. Andersson, P.-O. Norrby, *J. Am. Chem. Soc.* **2000**, 122, 8013; c) B. F. Straub, I. Gruber, F. Rominger, P. Hofmann, *J. Organomet. Chem.* **2003**, 684, 124; d) S.-L. Zhang, L. Liu, Y. Fu, Q.-X. Guo, *Organometallics* **2007**, 26, 4546; e) D. W. Macomber, M. D. Rausch, *J. Am. Chem. Soc.* **1983**, 105, 5325; f) D. Bouyssi, N. Monteiro, G. Balme, *Tetrahedron Lett.* **1999**, 40, 1297; g) N. T. Patil, Y. Yamamoto, *J. Org. Chem.* **2004**, 69, 5139.
- [5] For selected papers on the use of Cu^{II}/O₂ catalytic system, see: a) T. Hamada, X. Ye, S. S. Stahl, *J. Am. Chem. Soc.* **2008**, 130, 833; b) C. Zhang, N. Jiao, *J. Am. Chem. Soc.* **2010**, 132, 28; c) S. Chiba, L. Zhang, J.-Y. Lee, *J. Am. Chem. Soc.* **2010**, 132, 7266; d) Y. Wei, H. Q. Zhao, J. Kan, W. P. Su, M. C. Hong, *J. Am. Chem. Soc.* **2010**, 132, 2522; e) B.-X. Tang, R.-J. Song, C.-Y. Wu, Y. Liu, M.-B. Zhou, W.-T. Wei, G.-B. Deng, D.-L. Yin, J.-H. Li, *J. Am. Chem. Soc.* **2010**, 132, 8900; f) L. Huang, H. Jiang, C. Qi, X. Liu, *J. Am. Chem. Soc.* **2010**, 132, 17652.
- [6] Until now, only few examples of the oxidative 1,*n*-enyne cyclization have been reported, which focused on the palladium catalysts in the presence of oxidants: the results showed that the oxidative 1,*n*-enyne cyclization provided a different chemoselectivity from the classical methodology using the similar substrates. See: a) L. L. Welbes, T. W. Lyons, K. A. Cychoz, M. S. Sanford, *J. Am. Chem. Soc.* **2007**, 129, 5836; b) X. Tong, M. Beller, M. K. Tse, *J. Am. Chem. Soc.* **2007**, 129, 4906; c) T. Tsujihara, K. Takenaka, K. Onitsuka, K. Hatanaka, H. Sasai, *J. Am. Chem. Soc.* **2009**, 131, 3452.
- [7] a) *Biochemistry of Quinones*, (Ed. R. A. Morton), Academic Press: New York, **1965**; b) R. H. Thompson, *Naturally Occurring Quinones*, 2nd ed., Academic Press, New York, **1971**; c) R. H. Thomson, *In The Chemistry of Quinonoid Compounds, Part 1*, (Ed.: S. Patai), Wiley-VCH, New York, **1974**, p. 111.
- [8] a) P. Monnet, F. Reverdin, E. Nölting, *Ber.* **1879**, 12, 2306; b) F. R. Japp, N. H. J. Miller, *J. Chem. Soc.* **1881**, 39, 220; c) N. H. J. Miller, *J. Russ. Phys. Chem. Soc.* **1884**, 16, 414; d) F. Russig, *J. Prakt. Chem.* **1900**, 62, 31; e) L. F. Fieser, M. Fieser, *J. Am. Chem. Soc.* **1935**, 57, 491; f) L. F. Fieser, *J. Am. Chem. Soc.* **1948**, 70, 3165; g) R. T. Arnold, R. Larson, *J. Org. Chem.* **1940**, 5, 250; h) R. E. Harman, J. Cason, *J. Org. Chem.* **1952**, 17, 1058.
- [9] a) K. Itami, T. Nokami, J. Yoshida, *Angew. Chem.* **2001**, 113, 1108; *Angew. Chem. Int. Ed.* **2001**, 40, 1074; b) K. Itami, T. Nokami, J.-i. Yoshida, *Adv. Syn. Catal.* **2002**, 344, 441.
- [10] a) L. S. Liebeskind, S. L. Baysdon, M. S. South, *J. Am. Chem. Soc.* **1980**, 102, 7397; b) L. S. Liebeskind, S. L. Baysdon, V. Goedken, R. Chidambaram, *Organometallics* **1986**, 5, 1086; c) L. S. Liebeskind, S. L. Baysdon, M. S. South, S. Iyer, J. P. Leeds, *Tetrahedron* **1985**, 41, 5839.
- [11] a) L. S. Liebeskind, S. Iyer, C. F. Jewell, *J. Org. Chem.* **1986**, 51, 3067; b) J. C. Bradley, T. Durst, A. J. Williams, *J. Org. Chem.* **1992**, 57, 6575.
- [12] a) W. D. Wulff, P.-C. Tang, J. S. McCallum, *J. Am. Chem. Soc.* **1981**, 103, 7677; b) W. D. Wulff, K.-S. Chan, P.-C. Tang, *J. Org. Chem.* **1984**, 49, 2293; c) J. P. A. Harrity, W. J. Kerr, D. Middlemiss, *Tetrahedron* **1993**, 49, 5565; d) M. L. Waters, M. E. Bos, W. D. Wulff, *J. Am. Chem. Soc.* **1999**, 121, 6403; e) J. C. Anderson, J. W. Cran, N. P. King, *Tetrahedron Lett.* **2002**, 43, 3849; f) C. R. Wu, D. O. Berbasov, W. D. Wulff, *J. Org. Chem.* **2010**, 75, 4441, and references cited therein.
- [13] See Table S1 and Figure S1 in the Supporting Information for detailed data. The single-crystal X-ray diffraction analysis of **2q** is given in Supporting Information.
- [14] Generally, the oxygen atom of the carbonyl group is from water in the Wacker-type oxidation reaction. See: a) J. Tsuji, *Pure Appl. Chem.* **1999**, 71, 1539; b) J. M. Takacs, X. Jiang, *Curr. Org. Chem.* **2003**, 7, 369; c) C. N. Cornell, M. S. Sigman, *Inorg. Chem.*

- 2007, 46, 1903; d) W. Ren, Y. Xia, S.-J. Ji, Y. Zhang, X. Wan, J. Zhao, *Org. Lett.* **2009**, 11, 1841.
- [15] a) S. L. Buchwald, S. M. King, *J. Am. Chem. Soc.* **1991**, 113, 258; b) T. Hayashi, F. T. Smith, K.-H. Lee, *J. Med. Chem.* **1987**, 30, 2005; c) A. Mital, V. S. Negi, U. Ramachandran, *Med. Chem.* **2008**, 4, 492.
- [16] a) G. Brieger, J. N. Bennett, *Chem. Rev.* **1980**, 80, 63; b) A. S. K. Hashmi, M. Rudolph, J. P. Weyrauch, M. Wölfe, W. Frey, J. W. Bats, *Angew. Chem.* **2005**, 117, 2858; *Angew. Chem. Int. Ed.* **2005**, 44, 2798; c) J. A. Cooper, P. Cornwall, C. P. Dell, D. W. Knight, *Tetrahedron Lett.* **1998**, 29, 2107.
- [17] The ^{13}C NMR (CD_3Cl) data of the carbonyl group in the 2-benzoyl moiety (see the Supporting Information): 192.917 ppm (product **2a**) and 192.940 ppm (product [^{18}O]-**2a**). See: a) M. T. Molina, C. Navarro, A. Moreno, A. G. Csáký, *Org. Lett.* **2009**, 11, 4938; b) P. A. Waske, J. Mattaya, M. Oelgemöller, *Tetrahedron Lett.* **2006**, 47, 1329.
- [18] For a paper on the reaction of the copper carbene with diazo compounds, see: P. Hofmann, I. V. Shishkov, F. Rominger, *Inorg. Chem.* **2008**, 47, 11755.
- [19] The experiments suggest that the reaction does not proceed by the alkyne hydrolysis leading to an α -ketone, then oxidative formation of the α -keto radical, and addition of this radical to the alkene. For selected papers on copper oxidation of enolates/enols leading to radicals, see: a) B. Zhao, H. Du, Y. Shi, *J. Am. Chem. Soc.* **2008**, 130, 7220; b) M. P. DeMartino, K. Chen, P. S. Baran, *J. Am. Chem. Soc.* **2008**, 130, 11546; c) Y. Wang, H. Du, *J. Org. Chem.* **2010**, 75, 3503, and references cited therein.