

Gold(I)-Catalyzed Stereoconvergent, Intermolecular Enantioselective Hydroamination of Allenes**

Kristina L. Butler, Michele Tragni, and Ross A. Widenhoefer*

The intermolecular, enantioselective addition of the N–H bond of an amine or carboxamide derivative across a C–C multiple bond (hydroamination) represents an attractive, atom-economical approach to the synthesis of chiral, non-racemic amines and amine derivatives.^[1] Within this family of transformations, the intermolecular enantioselective hydroamination (EHA) of allenes is of interest as a potentially expedient route to enantiomerically enriched α -chiral allylic amines, which are important chiral building blocks that are utilized in the synthesis of complex nitrogen-containing molecules.^[2] However, despite considerable efforts in this area,^[1] effective intermolecular EHA processes are scarce,^[3] and the intermolecular EHA of allenes remains unknown.^[4]

One of the challenges associated with the intermolecular EHA of allenes is the regioselectivity of extant hydroamination catalysts, which form predominantly achiral products from electronically unbiased monosubstituted allenes.^[4,5] To circumvent this regiochemical bias, we envisioned the stereoconvergent, intermolecular EHA of chiral, racemic 1,3-disubstituted allenes catalyzed by chiral bis(gold) phosphine complexes. This approach builds upon our previous efforts in the area of gold-catalyzed allene hydroamination.^[6,7] In particular, we have shown that achiral gold(I) N-heterocyclic carbene (NHC) complexes catalyze the regio- and diastereoselective hydroamination of chiral 1,3-disubstituted allenes with carbamates, and that allene racemization was rapid under reaction conditions.^[6] Furthermore, both we^[7] and Toste and co-workers^[8] have demonstrated the enantioselective intramolecular hydroamination of allenes catalyzed by chiral bis(gold) phosphine complexes.^[9] Herein we describe the stereoconvergent, enantioselective, intermolecular hydroamination of chiral, racemic 1,3-disubstituted allenes with carbamates catalyzed by chiral bis(gold) phosphine complexes.^[10]

Initial experiments directed toward the intermolecular EHA of allenes were only modestly encouraging. The reaction of benzyl carbamate (0.72 M) with 1-phenyl-1,2-butadiene (**1**; 1 equiv) catalyzed by a 1:2 mixture of [(*R*)-**2**](AuCl)₂] ((*R*)-**2** = (*R*)-DTBM-MeOBIPHEP, see structure

of (*S*)-**2** under Table 1) and AgOTf (OTf = trifluoromethanesulfonate) in dioxane at 24 °C for 24 h led to isolation of *N*-allylic carbamate (*R*)-**3a** in 35 % yield as a single diastereo-


Table 1: Effect of supporting ligand, solvent, silver salt, and allene concentration on the gold(I)-catalyzed enantioselective hydroamination of **1** with benzyl carbamate.

$\text{Ph}-\text{CH}=\text{C}(\text{Me})-\text{CH}_2 + \text{H}_2\text{NCbz} \xrightarrow[\text{solvent, 24 } ^\circ\text{C, 24 h}]{[\text{L}(\text{AuCl})_2] (2.5 \text{ mol } \%), \text{AgX} (5 \text{ mol } \%)}$ 1 (0.72 M) (0.72 M) 3a						
Entry	L	AgX	Solvent	Yield [%] ^[a]	ee [%]	Config.
1	(<i>R</i>)- 2	AgOTf	dioxane	35	50	<i>R</i>
2	(<i>R</i>)- 4	AgOTf	dioxane	15	44	<i>R</i>
3	(<i>S</i>)- 2	AgSbF ₆	dioxane	20	60	<i>S</i>
4	(<i>S</i>)- 2	AgPF ₆	dioxane	12	66	<i>S</i>
5	(<i>S</i>)- 2	AgClO ₄	dioxane	49	63	<i>S</i>
6	(<i>S</i>)- 2	AgNTf ₂	dioxane	20	45	<i>S</i>
7	(<i>S</i>)- 2	AgBF ₄	dioxane	71	72	<i>S</i>
8	(<i>S</i>)- 2	AgBF ₄	toluene	65	50	<i>S</i>
9 ^[b]	(<i>S</i>)- 2	AgBF ₄	THF	36	61	<i>S</i>
10 ^[c]	(<i>S</i>)- 2	AgBF ₄	dioxane	89	72 ^[d]	<i>S</i>

[a] Yield of isolated product; d.r. $\geq$ 25:1. Cbz = benzyloxycarbonyl.

[b] [**1**] = [H₂NCbz] = 0.4 M, 48 h, yield determined by GC. [c] [**1**] = 1.1 M.

[d] 96 % ee after recrystallization.



mer (d.r. $\geq$ 25:1) with 50 % ee (Table 1, entry 1).^[11,12] Substitution of ligand (*R*)-**2** with the SEGPHOS ligand (*R*)-**4** led to deterioration in both the yield and enantioselectivity of hydroamination (Table 1, entry 2). Conversely, optimization with respect to silver salt revealed that the use of AgBF₄ instead of AgOTf led to marked improvement in both yield and enantioselectivity of gold-catalyzed allene hydroamination (Table 1, entry 7). The reaction yield was further improved through employment of a slight excess of allene relative to benzyl carbamate and, in an optimized procedure, treatment of benzyl carbamate (0.72 M) with **1** (1.5 equiv) and a catalytic 1:2 mixture of [(*S*)-**2**](AuCl)₂] and AgBF₄ in dioxane at 24 °C led to isolation of (*S*)-**3a** in 89 % yield with 72 % ee (Table 1, entry 10). A single recrystallization from warm hexanes increased the enantiopurity of (*S*)-**3a** to 96 % ee.

In addition to benzyl carbamate, methyl carbamate; 9-fluorenylmethyl carbamate; and trichloroethyl carbamate

[*] K. L. Butler, M. Tragni, Prof. R. A. Widenhoefer
French Family Science Center, Duke University
Durham, NC 27708–0346 (USA)
E-mail: rwidenho@chem.duke.edu

[**] We thank the NIH (R01 GM080422-01) for support of this research. K.L.B. was supported in part through a Charles Bradsher Fellowship (Duke University) and M.T. was supported in part through a Marco Polo Fellowship (Alma Mater Studiorum—University of Bologna).

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201201584>.

underwent gold-catalyzed reaction with **1** to form *N*-allylic carbamates **3b–3d** with enantiopurities comparable to that of **3a** (Table 2, entries 1–3).^[13] Likewise, gold-catalyzed intermolecular EHA was effective for a number of 1-aryl-1,2-butadienes (Table 2, entries 4–13). For example, gold(I)-


Table 2: Enantioselective hydroamination of allenes (1.1 M) with carbamates (0.71 M) catalyzed by a mixture of $[(S)\text{-}2](\text{AuCl})_2$ (2.5 mol %) and AgBF_4 (5 mol %) in dioxane at room temperature.

Entry	Ar	R ^[a]	Product	Yield [%] ^[b]	ee [%] ^[c]
1	Ph (1)	CO ₂ Me	3b	81	69
2	Ph (1)	Fmoc	3c	79	71
3	Ph (1)	Troc	3d	43	73
4	X = Me (5a)	Cbz	6a	97	72
5	X = OMe (5b)	Cbz	6b	85	60
6	X = CF ₃ (5c)	Cbz	6c	82	75
7	X = Br (5d)	Cbz	6d	99	69 ^[d]
8	X = Me (5e)	Cbz	6e	82	81
9 ^[e]	X = <i>i</i> -Pr (5f)	Cbz	6f	80	81
10	X = Ph (5g)	Cbz	6g	86	86
11	X = Br (5h)	Cbz	6h	69	80
12 ^[e]	(5i)	Cbz	6i	42	92
13 ^[e]	(5j)	Cbz	6j	44	72

[a] Troc = 2,2,2-trichloroethoxycarbonyl, Fmoc = fluorenylmethyloxycarbonyl. [b] Yield of isolated product; d.r. ≥ 25:1. [c] Enantiomeric excess determined by HPLC analysis using chiral support. Absolute configuration assigned as *S* based on analogy to (*S*)-**3a**. [d] 99% ee after recrystallization. [e] Reaction run for 48 h.

catalyzed reactions of benzyl carbamate with *p*-substituted 1-aryl-1,2-butadienes **5a–5d** led to isolation of *N*-allylic carbamates **6a–6d** in 82–99% yield with 60–76% ee, albeit without a clear relationship between the electron-donating or -withdrawing properties of the arenes and the enantioselectivity (Table 2, entries 4–7). Also worth noting is that the enantiopurity of **6d** increased from 69 to 99% ee after a single recrystallization from warm hexanes (Table 2, entry 7). In comparison, gold-catalyzed hydroamination of the *ortho*-substituted 1-aryl-1,2-butadienes **5e–5h** formed *N*-allylic carbamates **6e–6h** with 80–86% ee (Table 2, entries 8–11). Gold(I)-catalyzed intermolecular hydroamination of the more sterically hindered *o,o*-disubstituted 1-aryl-1,2-butadiene **5i** occurred with even higher enantioselectivity (92% ee) but with diminished yield (Table 2, entry 12). Gold-

catalyzed intermolecular EHA was not restricted to 1-aryl-1,2-butadienes, and gold-catalyzed reaction of 1,3-dialkyl-substituted allene **7** with benzyl carbamate led to isolation of *N*-allylic carbamate **8** in 94% yield with 68% ee [Eq. (1)]. Conversely, gold-catalyzed intermolecular EHA was not effective for 1,3-disubstituted allenes lacking a methyl substituent.^[14]




Congruent with our expectations, allene racemization occurred rapidly under reaction conditions. For example, a solution of enantiomerically enriched (*S*)-**5b** (89% ee), benzyl carbamate, and a catalytic mixture of $[(S)\text{-}2](\text{AuCl})_2$ and AgBF_4 was periodically analyzed by HPLC using a chiral stationary phase; the analysis revealed complete racemization of (*S*)-**5b** prior to any detectable formation of **6b** (≤ 5 min; Table 3). Interestingly, continued analysis of the reaction mixture showed that the enantiopurity of **6b** decreased from 69% ee at 21% conversion to a terminal value of 59% ee at $\geq 74\%$ conversion (Table 3). This phenomenon was not

Table 3: Enantiopurity of allene and *N*-allylic carbamate as a function of conversion for the gold-catalyzed hydroamination of (*S*)-**5b** (89% ee).


<i>t</i> [h]	Conv. [%]	5b ee [%]	6b ee [%]
≤ 0.08	≤ 2	≤ 2	–
1.0	21	≤ 2	69
1.5	39	≤ 2	65
2.0	44	≤ 2	63
4.0	74	≤ 2	59
5.0	81	≤ 2	58
6.0	> 90	≤ 2	59

restricted to the formation of **6b**, and the enantiopurity of **3a** formed in the gold-catalyzed reaction of **1** with benzyl carbamate likewise decreased from 78% ee at 22% conversion to a terminal value of 72% ee at $\geq 83\%$ conversion.

Although we initially considered that racemization of **3a** or **6b** under the reaction conditions was responsible for the conversion-dependent enantioselectivity displayed by the gold-catalyzed intermolecular EHA of **1** or **5b**, respectively, this possibility was excluded. For example, stirring a solution of (*S*)-**3a** (72% ee) and a catalytic 1:2 mixture of $[(S)\text{-}2](\text{AuCl})_2$ and AgBF_4 at room temperature for eight hours either in the presence (1 equiv) or absence of benzyl carbamate led to no detectable decrease in the enantiopurity of (*S*)-**3a** [Eq. (2)]. Rather, a pair of experiments pointed to the potentially deleterious effect of the *N*-allylic carbamate product on the enantioselectivity of intermolecular EHA. For example, treatment of a 1.5:1:1 mixture of **1**, benzyl carbamate, and (*S*)-**6c** with a catalytic mixture of $[(S)\text{-}2](\text{AuCl})_2$ / AgBF_4 at 24 °C for 24 h led to isolation of (*S*)-**3a** in 75% yield



with 74% *ee* (Scheme 1), which is a nominally higher enantioselectivity than was realized in the absence of (*S*)-**6c** (Table 1, entry 10). However, the corresponding reaction of **1**, benzyl carbamate, and (*S*)-**6c** catalyzed by $[(R)-2](AuCl)_2/AgBF_4$ led to isolation of (*R*)-**3a** in 71% yield with 62% *ee* (Scheme 1), which is a considerably lower enantioselectivity than was observed in the absence of (*S*)-**6c**.



Scheme 1. Effect of *N*-allylic carbamate (*S*)-**6c** on the hydroamination of **1** with benzyl carbamate.

Rapid allene racemization precluded stereochemical analysis of the gold-catalyzed intermolecular EHA of (*S*)-**5b** (Table 3). However, we have previously established the *anti* stereochemistry of gold(I)-catalyzed intramolecular enantioselective allene hydrofunctionalization,^[7,15] which is consistent with the outer-sphere addition of the nucleophile to a gold(I) π -allene complex. Although such a mechanism requires that one of the two gold centers of a bis(gold) catalyst participates in the allene activation/C–N bond formation process, there is no firm evidence that supports the direct participation of the second gold center in these steps.^[10,16] There is, however, evidence that the ligation state of this spectator gold center can affect the enantioselectivity of gold-catalyzed hydrofunctionalization.^[8] Given the noncoordinating nature of the BF_4^- counterion employed in the gold-catalyzed intermolecular EHA of allenes, the spectator gold center is presumably ligated with an allene, benzyl carbamate, or either enantiomer of the *N*-allylic carbamate product.^[17] Therefore, the decreasing enantioselectivity of the gold-catalyzed intermolecular EHA of allenes with increasing conversion presumably reflects the increasing concentration of a less selective *N*-allylic carbamate ligated catalyst species.

In summary, we have developed a gold(I)-catalyzed protocol for the stereoconvergent, intermolecular enantioselective hydroamination of chiral, racemic 1,3-disubstituted allenes with *N*-unsubstituted carbamates. Furthermore, the observed decrease in enantioselectivity with increasing con-

version of intermolecular EHA coupled with independent analysis of the effect of *N*-allylic carbamate on the enantioselectivity of these transformations suggests that the nature of the catalytically active species changes with increasing concentration of *N*-allylic carbamate. We continue to work toward the identification of more selective and more general catalytic systems for intermolecular EHA and toward the development of ligand-modulated enantioselective gold catalysis.

Received: February 28, 2012

Published online: April 11, 2012

Keywords: allenes · asymmetric catalysis · asymmetric synthesis · gold · hydroamination

- [1] For reviews on catalytic hydroamination, see: a) S. R. Chemler, *Org. Biomol. Chem.* **2009**, 7, 3009–3019; b) T. E. Müller, K. C. Hultsch, M. Yus, F. Foubelo, M. Tada, *Chem. Rev.* **2008**, 108, 3795–3892; c) I. Aillaud, J. Collin, J. Hannedouche, E. Schulz, *Dalton Trans.* **2007**, 5105–5118; d) R. A. Widenhoefer, X. Han, *Eur. J. Org. Chem.* **2006**, 4555–4563; e) K. C. Hultsch, *Org. Biomol. Chem.* **2005**, 3, 1819–1824; f) K. C. Hultsch, *Adv. Synth. Catal.* **2005**, 347, 367–391.
- [2] For reviews regarding the synthesis of α -chiral allylic amines see: a) J. F. Hartwig, L. M. Stanley, *Acc. Chem. Res.* **2010**, 43, 1461–1475; b) B. M. Trost, T. Zhang, J. D. Sieber, *Chem. Sci.* **2010**, 1, 427–440; c) Z. Lu, S. Ma, *Angew. Chem.* **2008**, 120, 264–303; *Angew. Chem. Int. Ed.* **2008**, 47, 258–297; d) G. Helmchen, A. Dahnz, P. Dübon, M. Schelwies, R. Weihofen, *Chem. Commun.* **2007**, 675–691; e) B. M. Trost, M. L. Crawley, *Chem. Rev.* **2003**, 103, 2921–2943; f) T. K. Hollis, L. E. Overman, *J. Organomet. Chem.* **1999**, 576, 290–299; g) M. Johannsen, K. A. Jørgensen, *Chem. Rev.* **1998**, 98, 1689–1708; h) E. Skucas, M.-Y. Ngai, V. Komanduri, M. J. Krische, *Acc. Chem. Res.* **2007**, 40, 1394–1401; i) S. Kobayashi, H. Ishitani, *Chem. Rev.* **1999**, 99, 1069–1094; j) R. Bloch, *Chem. Rev.* **1998**, 98, 1407–1438.
- [3] For notable examples of enantioselective, intermolecular hydroamination, see: a) S. Pan, K. Endo, T. Shibata, *Org. Lett.* **2012**, 14, 780–783; b) A. L. Reznichenko, H. N. Nguyen, K. C. Hultsch, *Angew. Chem.* **2010**, 122, 9168–9171; *Angew. Chem. Int. Ed.* **2010**, 49, 8984–8987; c) Z. Zhang, S. D. Lee, R. A. Widenhoefer, *J. Am. Chem. Soc.* **2009**, 131, 5372–5373; d) J. Zhou, J. F. Hartwig, *J. Am. Chem. Soc.* **2008**, 130, 12220–12221; e) A. Hu, M. Ogasawara, T. Sakamota, A. Okada, K. Nakajima, T. Takahashi, W. Lin, *Adv. Synth. Catal.* **2006**, 348, 2051–2056; f) K. Li, P. N. Horton, M. B. Hursthouse, K. K. Hii, *J. Organomet. Chem.* **2003**, 665, 250–257; g) M. Utsunomiya, J. F. Hartwig, *J. Am. Chem. Soc.* **2003**, 125, 14286–14287; h) O. Löber, M. Kawatsura, J. F. Hartwig, *J. Am. Chem. Soc.* **2001**, 123, 4366–4367; i) M. Kawatsura, J. F. Hartwig, *J. Am. Chem. Soc.* **2000**, 122, 9546–9547; j) R. Dorta, P. Egli, F. Zürcher, A. Togni, *J. Am. Chem. Soc.* **1997**, 119, 10857–10858.
- [4] Yamamoto has reported the stereospecific, intermolecular hydroamination of chiral allenes: N. Nishina, Y. Yamamoto, *Angew. Chem.* **2006**, 118, 3392–3395; *Angew. Chem. Int. Ed.* **2006**, 45, 3314–3317.
- [5] a) J. F. Beck, J. A. R. Schmidt, *RSC Advances* **2012**, 2, 128–131; b) R. Kinjo, B. Donnadieu, G. Bertrand, *Angew. Chem.* **2011**, 123, 5674–5677; *Angew. Chem. Int. Ed.* **2011**, 50, 5560–5563; c) G. Kuchenbeiser, A. R. Shaffer, N. C. Zingales, J. F. Beck, J. A. R. Schmidt, *J. Organomet. Chem.* **2011**, 696, 179–187; d) A. N. Duncan, R. A. Widenhoefer, *Synlett* **2010**, 419–422; e) K. L. Toups, R. A. Widenhoefer, *Chem. Commun.* **2010**, 46,

- 1712–1714; f) X. Zeng, G. D. Frey, R. Kinjo, B. Donnadieu, G. Bertrand, *J. Am. Chem. Soc.* **2009**, *131*, 8690–8696; g) X. Zeng, M. Soleilhavoup, G. Bertrand, *Org. Lett.* **2009**, *11*, 3166–3169; h) N. Nishina, Y. Yamamoto, *Tetrahedron* **2009**, *65*, 1799–1808; i) V. Lavallo, G. D. Frey, B. Donnadieu, M. Soleilhavoup, G. Bertrand, *Angew. Chem.* **2008**, *120*, 5302–5306; *Angew. Chem. Int. Ed.* **2008**, *47*, 5224–5228; j) N. Nishina, Y. Yamamoto, *Tetrahedron Lett.* **2008**, *49*, 4908–4911; k) N. Nishina, Y. Yamamoto, *Synlett* **2007**, 1767–1770; l) R. O. Ayinla, L. L. Schafer, *Inorg. Chim. Acta* **2006**, *359*, 3097–3102; m) L. L. Anderson, J. Arnold, R. G. Bergman, *Org. Lett.* **2004**, *6*, 2519–2522; n) J. S. Johnson, R. G. Bergman, *J. Am. Chem. Soc.* **2001**, *123*, 2923–2924; o) M. Al-Masum, M. Meguro, Y. Yamamoto, *Tetrahedron Lett.* **1997**, *38*, 6071–6074; p) L. Besson, J. Goré, B. Cazes, *Tetrahedron Lett.* **1995**, *36*, 3857–3860; q) P. J. Walsh, A. M. Baranger, R. G. Bergman, *J. Am. Chem. Soc.* **1992**, *114*, 1708–1719.
- [6] R. E. Kinder, Z. Zhang, R. A. Widenhoefer, *Org. Lett.* **2008**, *10*, 3157–3159.
- [7] a) H. Li, S. D. Lee, R. A. Widenhoefer, *J. Organomet. Chem.* **2011**, *696*, 316–320; b) Z. Zhang, C. F. Bender, R. A. Widenhoefer, *Org. Lett.* **2007**, *9*, 2887–2889; c) Z. Zhang, C. F. Bender, R. A. Widenhoefer, *J. Am. Chem. Soc.* **2007**, *129*, 14148–14149.
- [8] a) R. L. LaLonde, Z. J. Wang, M. Mba, A. D. Lackner, F. D. Toste, *Angew. Chem.* **2010**, *122*, 608–611; *Angew. Chem. Int. Ed.* **2010**, *49*, 598–601; b) R. L. LaLonde, B. D. Sherry, E. J. Kang, F. D. Toste, *J. Am. Chem. Soc.* **2007**, *129*, 2452–2453; c) G. L. Hamilton, E. J. Kang, M. Mba, F. D. Toste, *Science* **2007**, *317*, 496–499.
- [9] For recent reviews of enantioselective gold(I) catalysis, see: a) A. Pradal, P. Y. Toullec, V. Michelet, *Synthesis* **2011**, 1501–1514; b) S. Sengupta, X. Shi, *ChemCatChem* **2010**, *2*, 609–619; c) R. A. Widenhoefer, *Chem. Eur. J.* **2008**, *14*, 5382–5391; d) N. Bongers, N. Krause, *Angew. Chem.* **2008**, *120*, 2208–2211; *Angew. Chem. Int. Ed.* **2008**, *47*, 2178–2181.
- [10] Che has recently reported the gold(I)-catalyzed intermolecular enantioselective hydroarylation of 1,3-diaryllallenes with indoles with up to 63% *ee*: M.-Z. Wang, C.-Y. Zhou, Z. Guo, E. L.-M. Wong, M.-K. Wong, C.-M. Che, *Chem. Asian J.* **2011**, *6*, 812–824.
- [11] a) The absolute configuration of (*S*)-**3a** was determined by optical rotation $[\alpha]_D^{25} = -39.4$ ($c = 0.19$, CH_2Cl_2); lit. [for (*S*)-**3a**] $[\alpha]_D^{23} = -55.4$ ($c = 1.10$, CH_2Cl_2).^[11b] The absolute configurations of all remaining *N*-allylic carbamates were assigned by analogy; b) J. C. A. Hunt, P. Laurent, C. J. Moody, *J. Chem. Soc. Perkin Trans. 1* **2002**, 2378–2389.
- [12] a) Allenes depicted in Tables 1 and 2 were synthesized in two steps from 3-butyne-2-ol in 26–51% yield employing the procedures of Knochel and co-workers^[12c] and Myers and Zheng,^[12b] with the exception of **1**, which was synthesized in one step from 4-phenyl-3-butyne-2-ol; b) A. G. Myers, B. Zheng, *J. Am. Chem. Soc.* **1996**, *118*, 4492–4493; c) N. Harrington-Frost, H. Leuser, M. I. Calaza, F. F. Kneisel, P. Knochel, *Org. Lett.* **2003**, *5*, 2111–2114.
- [13] Sulfonamides were less effective than were carbamates. For example, gold-catalyzed reaction of **1** with *p*-toluenesulfonamide at room temperature for 48 h formed (*E*)-4-methyl-*N*-(4-phenylbut-3-en-2-yl)benzenesulfonamide in 57% yield with 28% *ee*.
- [14] For example, gold-catalyzed reaction of 1-phenyl-1,2-pentadiene with benzyl carbamate at room temperature for 48 h formed (*E*)-benzyl (1-phenylpent-1-en-3-yl)carbamate in 13% yield with 23% *ee*.
- [15] Z. Zhang, R. A. Widenhoefer, *Angew. Chem.* **2007**, *119*, 287–289; *Angew. Chem. Int. Ed.* **2007**, *46*, 283–285.
- [16] J. H. Kim, S.-W. Park, S. R. Park, S. Lee, E. J. Kang, *Chem. Asian J.* **2011**, *6*, 1982–1986.
- [17] R. E. M. Brooner, R. A. Widenhoefer, *Organometallics* **2012**, *31*, 768–771.