

Rearrangements

Gold-Catalyzed Rearrangement of Allylic Oxonium Ylides: Efficient Synthesis of Highly Functionalized Dihydrofuran-3-ones**


Junkai Fu, Hai Shang, Zhaofeng Wang, Le Chang, Wenbing Shao, Zhen Yang,* and Yefeng Tang*

Rearrangements of allylic oxonium ylides have been recognized as powerful tools for the syntheses of oxygenated carbocycles.^[1] This type of transformation typically proceeds through a cascade carbenoid generation/oxonium ylide formation/[2,3]-sigmatropic rearrangement. The most common method to generate an oxonium ylide is the decomposition of an α -carbonyl diazo compound with a metal catalyst to form the metalcarbenoid, which is then trapped by the oxygen atom of an allylic ether (**A** $\rightarrow$ **C**, Scheme 1A).^[2] Given that diazo compounds are usually hazardous and explosive, and their preparation and handling are not trivial, the synthetic

utility of this type of transformation has been somewhat limited. In this context, the search for a system that is free of diazo compounds and thus less hazardous to promote the rearrangement of allylic oxonium ylides is of great value.^[3]

Recently, α -oxo gold carbenoids have received considerable attentions because they were assumed to be the key intermediates in a range of intriguing transformations,^[4] such as gold-catalyzed C–H insertion,^[5] O–H insertion,^[6] cyclopropanation,^[7] dipolar cycloaddition^[8] and others.^[9] Readily generated through intra- or intermolecular oxidation of gold-activated alkynes, α -oxo gold carbenoids display versatile reactivity, analogous to that of metalcarbenoids derived from diazo carbonyl compounds. For instance, Davies and Albrecht reported a gold-catalyzed [2,3]-sigmatropic rearrangement of sulfur ylides through an internal redox-combination strategy.^[10] More recently, Zhang and co-workers developed a facile synthesis of dihydrofuran-3-one derivatives from the readily available homopropargylic alcohols through a gold-catalyzed intramolecular O–H insertion.^[6a] Inspired by these seminal contributions, we envisioned that a similar strategy could be used in the [2,3]-sigmatropic rearrangement of oxonium ylides, as rationalized by the transformations from **E** $\rightarrow$ **D** (Scheme 1B). The proposed strategy would enable the [2,3]-sigmatropic rearrangement of oxonium ylides to proceed with readily available, diazo-free precursors under exceptionally mild conditions and with an operationally simple procedure. Moreover, the resulting product **D** features a highly functionalized dihydrofuran-3-one scaffold, which could be further elaborated into the core structures of various bioactive natural products, such as that of griseofulvin.^[1c]

To test the hypothesis, we initiated our studies by treatment of model substrate **1** with the conditions employed by Zhang and co-workers in their studies (Scheme 2).^[6a] The reaction was completed in three hours and led to the formation of two major products, one of which was the by-product resulting from direct deallylation of **1** (product structure not shown), and the other was the benzofuran allylic ether **2**. Although the expected [2,3]-sigmatropic rearrangement product **3** was not observed, we were pleased to find that **2** was converted almost quantitatively into **3** upon standing for 72 hours in CDCl_3 . The structure of **3** was unambiguously confirmed by X-ray crystallography.^[11] Subsequently, the conversion of **1** to **3** could be effected in one pot simply by increasing the reaction temperature to 60 °C. The observations indicated that the transformation of **1** to **3**, albeit formally a [2,3]-sigmatropic rearrangement, may indeed proceed through a mechanism distinctly different from the



Scheme 1. A) [2,3]-Sigmatropic rearrangement promoted by a metalcarbenoid that was derived from a diazo carbonyl compound. B) Proposed mechanism for the [2,3]-sigmatropic rearrangement promoted by an α -oxo gold carbenoid.

[*] J. Fu, Z. Wang, Dr. L. Chang, W. Shao, Prof. Dr. Z. Yang
Laboratory of Chemical Genomics, Peking University
Shenzhen Graduate School
Xili, Nanshan District, Shenzhen 518055 (China)
E-mail: zyang@pku.edu.cn
Dr. H. Shang, Prof. Dr. Y. Tang
The Comprehensive AIDS Research Center, Department of
Pharmacology & Pharmaceutical Sciences, School of Medicine
Tsinghua University
Beijing 100084 (China)
E-mail: yefengtang@tsinghua.edu.cn

[**] We acknowledge financial support from the National Science Foundation of China (21102081, 21272133), New Teachers' Fund for Doctor Stations, Ministry of Education (20110002120011), and opening foundation of Laboratory of Chemical Genomics, Peking University of Shenzhen Graduate School. We thank Dr. Tuoping Luo (H3 Biomedicine Inc., MA (USA)) and Prof. Zhigang Wang (Peking University) for the helpful discussions.


Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201208305>.

to 45 % (entry 5). The effect of substituents on the phenyl ring was also evaluated. Both electron-donating (*t*Bu-, Me-, or MeO-, **6f–i**) and electron-withdrawing (F- or Cl-, **6j** and **6k**) groups were tolerated well (entries 6–11). Naphthalene-derived compound **6l** also turned out to be a good substrate for the reaction (entry 12). The reactions also worked with **6m** and **6n**, which bear a 2,2- and 1,2-disubstituted alkene, respectively (entries 13 and 14). Notably, we observed the corresponding benzofuran allylic ether intermediates in the reactions with **6a–m**, suggesting that these reactions follow a stepwise process similar to that of compound **1**. In contrast, no such intermediate was detected for the reaction with **6n**. Moreover, both the substitution pattern and *E/Z* ratio of the alkene unit were fully preserved in the transformation from **6n** to **7n**, thus indicating that it proceeds through a [1,2] rearrangement, rather than a [2,3] rearrangement.

In order to further extend the scope of the reaction, various aliphatic substrates were examined (Table 3). The reactions of **8a–c**, in which the alkene and alkyne units are incorporated into a cyclic framework with either *cis* or *trans* configuration, proceeded smoothly to afford the corresponding products **9a–c** in excellent yields (entries 1–3). Notably, **9a** and **9b** were obtained as a single diastereoisomer, whereas **9c** was obtained as a mixture of C2 diastereoisomers (1:1 ratio). Interestingly, both diastereoisomers are *cis*- rather than *trans*-configured at the ring junction.^[16] This result might be attributed to a thermodynamically less stable 5–6 *trans*-fused

product, which upon formation gradually epimerizes into the corresponding *cis*-fused product. Furthermore, substrates with linear tethers were examined. To our delight, all of the reactions furnished the corresponding products in good yields (entries 4–6). Notably, unlike for the reactions with **6a–m**, the corresponding vinyl allylic ether intermediates were not observed in the reactions listed in Table 3.

To gain deeper insight into the gold-catalyzed rearrangement of allylic oxonium ylides, we performed deuterium-labeling experiments (Scheme S1, see Supporting Information), which, combined with the aforementioned experimental results, allowed us to propose a plausible mechanism (Scheme 3). Oxonium ylide **G**, which is generated from **E** via



Scheme 3. Proposed mechanisms for the gold-catalyzed rearrangement of allylic oxonium ylides.

Table 3: Gold-catalyzed rearrangement of allylic oxonium ylides with aliphatic moieties.^[a]


Entry	Substrate	<i>n</i>	Product	Yield ^[b] [%]
1		8a	9a	65
2		8b	9b	84
3 ^[c]		8c	9c	84
4		8d	9d	85
5		8e	9e	65
6 ^[c]		8f	9f	67

[a] Reactions were performed as described in Table 1. [b] Yields of isolated products. [c] The diastereoselectivity was determined by ¹H NMR spectroscopy. [d] d.r. = 1:1 (at C2). [e] d.r. = 3:2 (at C2).

α -oxo gold carbenoid **F**, may undergo three different pathways to give dihydrofuran-3-one **D** or its isomer **J**, depending on the nature of the substrates. The first pathway features a concerted [2,3]-sigmatropic rearrangement (path a), and is most likely involved in the reactions with **8a–f**, considering that 1) no vinyl allylic ether intermediates were observed in these reactions, and 2) inversion of the allyl moiety was observed in the deuterium-labeling experiment. In contrast, a stepwise mechanism may account for the reactions with **1** and **6a–m**, in which **G**, once generated, could rapidly tautomerize into the thermodynamically more stable intermediate **H**. Compound **H** undergoes a 1,4-allyl migration to afford **I**,^[17] which is readily converted into **D** through a Claisen rearrangement (path b).^[18] Moreover, a [1,2] rearrangement (path c) that leads to the formation of **J** cannot be completely ruled out in certain cases. Indeed, the transformation from **6n** to **7n** may follow this pathway.

To highlight the synthetic utility of the new reaction, we finally attempted to realize its asymmetric version. Significant efforts have been devoted to the asymmetric synthesis of dihydrofuran-3-ones through a [2,3]-sigmatropic rearrangement of oxonium ylides,^[19] however, only modest successes have been achieved. For instance, **1** was frequently employed as model substrate in the previous studies,^[19a–c] however, to date the highest *ee* value reported for **3** was 62 %.^[19b] We envisioned that the unique mechanism of the gold-catalyzed rearrangement of allylic oxonium ylides may offer an opportunity to address the challenge. Indeed, for those reactions proceeding through path b (Scheme 3), it is feasible to attempt the enantioselective synthesis of **D** through an

asymmetric Claisen rearrangement.^[20] As expected, after extensive screening of reaction conditions with **2** as model substrate (Table S1, Supporting Information), we were pleased to find that the desired transformation could be achieved by employing chiral bis(oxazoline)copper(II) complex **11** as the catalyst and CH₂Cl₂/Et₂O (1:1) as the solvent.^[21] Thus, **3** could be obtained almost quantitatively and with good enantioselectivity (80% *ee*, Scheme 4). The absolute config-



Scheme 4. Asymmetric Claisen rearrangement with vinyl allylic ether intermediates.

uration at C2 of **3** was assigned as S by comparison of the rotation of a derivative with reported data (for details, see the Supporting Information).^[22] Moreover, the asymmetric reaction could be applied to several other benzofuran allylic ethers (**10**, Scheme 4), all of which afforded the corresponding products with high yields (>95%) and good enantioselectivity (78–91% *ee*). Thus, our method offers an alternative to the existing repertoire of asymmetric [2,3]-sigmatropic rearrangements of oxonium ylides for the enantioselective synthesis of benzofuranone-3-one derivatives that bear an oxygenated chiral quaternary carbon atom.

In summary, we developed a novel gold-catalyzed rearrangement of allylic oxonium ylides, allowing the rapid access of highly functionalized dihydrofuran-3-ones from readily available precursors in one pot. Mechanistically, the rearrangement of oxonium ylides may proceed through a concerted [2,3]-sigmatropic rearrangement or stepwise 1,4-allyl migration followed by Claisen rearrangement. We also succeeded in developing an efficient approach to the enantioselective synthesis of benzofuranone-3-one derivatives through a Cu^{II}-catalyzed asymmetric Claisen rearrangement. Owing to the absence of diazo compounds and the operationally simple and step-economical nature, this variant of the rearrangement of allylic oxonium ylides is not only conceptually novel, but also synthetically useful. Further mechanistic studies and applications of this methodology to the total synthesis of natural products are in progress and will be communicated in due course.

Received: October 16, 2012

Revised: November 30, 2012

Published online: March 6, 2013

Keywords: Claisen rearrangement · gold · carbenoids · oxonium ylide · sigmatropic rearrangement

[1] For leading reviews, see: a) D. M. Hodgson, F. Y. T. M. Pierard, P. A. Stupple, *Chem. Soc. Rev.* **2001**, 30, 50–61; b) J. S. Clark, in

Nitrogen, Oxygen and Sulfur Ylide Chemistry: A Practical Approach, Oxford University Press, Oxford, **2002**; for representative examples, see: c) M. C. Pirrung, W. L. Brown, S. Rege, P. Laughton, *J. Am. Chem. Soc.* **1991**, 113, 8561–8562; d) F. P. Marmsäter, F. G. West, *J. Am. Chem. Soc.* **2001**, 123, 5144–5145; e) J. S. Clark, T. C. Fessard, C. Wilson, *Org. Lett.* **2004**, 6, 1773–1776; f) J. S. Clark, S. T. Hayes, C. Wilson, L. Gobbi, *Angew. Chem.* **2007**, 119, 441–444; *Angew. Chem. Int. Ed.* **2007**, 46, 437–440; g) J. S. Clark, C. A. Baxter, A. G. Dosseter, S. Poigny, L. L. Castro, W. G. Wittingham, *J. Org. Chem.* **2008**, 73, 1040–1055; h) D. M. Hodgson, D. Angrish, S. P. Erickson, J. Kloesges, C. H. Lee, *Org. Lett.* **2008**, 10, 5553–5556; i) J. S. Clark, R. Berger, S. T. Hayes, L. H. Thomas, A. J. Morrison, L. Gobbi, *Angew. Chem.* **2010**, 122, 10063–10066; *Angew. Chem. Int. Ed.* **2010**, 49, 9867–9870; j) J. S. Clark, D. Vignard, A. Parkin, *Org. Lett.* **2011**, 13, 3980–3983.

- [2] For selected examples, see: a) M. C. Pirrung, J. A. Werner, *J. Am. Chem. Soc.* **1986**, 108, 6060–6062; b) E. J. Roskamp, C. R. Johnson, *J. Am. Chem. Soc.* **1986**, 108, 6062–6063; c) F. P. Marmsäter, J. A. Vanecko, F. G. West, *Org. Lett.* **2004**, 6, 1657–1660; d) K. J. Quinn, N. A. Biddick, B. A. DeChristopher, *Tetrahedron Lett.* **2006**, 47, 7281–7283; e) J. S. Clark, S. B. Walls, C. Wilson, S. P. East, M. J. Drysdale, *Eur. J. Org. Chem.* **2006**, 323–327; f) J. S. Clark, C. Guerot, C. Wilson, A. Blake, *Chem. Commun.* **2007**, 4134–4136.
- [3] G. K. Murphy, F. G. West, *Org. Lett.* **2006**, 8, 4359–4361.
- [4] For a leading review, see: a) J. Xiao, X. W. Li, *Angew. Chem.* **2011**, 123, 7364–7375; *Angew. Chem. Int. Ed.* **2011**, 50, 7226–7236; for other related reviews, see: b) A. Fürstner, P. W. Davies, *Angew. Chem.* **2007**, 119, 3478–3519; *Angew. Chem. Int. Ed.* **2007**, 46, 3410–3449; c) S. F. Kirsch, *Synthesis* **2008**, 3183–3204; d) N. T. Patil, Y. Yamamoto, *Chem. Rev.* **2008**, 108, 3395–3442; e) A. Corma, A. Leyva-Pérez, M. J. Sabater, *Chem. Rev.* **2011**, 111, 1657–1712; f) N. D. Shapiro, F. D. Toste, *Synlett* **2010**, 675–691.
- [5] a) N. D. Shapiro, F. D. Toste, *J. Am. Chem. Soc.* **2007**, 129, 4160–4161; b) G. Li, L. M. Zhang, *Angew. Chem.* **2007**, 119, 5248–5251; *Angew. Chem. Int. Ed.* **2007**, 46, 5156–5159; c) L. Cui, G. Zhang, Y. Peng, L. Zhang, *Org. Lett.* **2009**, 11, 1225–1228; d) A. B. Cuenca, S. Montserrat, K. M. Hossain, G. Mancha, A. Lledós, M. Medio-Simón, G. Ujaque, G. Asensio, *Org. Lett.* **2009**, 11, 4906–4909.
- [6] a) L. Ye, L. Cui, G. Zhang, L. Zhang, *J. Am. Chem. Soc.* **2010**, 132, 3258–3259; b) L. Ye, W. He, L. Zhang, *J. Am. Chem. Soc.* **2010**, 132, 8550–8551.
- [7] a) D. Y. Qian, J. L. Zhang, *Chem. Commun.* **2011**, 47, 11152–11154; b) D. Vasu, H. H. Hung, S. Bhunia, S. A. Gawade, A. Das, R. S. Liu, *Angew. Chem.* **2011**, 123, 7043–7046; *Angew. Chem. Int. Ed.* **2011**, 50, 6911–6914.
- [8] a) H. S. Yeom, J. E. Lee, S. Shin, *Angew. Chem.* **2008**, 120, 7148–7151; *Angew. Chem. Int. Ed.* **2008**, 47, 7040–7043; b) A. M. Jadhav, S. Bhunia, H. Y. Liao, R. S. Liu, *J. Am. Chem. Soc.* **2011**, 133, 1769–1771.
- [9] a) G. Y. Lin, C. W. Li, S. H. Hung, R. S. Liu, *Org. Lett.* **2008**, 10, 5059–5062; b) A. S. K. Hashmi, M. Bührle, R. Salathé, J. W. Bats, *Adv. Synth. Catal.* **2008**, 350, 2059–2064; c) A. Mukherjee, R. B. Dateer, R. Chaudhuri, S. Bhunia, S. N. Karad, R. S. Liu, *J. Am. Chem. Soc.* **2011**, 133, 15372–15375; d) L. Ye, W. He, L. Zhang, *Angew. Chem.* **2011**, 123, 3294–3297; *Angew. Chem. Int. Ed.* **2011**, 50, 3236–3239; e) H. Yeom, Y. Lee, J. Jeong, E. So, S. Hwang, J. Lee, S. Lee, S. Shin, *Angew. Chem.* **2010**, 122, 1655–1658; *Angew. Chem. Int. Ed.* **2010**, 49, 1611–1614; f) C. W. Li, K. Pati, G. Y. Lin, H. H. Hung, R. S. Liu, *Angew. Chem.* **2010**, 122, 10087–10090; *Angew. Chem. Int. Ed.* **2010**, 49, 9891–9894; g) B. Lu, C. Q. Li, L. M. Zhang, *J. Am. Chem. Soc.* **2010**, 132, 14070–14072; h) W. He, C. Li, L. Zhang, *J. Am. Chem. Soc.* **2011**, 133, 8482; i) Y. Z. Wang, K. G. Ji, S. Lan, L. M. Zhang, *Angew. Chem.*

- 2012, 124, 1951–1954; *Angew. Chem. Int. Ed.* **2012**, 51, 1915–1918.
- [10] P. W. Davies, S. J.-C. Albrecht, *Angew. Chem.* **2009**, 121, 8522–8525; *Angew. Chem. Int. Ed.* **2009**, 48, 8372–8375.
- [11] CCDC 904703 (**3**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [12] For the Rh^{II}- or Cu^I-catalyzed [2,3]-sigmatropic rearrangement of oxonium ylides through a concerted mechanism, see references [2a] and [3]. For other similar cases, see: a) N. McCarthy, M. A. Mckerverey, T. Ye, M. McCann, E. Murphy, M. P. Doyle, *Tetrahedron Lett.* **1992**, 33, 5983–5986; b) N. Pierson, C. Fernández-Carcía, M. A. Mckerverey, *Tetrahedron Lett.* **1997**, 38, 4705–4708.
- [13] Lewis acids have rarely been used in reactions promoted by α -oxo gold carbenoids. In the reactions presented herein, Yb(OTf)₃ may play multiple roles, including: 1) activate the alkyne unit toward the nucleophilic addition of the pyridine *N*-oxide, 2) neutralize the pyridine derivative that is generated in the reaction to avoid a side reaction, and 3) facilitate the Claisen rearrangement.
- [14] L. Z. Dai, M. Shi, *Chem. Eur. J.* **2010**, 16, 2496–2502.
- [15] We obtained some unexpected results with a substrate bearing an alkynone (for details, see Scheme S2 in the Supporting Information).
- [16] The relative stereochemistry of **9c** was assigned by the NOE studies (for details, see the Supporting Information).
- [17] In order to explore whether the 1,4-allyl migration proceeds through a concerted or stepwise (dissociation/recombination) mechanism, we performed the crossover experiment with **6g** and **6m**. Only **7g** and **7m** were formed, which partly supported the concerted mechanism.
- [18] To the best of our knowledge, there is only one report in which a similar mechanism was assumed for the rearrangement of allylic oxonium ylides: a) J. S. Clark, G. Whitlock, S. Jiangand, N. Onyia, *Chem. Commun.* **2003**, 2578–2579. Furthermore, 1,4 migration was reported as the competitive reaction pathway in the Rh^{II}- or Cu^I-catalyzed [1,2] rearrangement of benzyl ethers or tertiary amines: b) N. P. Karche, S. M. Jachak, D. D. Dhavale, *J. Org. Chem.* **2001**, 66, 6323–6332; c) F. G. West, B. N. Naidu, R. W. Tester, *J. Org. Chem.* **1994**, 59, 6892–6894.
- [19] a) D. M. Hodgson, M. Petrolia, *Tetrahedron: Asymmetry* **2001**, 12, 877–881; b) S. Kitagaki, Y. Yanamoto, H. Tsutsui, M. Anada, *Tetrahedron Lett.* **2001**, 42, 6361–6364; c) N. Shimada, S. Nakamura, M. Anada, M. Shiro, S. Hashimoto, *Chem. Lett.* **2009**, 38, 488–489.
- [20] For a review, see: a) C. M. Krishna, A. Safiul, C. Buddhadeb, *Tetrahedron* **2008**, 64, 597–643; for selected examples, see: b) L. Abraham, R. Czerwonka, M. Hiersemann, *Angew. Chem.* **2001**, 113, 4835–4837; *Angew. Chem. Int. Ed.* **2001**, 40, 4700–4703; c) E. C. Linton, M. C. Kozlowski, *J. Am. Chem. Soc.* **2008**, 130, 16162–16163; d) C. Uyeda, E. N. Jacobsen, *J. Am. Chem. Soc.* **2008**, 130, 9228–9229; e) C. Uyeda, A. R. Rötheli, E. N. Jacobsen, *Angew. Chem.* **2010**, 122, 9947–9950; *Angew. Chem. Int. Ed.* **2010**, 49, 9753–9756.
- [21] a) A. Gille, M. Hiersemann, *Org. Lett.* **2010**, 12, 5258–5261; b) J. Troendlin, J. Rehbein, M. Hiersemann, O. Trapp, *J. Am. Chem. Soc.* **2011**, 133, 16444–16450.
- [22] T. Akiyama, T. Katoh, K. Mori, *Angew. Chem.* **2009**, 121, 4290–4292; *Angew. Chem. Int. Ed.* **2009**, 48, 4226–4228.