

Total Synthesis

Synthesis of *ent*-Kaurane and Beyerane Diterpenoids by Controlled Fragmentations of Overbred Intermediates**

Emily C. Cherney, Jason C. Green, and Phil S. Baran*

Dedicated to R. W. Hoffmann on the occasion of his 80th birthday

The *ent*-kauranes present highly varied oxidation patterns (**3**, **4**; Figure 1A), undergo intriguing skeletal rearrangements, and possess antibacterial, antitumor, and antimalarial activity.^[1] These attributes make *ent*-kauranes interesting candidates for two-phase terpene total synthesis.^[2] Steviol (**1**) was chosen as the lowest oxidized logical target due to the known conversion of such structures into beyeranes^[3] (isosteviol, **2**) and the useful functionality present for an “oxidase phase.” The first total synthesis reported by Mori et al. provides steviol (**1**) in 35 steps and 0.013 % overall yield.^[4a,b,c] A 19-step synthesis of steviol methyl ester was reported subsequently by Ziegler and Klock, but it relied on a key step that gave only a 3 % yield of the [3.2.1]bicyclic system and 0.015 % overall yield.^[4d] An elegant approach to isosteviol (**2**) was reported by Snider et al. in 13–18 steps, 0.37–1.2 % overall yield.^[5] Conversion of isosteviol (**2**) into steviol (**1**), however, is unknown. Herein, an efficient synthesis of (±)-steviol (**1**) is presented.

In 2009, Hoffmann formalized the concept of “overbred intermediates” in synthesis design as intermediates having one or more excess C–C bonds that must be subsequently cleaved.^[6] The route to the [3.2.1]bicyclic system of steviol (**1**) relies on the controlled fragmentation of two overbred intermediates. Cyclopropane **5** (Figure 1B) would require preferential cleavage of the C12–C16 bond over the C12–C13 bond. Such a fragmentation would be ambitious because these two bonds appear to be nearly indistinguishable.^[7] Indeed, a similar system fragmented with only modest diastereoselectivity (2:1).^[4c] Cyclobutane **6** would then arise from a [2+2] photocycloaddition with allene. This strategy should install a very hindered quaternary center with high diastereoselectivity.^[8] It is known that strained cyclobutanones in similar systems will open upon nucleophilic attack to break the analogous C13–C14 bond.^[9]

Tricyclic system **7** presents a challenge when the issues of stereo- and regioselectivity are considered. In particular, the required axial C19-methyl oxidation and *para* regioselectivity

are not adequately addressed by known approaches (Figure 1C). Radical-mediated methods give the undesired *ortho*-methoxy product.^[10] C–H activation reactions directed from C3 preferentially oxidize the C18-methyl group.^[11,12] Cyclizations initiated from a terminal epoxide (i.e. **10**, Scheme 1)





Figure 1. A) Truncated oxidation pyramid for *ent*-kauranes and beyeranes. B) Cyclase-phase retrosynthetic strategy. C) Polycyclization methods for installation of C18 or C19-methyl group oxidation.

[*] E. C. Cherney,^[‡] Dr. J. C. Green,^[‡] Prof. Dr. P. S. Baran
Department of Chemistry, The Scripps Research Institute
10550 North Torrey Pines Road, La Jolla, CA 92037 (USA)
E-mail: pbaran@scripps.edu

[‡] These authors contributed equally to this work.

[**] Financial support for this work was provided by an unrestricted grant from TEVA, the NIH (GM097444-03), and NSF (a predoctoral fellowship to E.C.C.). We are grateful to Prof. A. Rheingold and Dr. C. E. Moore (UCSD) for X-ray crystallographic analysis.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201304609>.



Scheme 1. Total synthesis of (±)-steviol (**1**). Reagents and conditions: a) LiNEt₂ (3 equiv), THF, 60 °C, 2 h (92 %); b) [VO(acac)₃] (0.25 equiv), *t*BuOOH (5 M in decane) (1.6 equiv), benzene, 6 °C, 2 h (74 % or 83 % BRSM); c) CBr₄ (3 equiv), PPh₃ (2.9 equiv), *i*Pr₂NEt (3.3 equiv), CH₂Cl₂, –10 °C, 12 h (81 %); d) NaOtBu (2.4 equiv) benzyl alcohol (solvent), 100 °C, 3 h (92 %); e) FeCl₃ (2 equiv), CH₂Cl₂, RT, 3 h (52 %); f) Li⁰ (50 equiv), NH₃, THF; *t*BuOH, –78 °C to –45 °C, 2 h; 4 M HCl in dioxane, RT, 30 min (79 %); g) DEAD (5 equiv), PPh₃ (5 equiv), THF, 70 °C, 5 h (91 %); h) H₂, Pd/C (10 wt %; 10 mol %), EtOAc, RT, 7 h (93 %); i) allene, CH₂Cl₂, RT, 450 W Hg lamp, pyrex, 12 h (82 %); j) O₃, MeOH, –78 °C, 5 min; Me₂S, RT, 30 min; AcOH/PPA (9:1), 110 °C, 12 h (62 %); k) HCl(g), Ac₂O (solvent), act. Zn⁰ (60 equiv), 0 °C, 45 min; l) AcCl (3 M in MeOH), 0–6 °C, 12 h (79 % **18** and 11 % **17**); m) PPh₃ (6.6 equiv), [RhCl(PPh₃)₃] (5 mol %), THF, *i*PrOH; TMSCHN₂ (20 equiv), 48 h (63 %); n) PDC, DMF, NaClO₂, NaH₂PO₄ (10 equiv), 2-methyl-2-butene (10 equiv), THF/*t*BuOH, 0 °C to RT, 16 h (85 %). acac = acetylacetonate, BRSM = based on recovered starting material, DEAD = diethyl azodicarboxylate, PPA = polyphosphoric acid, PDC = pyridinium dichromate, DMF = *N,N*-dimethylformamide, TMS = trimethylsilyl.

provide oxidation on the equatorial C18-methyl group.^[13] Consequentially, a unique cyclization precursor (**8**) was designed with the following considerations: 1) the polycyclization should be Lewis acid initiated (rather than radically initiated) to give the correct *para* regioselectivity; 2) the epoxide should be internal (rather than terminal) to give the required C19 oxidation;^[13] and 3) the *Z* stereochemistry of this internal epoxide is imperative to give C19 oxidation.^[14]


The pursuit of cyclization precursor **8** began from epoxide **9** (see Scheme 1). Elimination to open epoxide **9** followed by vanadium-directed epoxidation gave the *erythro* product **10** in 68 % overall yield (5.3 gram scale). The secondary alcohol in **10** was inverted to give the *threo* bromide **11** in 81 % yield (7.2 gram scale). Nucleophilic addition of benzyloxide to open the epoxide followed by closure of the bromohydrin provided the cyclization precursor **8** (7.1 gram scale). The polycyclization was most efficiently effected by iron trichloride to give tricyclic system **12** (1.1 gram scale). Compound **12** was converted into crystalline enone **13** by Birch reduction/deprotection and isomerization. X-ray analysis confirmed the *para* regiochemistry, the crucial C19 axial methyl group oxidation, and the correct stereochemistry at the C9-methine group.

Next, the neopentyl alcohol was eliminated, followed by hydrogenation to furnish compound **7** (2.1 gram scale). Birch reduction and isomerization proceeded to give enone **14** (1.3 gram scale). Allene [2+2] photocycloaddition with **14**

formed the hindered C8 quaternary center in overbred cyclobutane **6** (1.1 gram scale).^[8] The formation of this overbred intermediate was strategic because all other attempts to form this quaternary center failed, including: copper-, indium-, and tin-mediated 1,4-additions, Sakurai and Keck allylations, as well as intramolecular bond formations through sigmatropic rearrangements. Cyclobutane **6** was transformed to **15** in a one-pot sequence (1.0 gram scale): ozonolysis, selective fragmentation with methanol to give the methyl ester, and finally acid-mediated condensation to forge the [2.2.2]bicyclic system.^[9]

Reductive cyclopropanation of **15** would generate an overbred cyclopropanediol (**16**), which could undergo divergent fragmentation pathways: C12–C13 cleavage or C12–C16 cleavage to give **17** or **18**, respectively. Mori et al. treated a similar system with Zn(Hg) amalgam in 6 M HCl/toluene at 110 °C for 1 h to get a 2:1 ratio in favor of the analogous desired isomer in 41 % yield.^[4c] Treatment of diketone **15** with these conditions for 45 min gave *only* the undesired isomer **17** in 26 % yield (see Table 1, entry 1). Encouragingly, when the reaction was stopped after 5 min, a 2.2:1 ratio in favor of the desired isomer **18** was observed (entry 3). Moreover, the desired product **18** was found to rearrange to the undesired isomer **17** under acidic conditions (Scheme 2A). It seemed that Mori's conditions were unsuitable due to the high temperatures, which caused the desired kinetic isomer (**18**) to rearrange to the thermodynamic isomer (**17**). Under the

Table 1: Attempts at conversion of **15** or **5** into **18**.

diketone 15 or cyclopropane 5		conditions		
Entry	Conditions	Yield 18	Yield 17	Ratio 18 / 17
1	15 , Zn(Hg), 6 M HCl, PhMe, 110 °C, 45 min	0%	26%	0:1
2	15 , Zn(Hg), 6 M HCl, PhMe, 110 °C, 30 min	13%	9%	1.4:1
3	15 , Zn(Hg), 6 M HCl, PhMe, 110 °C, 5 min	24%	11%	2.2:1
4	15 , act. Zn ⁰ , HCl in Et ₂ O, 0 °C, 15 min	—	trace	—
5	5 , AcCl, MeOH, 0–6 °C, 12 h	79%	11%	7.2:1

reaction conditions, the ketones in products **17** and **18** likely undergo further reduction. Attempts to run this reaction at lower temperature with activated zinc led to decomposition, with trace formation of **17** (entry 4).

To overcome these issues, cyclopropane diol **16** was trapped as diacetate **5**.^[15] This would allow for fragmentation at low temperatures, thereby avoiding isomerization of kinetic product **18** to thermodynamic product **17**. It would also avoid over-reduction. Treatment of **5** with methanolic HCl at 0–6 °C gave **18** and **17** in 79% and 11% yield, respectively (greater than 7:1 ratio; entry 5). Undesired isomer **17** can also be recycled to **15** (Scheme 2B).

With suitable quantities of **18** in hand, installation of the methylene group was attempted (Scheme 1). While the Wittig olefination of a similar substrate has been reported,^[4a,c] this procedure as well as salt-free variations either yielded

rearranged material or gave no reaction, respectively. A modified Wittig procedure proceeded to give olefin **19**,^[16] which was oxidized to give (±)-steviol (**1**) in 17 steps from geranyl acetate.^[17] Acid-induced rearrangement of steviol (**1**) provided isosteviol (**2**; Scheme 2C). Alternatively, compound **19** could first be rearranged to the beyerane skeleton followed by Jones oxidation to provide isosteviol (**2**) in 17 steps (Scheme 2D).


To summarize, a synthetic route that offers efficient access to minimally oxidized members of the *ent*-kaurane and beyerane class of terpenes has been developed. This route could conceptually be rendered enantioselective.^[18] The challenging axial C19 oxidation and [3.2.1]bicyclic motifs prompted a reevaluation and strategic modification of literature precedent. The first challenge was addressed with a unique polycyclization precursor (**8**) while the second necessitated the use of overbred intermediates (**6** and **5**) and their controlled fragmentations. Such strained intermediates enabled and simplified the overall synthetic route. The completion of this cyclase phase sets the stage for an in-depth study of the oxidation chemistry of these complex terpenes.

Received: May 28, 2013

Revised: June 20, 2013

Published online: July 16, 2013

Keywords: beyerane · diterpenes · *ent*-kaurane · overbred intermediates · total synthesis



Scheme 2. Reagents and conditions: a) 12 M HCl, PhMe, 110 °C, 30 min (79%); b) CAN (3 equiv), MeCN, 0 °C, 10 min; c) PPA, AcOH, 110 °C, 12 h (72% over 2 steps); d) HBr (48% aq) Et₂O, RT, 15 h (90%); e) HBr (48% aq) Et₂O, RT, 18 h (87%); f) CrO₃ (10 equiv), acetone, 0 °C to RT, 3 h (81%). CAN = cerium ammonium nitrate.

- [1] H.-D. Sun, S.-X. Huang, Q.-B. Han, *Nat. Prod. Rep.* **2006**, *23*, 673.
- [2] a) Y. Ishihara, P. S. Baran, *Synlett* **2010**, 1733; b) K. Chen, P. S. Baran, *Nature* **2009**, *459*, 824.
- [3] M. Ruddat, E. Heftmann, A. Lang, *Arch. Biochem. Biophys.* **1965**, *110*, 496.
- [4] a) K. Mori, Y. Nakahara, M. Matsui, *Tetrahedron Lett.* **1970**, *11*, 2411; b) Y. Nakahara, K. Mori, M. Matsui, *Agric. Biol. Chem.* **1971**, *35*, 918; c) K. Mori, Y. Nakahara, M. Matsui, *Tetrahedron* **1972**, *28*, 3217; d) F. E. Ziegler, J. A. Klock, *Tetrahedron* **1977**, *33*, 373; e) I. F. Cook, J. R. Knox, *Tetrahedron Lett.* **1970**, *11*, 4091.
- [5] B. B. Snider, J. Y. Kiselgof, B. M. Foxman, *J. Org. Chem.* **1998**, *63*, 7945.
- [6] R. W. Hoffmann, *Elements of Synthesis Planning*, Springer, Berlin, **2009**, pp. 106–117, and references therein.
- [7] A. Abad, C. Agullo, I. Alfonso Marzal, I. Navarro, A. Gris, *Tetrahedron* **2006**, *62*, 3266.
- [8] a) M. T. Crimmins, T. L. Reinhold, *Org. React.* **1994**, *44*, 297; b) K. Wiesner, *Tetrahedron* **1975**, *31*, 1655.
- [9] a) H. Nagaoka, M. Shimano, Y. Yamada, *Tetrahedron Lett.* **1989**, *30*, 971; b) M.-A. Poupart, L. Paquette, *Tetrahedron Lett.* **1988**, *29*, 269.
- [10] a) B. B. Snider, R. Mohan, S. A. Kates, *J. Org. Chem.* **1985**, *50*, 3659; b) P. A. Zoretic, X. Weng, M. L. Caspar, D. G. Davis, *Tetrahedron Lett.* **1991**, *32*, 4819; c) L. Chen, G. B. Gill, G. Pattenden, *Tetrahedron Lett.* **1994**, *35*, 2593; d) S. Rendler, D. W. C. MacMillan, *J. Am. Chem. Soc.* **2010**, *132*, 5027.
- [11] a) J. E. Baldwin, R. H. Jones, C. Najera, M. Yus, *Tetrahedron* **1985**, *41*, 699; b) L. V. Desai, K. L. Hull, M. S. Sanford, *J. Am. Chem. Soc.* **2004**, *126*, 9542; c) E. M. Simmons, J. F. Hartwig, *Nature* **2012**, *483*, 70.

- [12] A. García-Granados, P. E. Lopez, E. Melguizo, A. Parra, Y. Simeo, *J. Org. Chem.* **2007**, 72, 3500.
- [13] a) E. E. Van Tamelen, S. R. Zawacky, R. K. Russell, J. G. Carlson, *J. Am. Chem. Soc.* **1983**, 105, 142; b) E. J. Corey, K. Liu, *J. Am. Chem. Soc.* **1997**, 119, 9929.
- [14] a) G. Stork, A. W. Burgstahler, *J. Am. Chem. Soc.* **1955**, 77, 5068; b) A. Eschenmoser, L. Ruzicka, O. Jeger, D. Arigoni, *Helv. Chim. Acta* **1955**, 38, 1890.
- [15] M. Toda, M. Hayashi, Y. Hirata, S. Yamamura, *Bull. Chem. Soc. Jpn.* **1972**, 45, 264.
- [16] H. Lebel, D. Guay, V. Paquet, K. Huard, *Org. Lett.* **2004**, 6, 3047.
- [17] E. J. Corey, G. Schmidt, *Tetrahedron Lett.* **1979**, 20, 399.
- [18] E. J. Corey, M. C. Noe, W.-C. Shieh, *Tetrahedron Lett.* **1993**, 34, 5995.
-