


Pyridine-Directed Palladium-Catalyzed Phosphonation of C(sp²)–H Bonds**

Changkun Li, Takaaki Yano, Naoki Ishida, and Masahiro Murakami*

Direct functionalization of carbon–hydrogen (C–H) bonds provides a straightforward means of molecular transformation, and thus has been extensively investigated in the past two decades.^[1,2] Although a wide variety of transition-metal-catalyzed reactions for functionalizing C–H bonds are currently available, examples of carbon–phosphorus bond formation are significantly limited, presumably owing to the strong coordinating character of phosphorus reagents.^[3,4] Existence of an excess amount of coordinative phosphorus over metal in a reaction media would hamper a process to activate less coordinative C–H bonds. Very recently, Yu and co-workers reported a pyridine-directed C–H phosphonation reaction catalyzed by palladium, in which *H*-phosphonates were directly used.^[5] Expedient deactivation of the catalyst was avoided by adding *H*-phosphonate slowly with a syringe pump. Herein, we describe our independent study of the analogous phosphonation reaction of 2-arylpyridines; an α -hydroxyalkylphosphonate generates *H*-phosphonate upon treatment with a base,^[6] which serves as the masked phosphonating reagent^[7] to save the catalyst from deactivation. Furthermore, step-by-step stoichiometric reactions clearly delineate the mechanistic features.

2-Phenylpyridine (**1a**) was treated with commercially available *H*-phosphonate **2** in the presence of palladium(II) acetate (10 mol %), *N*-methylmaleimide (NMMI, 40 mol %), silver(I) acetate (2.5 equiv), and K₂HPO₄ (4.5 equiv) at 120 °C for 48 h (Scheme 1). The *ortho* C–H bond was phosphonated to give product **4a** in only 12 % yield. Next, *H*-phosphonate **2** was replaced by α -hydroxyalkylphosphonate **3**, which was easily prepared from **2** and acetone in one step according to the reported procedure.^[8] To our surprise, **4a** was obtained in 70 % yield, together with a small amount of diphosphonated product (6 %). Thus, α -hydroxyphosphonate **3** proved to be superior to **2** as the phosphonating reagent, probably because it gradually generated *H*-phosphonate **2**.^[9] Other reaction conditions were examined using **3** as the masked phospho-



Scheme 1. C–H phosphonation of **1a**.

nating reagent. The use of oxidants such as Cu(OAc)₂ and Ag₂CO₃ were less effective. Stronger bases such as K₃PO₄ gave inferior results. Reaction in other solvents, including toluene, dioxane, and acetonitrile, also gave the phosphonated products, but the yields were lower.^[10]

Various substituted 2-arylpyridines were phosphonated using phosphonating reagent **3** (Scheme 2). The substrate 2-(*o*-tolyl)pyridine (**1b**) allowed the isolation of monophosphonated product **4b** in 66 % yield. The *m*-tolyl group was site-selectively phosphonated on the sterically less-hindered side to afford **4c** in 82 % yield. Methoxy (**4d**) and chloro (**4e**) groups were tolerated on the phenyl ring. Benzothiophene (**4f**) and alkene (**4g**) were also phosphonated in good yields. Not only pyridine, but also quinoline (**4i**) and pyrimidine (**4k**) were suitable as directing groups.


A proposed mechanism is shown in Scheme 3. Initially, cyclopalladation of **1a** with palladium(II) acetate gives palladacycle **A**.^[11] The α -hydroxyalkylphosphonate **3** gradually releases acetone under the reaction conditions to generate a small amount of *H*-phosphonate **2**, which reacts with palladacycle **A** to displace the acetate ligand on palladium. The resulting palladium(II) complex **B** undergoes reductive elimination with the aid of NMMI (see below). Arylphosphonate **4a** is thus produced, along with a palladium(0) species, which is oxidized back into palladium(II) by silver(I) acetate.^[12]

We carried out some stoichiometric reactions to corroborate the steps constituting the proposed catalytic cycle. Complex **A** was prepared by treatment of **1a** with palladium(II) acetate (1.0 equiv) in MeOH and isolated in a form of the dimer **5**.^[11] Next, a dioxane solution containing complex **5**, phosphonate **3** (2.0 equiv), and K₂HPO₄ (2.2 equiv) was heated at 120 °C (Scheme 4). The acetate ligand of **A** was displaced with a phosphonate ligand to afford complex **B**, which was isolated in the dimeric form **6**. Single crystals of **6**,

[*] Dr. C. Li, T. Yano, Dr. N. Ishida, Prof. Dr. M. Murakami
Department of Synthetic Chemistry and Biological Chemistry
Kyoto University, Katsura, Kyoto 615-8510, (Japan)
E-mail: murakami@sbchem.kyoto-u.ac.jp
Homepage: <http://www.sbchem.kyoto-u.ac.jp/murakami-lab/>


[**] We thank Dr. Y. Nagata (Kyoto Univ.) for his assistance with X-ray crystallographic analysis. This work was supported in part by a Grant-in-Aid for Scientific Research on Innovative Areas "Molecular Activation Directed toward Straightforward Synthesis" and a Grant-in-Aid for Scientific Research (B) from MEXT and The Asahi Glass Foundation. C.L. thanks the Japan Society for the Promotion of Science for a Research Fellowship.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201305202>.




Scheme 2. Palladium-catalyzed phosphonation of **1**. Reaction conditions: **1** (0.20 mmol), **3** (1.5 equiv), Pd(OAc)₂ (10 mol %), NMMI (40 mol %), AgOAc (2.5 equiv), K₂HPO₄ (4.5 equiv), *t*BuOH (1.5 mL), 120°C, 48 h. Yields shown are of isolated products.

suitable for X-ray crystallographic analysis, were obtained by recrystallization from an ether/hexane solution. The six-membered dinuclear palladacycle is bridged with the phosphonate ligand through its oxygen and phosphorus atoms (Figure 1).^[13] It assumes a boat-like form; the O(2), Pd(1), P(1), O(1) atoms are located on a plane [O(2)–Pd(1)–P(1)–O(1)=3.54°] and the torsion angle of the Pd(1)–P(1)–O(1)–



Scheme 3. Proposed mechanism.



Scheme 4. Ligand exchange.





Figure 1. Crystal structure of **6**. Thermal ellipsoids set at 50%. Hydrogen atoms and three carbon atoms of the *n*-butyl groups are omitted for clarity. Selected bond lengths (Å) and angles (°): Pd(1)–P(1) 2.227(1), Pd(1)–C(1) 1.998(2), C(1)–Pd(1)–P(1) 96.00(6), O(2)–Pd(1)–P(1)–O(1) 3.54, Pd(1)–P(1)–O(1)–Pd(2) 55.85.


Pd(2) was 55.85°. The phosphorus atom was located *cis* to the phenyl carbon, for which stronger *trans* influence was expected than for the pyridine nitrogen.^[14]

The following reductive elimination step was also examined.^[15] No reaction was observed upon heating a CD₃CN solution of **6** at reflux. On the contrary, reductive elimination occurred when heated at reflux (bath temperature: 120°C) in the presence of NMMI (2.0 equiv), resulting in the isolation of arylphosphonate **4a** in 97% yield (Scheme 5). The use of benzoquinone and maleic anhydride instead of NMMI was equally effective in inducing reductive elimination (60% after 48 h). These results suggested that electron-accepting ligands are required to facilitate reductive elimination of arylphosphonate **4a** from palladacycle **B**.^[16]

The reductive elimination giving **4a** from **6** was accompanied by the formation of black precipitates, presumably an aggregate of palladium(0). The formation of a palladium(0)



Scheme 5. Reductive elimination.



Scheme 6. Detection of palladium(0) complex 7.

species was confirmed by the reaction in the presence of acetylene dicarboxylate and BINAP (Scheme 6). The palladium(0) complex **7**^[17] was isolated in 29% yield together with **4a** (90%). Thus, the mechanistic pathway of the phosphonation of 2-phenylpyridine, proceeding through cyclometallation, ligand exchange, and reductive elimination, was followed step-by-step on the basis of stoichiometric reactions, thus providing an experimental basis for the postulated mechanism depicted in Scheme 3.

In conclusion, we have reported that the palladium-catalyzed phosphonation of a C(sp²)-H bond occurs by the use of a pyridyl group as the directing group. This work demonstrates the potential of α -hydroxyalkylphosphonates as masked phosphonating agents to save the catalyst from deactivation.

Experimental Section

Palladium-catalyzed phosphonation of **1a**: Pd(OAc)₂ (4.5 mg, 0.02 mmol), *N*-methylmaleimide (8.8 mg, 0.08 mmol), AgOAc (84 mg, 0.5 mmol), and K₂HPO₄ (156 mg, 0.9 mmol). *t*BuOH (1.5 mL), α -hydroxyalkylphosphonate **3** (75 mg, 0.30 mmol), and 2-phenylpyridine **1a** (31 mg, 0.20 mmol) were added by syringe into a 25 mL screw-capped tube equipped with a magnetic stir bar. The reaction tube was capped in the air and heated at 120°C for 48 h. After cooling to room temperature, the reaction mixture was filtered through a short pad of celite, using ethyl acetate as the eluent. The solvent was removed under reduced pressure and the residue was then purified by chromatography to afford **4a** (48.6 mg, 70%).

Received: June 17, 2013

Published online: July 23, 2013

Keywords: C–H activation · palladium catalysis · phosphonation · synthetic methods · transition metals

- [1] a) *Activation of Unreactive Bonds and Organic Synthesis* (Ed.: S. Murai), Springer, Berlin, **1999**; b) *Handbook of C–H Transformations* (Ed.: G. Dyker), Wiley-VCH, Weinheim, **2005**; c) *Topics in Current Chemistry, C–H Activation* (Eds.: J.-Q. Yu, Z.-J. Shi), Springer, Berlin, **2010**; d) O. Daugulis, H.-Q. Do, D. Shabashov, *Acc. Chem. Res.* **2009**, *42*, 1074; e) R. Giri, B.-F. Shi, K.-M. Engle, N. Maugel, J.-Q. Yu, *Chem. Soc. Rev.* **2009**, *38*, 3242; f) D. A. Colby, R. G. Bergman, J. A. Ellman, *Chem. Rev.* **2010**, *110*, 624; g) I. A. I. Mkhaliid, J. H. Barnard, T. B. Marder, J. M. Murphy, J. F. Hartwig, *Chem. Rev.* **2010**, *110*, 890; h) T. W. Lyons, M. S. Sanford, *Chem. Rev.* **2010**, *110*, 1147; i) L. Ackermann, *Chem. Commun.* **2010**, *46*, 4866; j) T. Newhouse, P. S. Baran, *Angew. Chem.* **2011**, *123*, 3422; *Angew. Chem. Int. Ed.* **2011**, *50*, 3362; k) L. McMurray, F. O'Hara, M. J. Gaunt, *Chem. Soc. Rev.* **2011**, *40*, 1885; l) C. S. Yeung, V. M. Dong, *Chem. Rev.*

- 2011**, *111*, 1215; m) C.-L. Sun, B.-J. Li, Z.-J. Shi, *Chem. Rev.* **2011**, *111*, 1293; n) S. H. Cho, J. Y. Kim, J. Kwak, S. Chang, *Chem. Soc. Rev.* **2011**, *40*, 5068; o) J. Yamaguchi, A. D. Yamaguchi, K. Itami, *Angew. Chem.* **2012**, *124*, 9092; *Angew. Chem. Int. Ed.* **2012**, *51*, 8960; p) N. Kuhl, M. N. Hopkinson, J. Wencel-Delord, F. Glorius, *Angew. Chem.* **2012**, *124*, 10382; *Angew. Chem. Int. Ed.* **2012**, *51*, 10236; q) P. B. Arockiam, C. Bruneau, P. H. Dixneuf, *Chem. Rev.* **2012**, *112*, 5879.
- [2] For palladium-catalyzed C–H functionalization of 2-phenylpyridines, see: a) A. R. Dick, K. L. Hull, M. S. Sanford, *J. Am. Chem. Soc.* **2004**, *126*, 2300; b) D. Kalyani, N. R. Deprez, L. V. Desai, M. S. Sanford, *J. Am. Chem. Soc.* **2005**, *127*, 7330; c) K. L. Hull, W. Q. Anani, M. S. Sanford, *J. Am. Chem. Soc.* **2006**, *128*, 7134; d) D. Kalyani, A. R. Dick, W. Q. Anani, M. S. Sanford, *Tetrahedron* **2006**, *62*, 11483; e) H.-Y. Thu, W.-Y. Yu, C.-M. Che, *J. Am. Chem. Soc.* **2006**, *128*, 9048; f) X. Chen, C. E. Goodhue, J.-Q. Yu, *J. Am. Chem. Soc.* **2006**, *128*, 12634; g) D. C. Powers, T. Ritter, *Nat. Chem.* **2009**, *1*, 302; h) X. Zhao, E. Dimitrijevic, V. M. Dong, *J. Am. Chem. Soc.* **2009**, *131*, 3466; i) X. Wang, L. Truesdale, J.-Q. Yu, *J. Am. Chem. Soc.* **2010**, *132*, 3648; j) Y. Kuninobu, T. Iwanaga, T. Omura, K. Takai, *Angew. Chem.* **2013**, *125*, 4527; *Angew. Chem. Int. Ed.* **2013**, *52*, 4431.
- [3] a) T. Kagayama, A. Nakano, S. Sakaguchi, Y. Ishii, *Org. Lett.* **2006**, *8*, 407; b) Y. Gao, G. Wang, L. Chen, P. Xu, Y. Zhao, Y. Zhou, L.-B. Han, *J. Am. Chem. Soc.* **2009**, *131*, 7956; c) Y. Kuninobu, T. Yoshida, K. Takai, *J. Org. Chem.* **2011**, *76*, 7370; d) C. Hou, Y. Ren, R. Lang, X. Hu, C. Xia, F. Li, *Chem. Commun.* **2012**, *48*, 5181; e) C.-B. Xiang, Y. J. Bian, X.-R. Mao, Z.-Z. Huang, *J. Org. Chem.* **2012**, *77*, 7706; f) O. Berger, C. Petit, E. L. Deal, J.-L. Montchamp, *Adv. Synth. Catal.* **2013**, *355*, 1361.
- [4] For phosphonation of aromatic C–H bonds using a stoichiometric amount of a transition-metal complex, see: a) V. I. Sokolov, L. L. Troitskaya, O. A. Reutov, *J. Organomet. Chem.* **1980**, *202*, C58; b) C. Bolm, K. Wenz, G. Raabe, *J. Organomet. Chem.* **2002**, *662*, 23; c) V. A. Stepanova, V. V. Dunina, I. P. Smoliakova, *Organometallics* **2009**, *28*, 6546.
- [5] C.-G. Feng, M. Ye, K.-J. Xiao, S. Li, J.-Q. Yu, *J. Am. Chem. Soc.* **2013**, *135*, 9322.
- [6] M. S. Kharasch, R. A. Mosher, I. S. Bengelsdorf, *J. Org. Chem.* **1960**, *25*, 1000.
- [7] a) D. W. Allen, D. E. Hibbs, M. B. Hursthouse, K. M. A. Malik, *J. Organomet. Chem.* **1999**, *572*, 259; b) G. A. Stark, T. H. Riermeier, M. Beller, *Synth. Commun.* **2000**, *30*, 1703; c) M. Hayashi, T. Matsuura, I. Tanaka, H. Ohta, Y. Watanabe, *Org. Lett.* **2013**, *15*, 628.
- [8] M. Kyoda, M. Hirata, *PCT Int. Appl.* 2006049010, A1, **2006**.
- [9] When **3** was treated with K₂HPO₄ in CD₃CN at 120°C for 18 h, a trace amount of *H*-phosphonate **2** was produced.
- [10] See the Supporting Information for details.
- [11] a) A. Kasahara, *Bull. Chem. Soc. Jpn.* **1968**, *41*, 1272; b) E. C. Constable, A. M. W. C. Thompson, T. A. Leese, D. G. F. Reese, D. A. Tocher, *Inorg. Chim. Acta* **1991**, *182*, 93.
- [12] a) Y. Fujiwara, I. Moritani, M. Matsuda, S. Teranishi, *Tetrahedron Lett.* **1968**, *9*, 3863; b) D. R. Stuart, E. Villemure, K. Fagnou, *J. Am. Chem. Soc.* **2007**, *129*, 12072; c) Y. Lu, D. Leow, X. Wang, K. M. Engle, J.-Q. Yu, *Chem. Sci.* **2011**, *2*, 967.
- [13] CCDC 945235 (**6**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [14] The crystal structure of palladacycle **5**^[2e] indicates stronger *trans* influence of the phenyl carbon than the pyridine nitrogen; the Pd–O bond *trans* to carbon (avg. 2.158 Å) is longer than that *trans* to pyridine (avg. 2.061 Å).
- [15] For reductive elimination of arylphosphonates from organopalladium complexes, see: a) R. A. Stockland, Jr., A. M. Levine, M. T. Giovine, I. A. Guzei, J. C. Cannistra, *Organometallics*

2004, 23, 647; b) M. C. Kohler, R. A. Stockland Jr., N. P. Rath, *Organometallics* **2006**, 25, 5746; c) M. Kalek, J. Stawinski, *Organometallics* **2008**, 27, 5876; d) M. C. Kohler, T. V. Grimes, X. Wan, T. R. Cundari, R. A. Stockland, Jr., *Organometallics* **2009**, 28, 1193.

- [16] For reductive elimination promoted by electron deficient alkenes, see: a) T. Yamamoto, A. Yamamoto, S. Ikeda, *J. Am. Chem. Soc.* **1971**, 93, 3350; b) R. Sustmann, J. Lau, *Chem. Ber.* **1986**, 119, 2531.
- [17] J. Caeiro, D. Peña, A. Cobas, D. Pérez, E. Guitián, *Adv. Synth. Catal.* **2006**, 348, 2466.
-