

Enantioselective N–H Insertion Reaction of α -Aryl α -Diazoketones: An Efficient Route to Chiral α -Aminoketones**

Bin Xu, Shou-Fei Zhu,* Xiao-Dong Zuo, Zhi-Chao Zhang, and Qi-Lin Zhou*

Abstract: A highly enantioselective N–H insertion reaction of α -diazoketones was developed by using cooperative catalysis by dirhodium(II) carboxylates and chiral spiro phosphoric acids. The insertion reaction provides a new access route to diverse chiral α -aminoketones, which are versatile building blocks in organic synthesis, with fast reaction rates, good yields and high enantioselectivity under mild and neutral conditions.

The construction of carbon–heteroatom bonds (C–X) is a central task for studies in organic synthesis.^[1] C–X bonds can be formed directly by transition-metal-catalyzed insertion of carbenes into heteroatom–hydrogen bonds (X–H, where X = N, O, S, etc.).^[2] Recently, significant progress in catalytic asymmetric X–H insertion reactions has been made, for example, high enantioselectivities have been achieved with carbene insertions into N–H, O–H, and S–H bonds, which all contain heteroatoms with lone-pair electrons.^[3] However, α -diazoesters are the only carbene precursors that have been used for these asymmetric X–H insertion reactions, which produce chiral α -heteroatom-substituted esters. Therefore, the potential applications of these reactions are severely limited. α -Diazoketones, another readily available carbene precursor, have been used in a few rhodium-catalyzed asymmetric cyclopropanation reactions, in C–H and Si–H insertion reactions, and in tandem reactions involving ylides.^[2,4] X–H insertion reactions of α -diazoketones produce α -heteroatom-substituted ketones, which are useful building blocks for organic synthesis. However, when we used 1-diazo-1-phenylpropan-2-one (**2a**) in the asymmetric insertion reaction with *tert*-butyl carbamate (BocNH₂) catalyzed by [Rh₂(*R*-DOSP)₄], [Rh₂(*S*-PTAD)₄], or a chiral copper complex with a spirobisoxazoline ligand (**4**; Figure 1), no enantioselectivity was observed (Table 1, entries 1–3).^[5] The problem with the asymmetric insertion of α -diazoketones





Figure 1. The catalysts and ligands used in this study.

into X–H bonds in which X has lone-pair electrons is that the strongly electron-withdrawing carbonyl group promotes the formation of free ylide **III**^[2,6] or enol **IV**,^[7] and chiral catalyst is thus not present during the chirality-determining proton-transfer step (Scheme 1).



Scheme 1. Proposed pathway of X–H insertions of α -diazoketones.

Herein, we report a highly enantioselective N–H insertion reaction of α -diazoketones co-catalyzed by dirhodium(II) carboxylates and chiral spiro phosphoric acids (SPAs, **1**), in which chiral induction was achieved in the proton-transfer step.^[8] The mild, neutral conditions bring special advantages for the preparation of synthetically important chiral α -aminoketones,^[9] which can epimerize under harsh conditions.

[*] B. Xu, Prof. S.-F. Zhu, X.-D. Zuo, Z.-C. Zhang, Prof. Q.-L. Zhou
State Key Laboratory and Institute of Elemento-organic Chemistry,
Nankai University, Tianjin 300071 (China)

and

Collaborative Innovation Center of Chemical Science and

Engineering, Tianjin (China)

E-mail: sfzhu@nankai.edu.cn

qlzhou@nankai.edu.cn

Homepage: <http://zhou.nankai.edu.cn/>


[**] We thank the National Natural Science Foundation of China and the National Basic Research Program of China (2012CB821600), the “111” project (B06005) of the Ministry of Education of China, and the Program for New Century Excellent Talents in University (NCET-10-0516) for financial support.



Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201400236>.

Initially, we performed the N–H insertion reaction of α -diazoketone **2a** with BocNH₂ by using 1 mol % [Rh₂(OAc)₄] and 1 mol % SPA **1** as the catalysts (Table 1). SPAs with various substituents at the 6 and 6' positions were evaluated (Table 1, entries 4–11), and the 6,6'-di-(4-PhC₆H₄)-substituted SPA **1c** afforded the highest enantioselectivity (52% *ee*; Table 1, entry 6). The type of dirhodium(II) complex strongly

Table 1: Enantioselective N–H insertion reaction of α -diazoketone **2a**: optimization of the reaction conditions.



Entry ^[a]	[M]	SPA	Yield [%] ^[b]	<i>ee</i> [%] ^[c]
1 ^[d]	Cu ^I /4	none	78	0
2	[Rh ₂ (R-DOSP) ₄]	none	70	1
3	[Rh ₂ (S-PTAD) ₄]	none	72	1
4	[Rh ₂ (OAc) ₄]	1a	79	14
5	[Rh ₂ (OAc) ₄]	1b	76	50
6	[Rh ₂ (OAc) ₄]	1c	85	52
7	[Rh ₂ (OAc) ₄]	1d	84	25
8	[Rh ₂ (OAc) ₄]	1e	88	37
9	[Rh ₂ (OAc) ₄]	1f	75	17
10	[Rh ₂ (OAc) ₄]	1g	85	8
11	[Rh ₂ (OAc) ₄]	1h	76	4
12	[Rh ₂ (oct) ₄]	1c	75	47
13	[Rh ₂ (piv) ₄]	1c	89	58
14	[Rh ₂ (TPA) ₄]	1c	87	67
15	[Rh ₂ (TFA) ₄]	1c	85	93
16 ^[e]	[Rh ₂ (TFA) ₄]	1c	80	96
17 ^[f]	none	1c	–	–


[a] Reaction conditions: [Rh]/SPA/**2a**/BocNH₂ = 0.002:0.002:0.21:0.2 (mmol) in 2 mL of CHCl₃ at 25 °C. [b] Yield of isolated product.

[c] Determined by HPLC. [d] With 5 mol % catalyst, 4 h. [e] A mixed solution of two substrates was added over 5 min by a pump. [f] No insertion product was detected and **2a** was recovered in 93% yield.

affected the enantioselectivity of the reaction (Table 1, entries 12–15). Specifically, when we used dirhodium(II) carboxylates, including [Rh₂(TPA)₄], which gives the best results in the N–H insertion reaction of α -diazoesters,^[8a] only moderate enantioselectivity was obtained. In sharp contrast, dirhodium(II) trifluoroacetate, a strong Lewis acid, significantly increased the enantioselectivity to 93% *ee* (Table 1, entry 15). The enantioselectivity was further improved to 96% *ee* by addition of the mixed solution of **2a** and BocNH₂ with a syringe pump over the course of 5 min (Table 1, entry 16). A control experiment showed that the SPA did not catalyze the insertion reaction in the absence of a dirhodium complex (Table 1, entry 17). Although the reaction afforded comparable enantioselectivity (93% *ee*) when it was performed without argon protection, the yield of desired N–H insertion product markedly decreased to 32%. The lower yield is attributed to significant dimerization of α -diazoketones.

Using the optimal reaction conditions, we investigated the substrate scope of the asymmetric N–H insertion reaction of

Table 2: Enantioselective N–H insertion reaction of α -diazoketones: substrate scope.




Entry	R ¹	R ²	Product	Yield [%]	<i>ee</i> [%] ^[11]
1	C ₆ H ₅	Me	3a	80	96(R)
2	4-MeC ₆ H ₄	Me	3b	79	94
3	4-BrC ₆ H ₄	Me	3c	82	93
4	4-FC ₆ H ₄	Me	3d	78	93
5	3-MeOC ₆ H ₄	Me	3e	80	95
6	3-MeC ₆ H ₄	Me	3f	82	96
7	3-FC ₆ H ₄	Me	3g	75	90
8	2-FC ₆ H ₄	Me	3h	86	86
9	2-Naph	Me	3i	95	97
10	C ₆ H ₅	Et	3j	85	97(R)
11	C ₆ H ₅	<i>n</i> Bu	3k	78	98(R)
12	C ₆ H ₅		3l	82	96(R)
13	C ₆ H ₅		3m	46	95(R)
14	C ₆ H ₅		3n	56	92
15	C ₆ H ₅		3o	85	95(R)
16	C ₆ H ₅	C ₆ H ₅	3p	85	96(R)
17	C ₆ H ₅	4-MeOC ₆ H ₄	3q	89	98
18	C ₆ H ₅	4-FC ₆ H ₄	3r	78	96
19	C ₆ H ₅		3s	79	94

The reaction conditions and analysis method were the same as those described for Table 1, entry 16. All of the reactions had run to completion within 1 min after the addition of diazo compound **2**.

α -diazoketones^[10] (Table 2). All of the tested α -aryl- α -diazoketones (**2a–2i**) reacted completely in 1 min to afford the corresponding chiral α -aminoketones **3a–3i** in good yields with high enantioselectivity (Table 2, entries 1–9). The electronic properties of the substituent at the *para* or *meta* position of the phenyl ring of the diazoketone affected the yields and enantioselectivity only slightly, whereas an *ortho*-fluoro substituent lowered the enantioselectivity slightly (86% *ee*; Table 2, entry 8). We also studied the effect of the R² substituent. Excellent enantioselectivity (92–98% *ee*) was obtained when R² was either an alkyl group (**2j–2o**; Table 2, entries 10–15) or an aryl group (**2p–2s**; Table 2, entries 16–19). The insertion reaction tolerated a range of functional groups, such as alkenyl, methoxy, silyloxy, and *N*-methyl pyrrole (Table 2, entries 13–15 and 19). In the reactions of **2m**, which has a terminal olefin, and **2n**, which has a methoxy group, the occurrence of side reactions (intramolecular cyclopropanation and C–H insertion) decreased the yields (Table 2, entries 13 and 14).

To demonstrate the utility of this asymmetric N–H insertion reaction of α -diazoketones, we carried out various reactions with the chiral α -aminoketone products (Scheme 2). Ketone **3p** could be reduced with NaBH₄ to chiral α -amino alcohol **5a** in quantitative yield with high stereoselectivity (*anti/syn* > 20:1, 97% *ee*; Scheme 2A).^[12] Upon addition of



Scheme 2. Transformations of the chiral α -aminoketone products.

a Grignard reagent, **3p** could also be easily converted to α -amino alcohol **5b**, which has a quaternary chiral center, in 90% yield and with excellent stereoselectivity (*anti/syn* > 20:1, 97% ee; Scheme 2B). The configuration of **5b** was determined to be (1*R*, 2*S*) by means of X-ray diffraction analysis of a single crystal.^[13] Removal of the protecting group from **3s** afforded **6**, which is a key intermediate for the synthesis of monoamine oxidase inhibitor **7** (Scheme 2C).^[9e] According to the reported procedure,^[9g] chiral insertion product **3o** could be readily converted to (L)-(-)-733,061, which is a selective antagonist for the NK1 receptor (Scheme 2D).

In conclusion, we have described the first asymmetric N–H insertion reaction of α -diazoketones, which was achieved by cooperative catalysis by achiral dirhodium(II) carboxylates and chiral SPAs. This new reaction provides an efficient and straightforward route to synthetically important chiral α -aminoketones with good yields and high enantioselectivity. This study markedly broadens the scope of asymmetric X–H insertion reactions.

Experimental Section

The $[\text{Rh}_2(\text{F}_3\text{CCO}_2)_4]$ (1.3 mg, 0.002 mmol, 1 mol%) and (*R*)-**1c** (1.2 mg, 0.002 mmol, 1 mol%) were introduced into an oven-dried Schlenk tube in an argon-filled glovebox. After 2 mL CHCl_3 was injected into the Schlenk tube, the mixture was stirred at 25 °C. A solution of 1-diazo-1-phenylpropan-2-one **2a** (33.6 mg, 0.21 mmol) and BocNH_2 (23.4 mg, 0.2 mmol) in 1 mL CHCl_3 was introduced into the mixture by a syringe pump over 5 min. The color of the diazo compound disappeared immediately after the addition. TLC showed that the reaction ran to completion as soon as the addition finished. The reaction mixture was then concentrated and purified by flash

chromatography on silica gel (petroleum ether/ethyl acetate = 8:1) to give **3a** as a white solid.

Received: January 9, 2014

Published online: March 5, 2014

Keywords: asymmetric catalysis · carbenes · chiral proton shuttle · chiral α -aminoketones · diazo compounds

- [1] A. K. Yudin, *Catalyzed Carbon–Heteroatom Bond Formation*, Wiley-VCH, Weinheim, **2011**.
- [2] For reviews: a) M. P. Doyle, M. A. McKerverey, T. Ye, *Modern Catalytic Methods for Organic Synthesis with Diazo Compounds*, Wiley, New York, **1998**; b) C. J. Moody, *Angew. Chem.* **2007**, *119*, 9308–9310; *Angew. Chem. Int. Ed.* **2007**, *46*, 9148–9150; c) Z.-H. Zhang, J.-B. Wang, *Tetrahedron* **2008**, *64*, 6577–6605.
- [3] For a review: a) S.-F. Zhu, Q.-L. Zhou, *Acc. Chem. Res.* **2012**, *45*, 1365–1377; for O–H insertions: b) T. C. Maier, G. C. Fu, *J. Am. Chem. Soc.* **2006**, *128*, 4594–4595; c) C. Chen, S.-F. Zhu, B. Liu, L.-X. Wang, Q.-L. Zhou, *J. Am. Chem. Soc.* **2007**, *129*, 12616–12617; d) S.-F. Zhu, Y. Cai, H.-X. Mao, J.-H. Xie, Q.-L. Zhou, *Nat. Chem.* **2010**, *2*, 546–551; e) T. Osako, D. Panichakul, Y. Uozumi, *Org. Lett.* **2012**, *14*, 194–197; f) X.-G. Song, S.-F. Zhu, X.-L. Xie, Q.-L. Zhou, *Angew. Chem.* **2013**, *125*, 2615–2618; *Angew. Chem. Int. Ed.* **2013**, *52*, 2555–2558; for N–H insertions: g) B. Liu, S.-F. Zhu, W. Zhang, C. Chen, Q.-L. Zhou, *J. Am. Chem. Soc.* **2007**, *129*, 5834–5835; h) E. C. Lee, G. C. Fu, *J. Am. Chem. Soc.* **2007**, *129*, 12066–12067; i) Z.-R. Hou, J. Wang, P. He, J. Wang, B. Qin, X.-H. Liu, L.-L. Lin, X.-M. Feng, *Angew. Chem.* **2010**, *122*, 4873–4876; *Angew. Chem. Int. Ed.* **2010**, *49*, 4763–4766; j) X.-F. Xu, P. Y. Zavalij, M. P. Doyle, *Angew. Chem.* **2012**, *124*, 9967–9971; *Angew. Chem. Int. Ed.* **2012**, *51*, 9829–9833; for S–H insertions: k) Y.-Z. Zhang, S.-F. Zhu, Y. Cai, H.-X. Mao, Q.-L. Zhou, *Chem. Commun.* **2009**, 5362–5364.
- [4] For selected examples of cyclopropanation: a) A. Nakamura, A. Konishi, Y. Tatsuno, S. Otsuka, *J. Am. Chem. Soc.* **1978**, *100*, 3443–3448; b) M. P. Doyle, M. Y. Eismont, Q.-L. Zhou, *Russ. Chem. Bull.* **1997**, *46*, 955–958; c) M. Barberis, J. Pérez-Prieto, S.-E. Stiriba, P. Lahuerta, *Org. Lett.* **2001**, *3*, 3317–3319; d) I. Nicolas, P. L. Maux, G. Simonneaux, *Tetrahedron Lett.* **2008**, *49*, 2111–2113; for selected examples of C–H insertion: e) P. Müller, E. Maîtrejean, *Collect. Czech. Chem. Commun.* **1999**, *64*, 1807–1826; f) D. J. Wardrop, R. E. Forslund, *Tetrahedron Lett.* **2002**, *43*, 737–739; g) C. H. Hwang, Y. H. Chong, S. Y. Song, H. S. Kwak, E. Lee, *Chem. Commun.* **2004**, 816–817; h) D. F. Taber, W. Tian, *J. Org. Chem.* **2007**, *72*, 3207–3210; i) D. F. Taber, W. Tian, *J. Org. Chem.* **2008**, *73*, 7560–7564; j) J. R. Denton, H. M. L. Davies, *Org. Lett.* **2009**, *11*, 787–790; k) D. F. Taber, C. M. Paquette, P.-M. Gu, W. Tian, *J. Org. Chem.* **2013**, *78*, 9772–9780; for a Si–H insertion: l) M. Ge, E. J. Corey, *Tetrahedron Lett.* **2006**, *47*, 2319–2321; for selected examples of ylide-involving tandem reactions: m) S. Kitagaki, M. Anada, O. Kataoka, K. Matsuno, C. Umeda, N. Watanabe, S. Hashimoto, *J. Am. Chem. Soc.* **1999**, *121*, 1417–1418; n) H. Suga, D. Ishimoto, S. Higuchi, M. Ohtsuka, T. Arikawa, T. Tsuchida, A. Kakehi, T. Baba, *Org. Lett.* **2007**, *9*, 4359–4362; o) X.-F. Xu, Y. Qian, L.-P. Yang, W.-H. Hu, *Chem. Commun.* **2011**, *47*, 797–799; p) Y. Qian, C.-C. Jing, S.-Y. Liu, W.-H. Hu, *Chem. Commun.* **2013**, *49*, 2700–2702.
- [5] An N–H insertion of α -diazoketones catalyzed by a chiral iron porphyrin was investigated; however, no chiral induction was obtained. I. Nicolas, T. Roisnel, P. L. Maux, G. Simonneaux, *Tetrahedron Lett.* **2009**, *50*, 5149–5151.
- [6] For a review: a) X. Guo, W.-H. Hu, *Acc. Chem. Res.* **2013**, *46*, 2427–2440; for selected examples: b) H.-X. Huang, Y.-H. Wang, Z.-Y. Chen, W.-H. Hu, *Adv. Synth. Catal.* **2005**, *347*, 531–534;

- c) X. Guo, H.-X. Huang, L.-P. Yang, W.-H. Hu, *Org. Lett.* **2007**, *9*, 4721–4723; d) X. Zhang, J.-J. Ji, Y.-G. Zhu, C.-C. Jing, M. Li, W.-H. Hu, *Org. Biomol. Chem.* **2012**, *10*, 2133–2138.
- [7] a) J. L. Wood, G. A. Moniz, D. A. Pflum, B. M. Stoltz, A. A. Holubec, H.-J. Dietrich, *J. Am. Chem. Soc.* **1999**, *121*, 1748–1749; b) J. L. Wood, G. A. Moniz, *Org. Lett.* **1999**, *1*, 371–374; c) G. A. Moniz, J. L. Wood, *J. Am. Chem. Soc.* **2001**, *123*, 5095–5097; d) Z.-J. Li, H. M. L. Davies, *J. Am. Chem. Soc.* **2010**, *132*, 396–401; e) Z.-J. Li, V. Boyarskikh, J. H. Hansen, J. Autschbach, D. G. Musaev, H. M. L. Davies, *J. Am. Chem. Soc.* **2012**, *134*, 15497–15504.
- [8] For the initial investigation into the enantioselective insertion reactions of α -diazoesters by using chiral proton-shuttle catalysts: a) B. Xu, S.-F. Zhu, X.-L. Xie, J.-J. Shen, Q.-L. Zhou, *Angew. Chem.* **2011**, *123*, 11685–11688; *Angew. Chem. Int. Ed.* **2011**, *50*, 11483–11486; for further applications: b) H. Saito, D. Morita, T. Uchiyama, M. Miyake, S. Miyairi, *Tetrahedron Lett.* **2012**, *53*, 6662–6664; c) H. Qiu, D. Zhang, S.-Y. Liu, L. Qiu, J. Zhou, Y. Qian, C.-W. Zhai, W.-H. Hu, *Acta Chim. Sin.* **2012**, *70*, 2484–2488; d) B. Xu, S.-F. Zhu, Z.-C. Zhang, Z.-X. Yu, Y. Ma, Q.-L. Zhou, *Chem. Sci.* **2014**, DOI: 10.1039/c3sc52807c.
- [9] a) D. J. Ager, I. Prakash, D. R. Schaad, *Chem. Rev.* **1996**, *96*, 835–875; b) M. P. Gamble, G. M. P. Giblin, J. G. Montana, P. O'Brien, T. P. Ockendon, R. J. K. Taylor, *Tetrahedron Lett.* **1996**, *37*, 7457–7460; c) R. Tripathy, M. A. Ator, J. P. Mallamo, *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2315–2319; d) K. F. Foley, N. V. Cozzi, *Drug Dev. Res.* **2003**, *60*, 252–260; e) R. Di Santo, R. Costi, A. Roux, M. Artico, O. Befani, T. Meninno, E. Agostinelli, P. Palmegiani, P. Turini, R. Cirilli, R. Ferretti, B. Gallinella, F. L. Torre, *J. Med. Chem.* **2005**, *48*, 4220–4223; f) F. A. Davis, T. Ramachandar, *Tetrahedron Lett.* **2008**, *49*, 870–872; g) G. Kumaraswamy, A. Pitchaiah, *Tetrahedron* **2011**, *67*, 2536–2541.
- [10] The diazoketones were prepared according to reported procedures: D. F. Taber, R. E. Ruckle, M. J. Hennessy, *J. Org. Chem.* **1986**, *51*, 4077–4078 and ref. [4j].
- [11] The absolute configurations of the products were determined by comparison to reported values; for **3a**, **3j**, **3l**, and **3m**: a) D. A. DiRocco, T. Rovis, *Angew. Chem.* **2012**, *124*, 6006–6008; *Angew. Chem. Int. Ed.* **2012**, *51*, 5904–5906; for **3k**: b) M. K. Ghorai, A. Kumar, D. P. Tiwari, *J. Org. Chem.* **2010**, *75*, 137–151; for **3p**: c) T. Hashimoto, M. Hirose, K. Maruoka, *J. Am. Chem. Soc.* **2008**, *130*, 7556–7557; for **3o**, see Ref. [9g].
- [12] The absolute configuration of **5a** was determined by comparison to reported values: Y. Qin, C.-H. Wang, Z.-Y. Huang, X. Xiao, Y.-Z. Jiang, *J. Org. Chem.* **2004**, *69*, 8533–8536.
- [13] CCDC 980485 (**5b**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.