

Homogeneous Catalysis


Gold-Catalyzed 1,2-Oxoarylations of Nitriles with Pyridine-Derived Oxides**

Somnath Narayan Karad and Rai-Shung Liu*

Abstract: We report the first success in the gold-catalyzed oxoarylations of nitriles with pyridine-derived *N*-oxides using gold carbenes as initiators. These oxoarylations were also achieved satisfactorily in intermolecular three-component oxidations, including diverse alkenyldiazo esters, nitriles, and pyridine-based oxides.

Catalytic oxidations of alkynes with organic oxides emerge as thriving topics in current gold and platinum catalysis.^[1–5] These oxidations lead to 1,2-oxofunctionalization products by two paths. With pyridine-derived *N*-oxides, the oxidations typically generate α -oxo carbenes which can be functionalized with a tethered or external nucleophile.^[2,3] For sulfoxides and some nitrones, the initial intermediates (**I** and **II**, respectively) undergo 3,3-sigmatropic shifts to afford the 1,2-oxoarylation products **2** and **3**, respectively (Scheme 1a).^[4,5] Although nitriles and alkynes are two common triple-bond motifs, catalytic oxidations of nitriles with organic or inorganic oxides still remain a formidable task.^[6,7] Commonly used nitrile oxides are not prepared from direct oxidations of nitriles.^[8] In the presence of an oxidant, nitriles typically undergo either a hydration reaction^[6] or a C–H oxidation with an elimination of the nitrile.^[7] The development of a new strategy to achieve a nitrile oxidation is significant and highly desired in catalysis. Herein, we report the catalytic 1,2-oxoarylations^[9] of nitriles through a three-component coupling reaction (Scheme 1b). Our approach involves the initial generation of gold carbenes to trap the nitrile and generate a gold-containing nitrilium species, which is subsequently oxidized by pyridine-based *N*-oxides to afford *N*-(pyridin-2-yl)amide derivatives in an atom-economic fashion. Related to this work are the stoichiometric reactions between stable nitrilium salts and quinoline *N*-oxides, which produced distinctly different products resulting from a 1,5-shift.^[10] Although similar *N*-(pyridin-2-yl)amides were yielded from the reactions between imidoyl chlorides and pyridine *N*-oxides, these noncatalytic reactions also produced the undesired 3-chloropyridine in an appreciable amount.^[11]

The utility of this synthesis allows a rapid entry to the *N*-(pyridin-2-yl)amide framework, which is found as the core structure in several bioactive molecules, including piroxiocam,



Scheme 1. 1,2-Oxoarylation reactions.




Figure 1. Representative bioactive molecules.


cam,^[12] variolin B,^[13] pirenzepin,^[14] and piketoprofen (Figure 1).^[15]

Table 1 shows the realization of an unprecedented 1,2-oxoarylation of 2-cyanomethyl-1-ethynylbenzene (**4a**) with 8-methyl quinoline *N*-oxide (3 equiv) using various gold catalysts. Our initial trial with AuCl₃ (5 mol %) in hot 1,2-dichloroethane (80 °C, 12 h) resulted in an incomplete conversion, from which the double oxidation product **5a** was isolated in 22 % yield, together with unreacted **4a** (55 %; entry 1). We switched to cationic gold catalysts such as PPh₃AuCl/AgSbF₆, P(*t*Bu)₂(*o*-biphenyl)AuCl/AgSbF₆, and IPrAuCl/AgSbF₆ (IPr = 1,3-bis(diisopropylphenyl)imidazol-2-ylidene), which provided **5a** in 48, 75, and 82 % yield, respectively (entries 2–4). Different counter anions to IPrAuCl/AgX (X = OTf and NTf₂) maintained satisfactory product yields (entries 5 and 6). AgSbF₆ and HNTf₂ were catalytically inactive (entries 7 and 8). We tested the reactions

[*] S. N. Karad, Prof. Dr. R.-S. Liu
Department of Chemistry, National Tsing-Hua University
Hsinchu (30013) (Taiwan)
E-mail: rslu@mx.nthu.edu.tw

[**] We thank the National Science Council, Taiwan, for financial support of this work.

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201403015>.


Table 1: Catalyst screening for 1,2-oxoarylations of nitriles.


Entry	Catalyst	Solvent ^[b]	t [h]	Yield [%] ^[c]	
				4a	5a
1	AuCl ₃	DCE	12	55	22
2	PPh ₃ AuCl/AgSbF ₆	DCE	12	15	48
3	LAuCl/AgSbF ₆ ^[a]	DCE	10	–	75
4	IPrAuCl/AgSbF ₆	DCE	7.5	–	82
5	IPrAuCl/AgOTf	DCE	9	–	80
6	IPrAuCl/AgNTf ₂	DCE	9.5	–	79
7	AgSbF ₆	DCE	9.5	85	–
8	HNTf ₂	DCE	24	80	–
9	IPrAuCl/AgSbF ₆	C ₆ H ₅ Cl	10.5	–	72
10	IPrAuCl/AgSbF ₆	1,4-dioxane	12.5	–	54

[a] L = P(*t*Bu)₂(*o*-biphenyl). [b] [4a] = 0.20 m. [c] Product yields are reported for product isolated after purification using a silica gel column. DCE = 1,2-dichloroethane, IPr = 1,3-bis(diisopropyl phenyl)imidazol-2-ylidene, Tf = trifluoromethanesulfonyl.

in other solvents, such as chlorobenzene and 1,4-dioxane, thus affording **5a** in 72 and 54 % yields, respectively (entries 9 and 10). The molecular structure of **5a** was verified by an X-ray diffraction study,^[16] thus confirming the working mechanism (Scheme 1 b), which involves the initial formation of the α -oxo gold carbene **A**.


Table 2 shows the substrate scope of such 1,2-oxoarylations using various 5-cyano-3-en-1-ynes and pyridine-derived *N*-oxides. The results show the reactions for the benzenoid substrates **4b–d** bearing various 4-phenyl substituents (R¹ = F, Cl and OMe), thus yielding the desired oxoarylation products **5b–d** in 75–81 % with 8-methylquinoline *N*-oxide as an oxidant. The same reactions were extended to the analogous benzenoid substrates **4e–g**, and yielded the desired **5e–g** in 72–85 % yields. For the nonbenzenoid substrates **4h** and **4i**, the corresponding products **5h** and **5i** were obtained in 84 and 62 % yield, respectively. Such oxoarylations were compatible with 2-substituted pyridine *N*-oxides (R' = phenyl, 2-naphthyl, isobutyl; R'' = H), thus affording the oxoarylation products **5j–l** in 31–53 % yields. The presence of the 2-pyridinyl substituents impede its coordination with gold(I) to maintain the acidity of the gold species. Gold-catalyzed reactions between **4h** and 2-substituted pyridine *N*-oxides (R' = phenyl and 2-naphthyl) proceeded smoothly to afford the desired products **5m** and **5n** in 56 and 65 % yield, respectively. We also prepared 2-cyano-1-ethynylbenzene (**4o**) which was recovered in 58 % yield after a longer reaction time (18 h). For the internal alkyne substrates **4p** (R = cyclopropyl) and **4q** (R = phenyl) we obtained the 1,2-diketone species **4p'** (75 %) and **4q'** (77 %), respectively. The reactions were inapplicable to 2-chloro- or bromo-substituted pyridine *N*-oxides, which worked well in the intermolecular three-

Table 2: Gold-catalyzed oxoarylations of 5-cyano-3-en-1-ynes.



[a] *N*-Oxide (3.0 equiv). [4] = 0.20 m. [b] Product yields are reported for product isolated after purification using a silica gel column.

component system (see Table 3) and they failed to oxidize 5-cyano-3-en-1-yne (**4a**).

Although a catalytic 1,2-oxoarylation of nitriles was proven to be feasible, it is imperative to achieve an intermolecular reaction among gold carbenes, nitriles, and organic oxides to access acyclic molecules. The α -oxo metal carbenes **III** were reported to undergo facile [3+2] cycloadditions to form the oxazole products **6** [Eq. (1)].^[17] We thus




sought to use the alkenyldiazo ester **7a** to form a gold alkenyl carbene species to induce a nucleophilic attack of the nitrile at the alkenyl C3. Such regioselectivity avoids the formation of undesired oxazole products.^[18b, 19] The reaction between **7a**, IPrAuSbF₆ (8 mol %), 8-methylquinoline *N*-oxide (1.1 equiv), and CH₃CN (8 equiv) in DCE (28 °C, 6 h, 0.2 m) gave the oxoarylation product **8a** in 80 % yield [Eq. (2)]. Notably, we found no pyrrole derivatives resulting from a [3+2] cycloaddition of the alkenylgold carbene with nitriles.^[18b] The yield was slightly decreased to 74 % when using a smaller amount of acetonitrile (4 equiv).



The scope of this intermolecular process was assessed with various diazo species (**7**), IPrAuSbF₆ (8 mol %), nitriles (4 equiv), and *N*-oxides (1.1 equiv), as shown in Table 3. Notably, the resulting 1,2-oxoarylation products **8c** and **8d** resulted from the addition of benzonitrile and styrylnitrile, respectively, which reacted with the vinylgold carbene intermediates, and is consistent with the reported regiochemistry in the literature.^[18b,19,20] The results also show the applicability of this oxoarylation to *n*-butynitrile to afford the desired product **8b** in satisfactory yield. These gold-catalyzed reactions were amenable to halo-containing pyridine *N*-oxides, including 2-bromo, 2-chloro, and 2,4-dichloro substrates, thus yielding desired products **8e–g** in 42–71 % yields. The less basic pyridines tended to bind to gold(I) weakly to maintain the acidity of the gold complex. For 2-phenyl- and 2-(2-naphthyl)pyridine *N*-oxides, the corresponding products **8h,i** were obtained in reasonable yields (62–68 %). We prepared the 2-substituted alkenyldiazo esters **7b** (*R*¹=Me, *R*²=H) and **7c** (*R*¹=Ph, *R*²=H), both of which provided desired the 1,2-


oxoarylation products **8j** and **8k**, respectively, in 58–71 % yield. The reactions were also suitable for the 3-substituted alkenyldiazo ester **7d** (*R*¹=H, *R*²=Et), which reacted with 8-methylquinoline *N*-oxide and 2-naphthylpyridine *N*-oxide to afford desired compounds **8l** (68 %) and **8m** (65 %), respectively.

We postulate a mechanism to rationalize the 1,2-oxoarylations of 2-cyanomethyl-1-ethynylbenzene (**4a**) with pyridine-based *N*-oxides, as depicted in Scheme 2. In the presence



Scheme 2. Proposed mechanism for 1,2-oxoarylations of nitriles.

Table 3: Gold-catalyzed intermolecular 1,2-oxoarylations of nitriles.



[a] [**7**] = 0.20 M, nitrile (4 equiv), *N*-oxide (1.1 equiv). [b] Product yields are reported for product isolated after purification using a neutral alumina column.

of IPrAuSbF₆, this mixture is expected to generate the α -oxo gold carbene **A** which is attacked by a tethered nitrile to form the seven-membered nitrilium species **B**. The species **B** might possess a vinyl cation resonance form (**B'**) to reduce the ring strain. A subsequent attack of the pyridine-based oxide on **B'** (or **B**) is expected to form the adduct **C**, and the pyridinium ring then undergoes attack by the iminyl nitrogen atom to produce the azacyclic species **D**. A ring cleavage of this azacyclic species forms the gold-containing *N*-(pyridin-2-yl)amide **E**, together with a loss of proton. Notably, a 2,3-sigmatropic shift in the transformation **C**→**E** is distinct from the reported 1,5-shift for the reactions of quinoline *N*-oxide with imidoyl chlorides or nitrilium species, and thus yields a pyridine derivative.^[10,21–22] The proton ultimately implements the protodeauration of **E** to afford the observed *N*-(pyridin-2-yl)amides **5a** and **5j–l**. This nitrile oxoarylation failed to work with **4o** (Table 2) because the corresponding six-membered nitrilium intermediate is too strained. This proposed mechanism also rationalizes the 1,2-oxoarylation products **8** resulting from the three-component couplings involving nitriles, alkenyldiazo esters, and pyridine-based oxides as depicted in Table 3 (see details in the Supporting Information).

Although alkynes and nitriles are two common triple-bond species, catalytic oxidations of nitriles with oxidants have no precedent. We report herein the first successful gold-catalyzed oxoarylation of nitriles with pyridine-derived *N*-oxides using gold carbenes as initiators. Initial investigations indicate that a variety of oxoarylation products can be accessed from a number of 5-cyano-3-en-1-yne and pyridine-based *N*-oxides.^[23] Seven-membered cyclic nitriliums were involved as key intermediates. These gold-catalyzed nitrile oxoarylations were also achieved satisfactorily for

intermolecular three-component oxidations, including various alkenyldiazo esters, nitriles, and pyridine-based *N*-oxides. This work reports the first examples of the oxidations of nitriles with organic *N*-oxides. Expansion of this nitrile oxoarylation, which involves distinct nitrilium species,^[21,22] to generate organic and inorganic oxides is under current investigation.



Received: March 5, 2014

Published online: April 22, 2014

Keywords: carbenes · gold · heterocycles · nitrogen oxides · synthetic methods

- [1] Review: a) J. Xiao, X. Li, *Angew. Chem.* **2011**, *123*, 7364–7375; *Angew. Chem. Int. Ed.* **2011**, *50*, 7226–7236; b) L. Zhang, *Acc. Chem. Res.* **2014**, *47*, 877–888.
- [2] a) L. Ye, L. Cui, G. Zhang, L. Zhang, *J. Am. Chem. Soc.* **2010**, *132*, 3258–3259; b) L. Ye, W. He, L. Zhang, *J. Am. Chem. Soc.* **2010**, *132*, 8550–8551; c) L. Ye, W. He, L. Zhang, *Angew. Chem.* **2011**, *123*, 3294–3297; *Angew. Chem. Int. Ed.* **2011**, *50*, 3236–3239; d) S. Shi, T. Wang, W. Yang, M. Rudolph, A. S. K. Hashmi, *Chem. Eur. J.* **2013**, *19*, 6576–6580; e) A. S. K. Hashmi, T. Wang, S. Shi, M. Rudolph, *J. Org. Chem.* **2012**, *77*, 7761–7767.
- [3] a) B. Lu, C. Li, L. Zhang, *J. Am. Chem. Soc.* **2010**, *132*, 14070–14072; b) W. He, C. Li, L. Zhang, *J. Am. Chem. Soc.* **2011**, *133*, 8482–8485; c) Y. Luo, K. Ji, Y. Li, L. Zhang, *J. Am. Chem. Soc.* **2012**, *134*, 17412–17415; d) S. Ghorpade, M.-D. Su, R.-S. Liu, *Angew. Chem.* **2013**, *125*, 4323–4328; *Angew. Chem. Int. Ed.* **2013**, *52*, 4229–4234; e) S. Bhunia, S. Ghorpade, D. B. Huplé, R.-S. Liu, *Angew. Chem.* **2012**, *124*, 2993–2996; *Angew. Chem. Int. Ed.* **2012**, *51*, 2939–2942; f) D. Vasu, H.-H. Hung, S. Bhunia, S. A. Gawade, A. Das, R.-S. Liu, *Angew. Chem.* **2011**, *123*, 7043–7046; *Angew. Chem. Int. Ed.* **2011**, *50*, 6911–6914; g) L. Wang, X. Xie, Y. Liu, *Angew. Chem.* **2013**, *125*, 13544–13548; *Angew. Chem. Int. Ed.* **2013**, *52*, 13302–13306; h) J. Fu, H. Shang, Z. Wang, L. Chang, W. Shao, Z. Yang, Y. Tang, *Angew. Chem.* **2013**, *125*, 4292–4296; *Angew. Chem. Int. Ed.* **2013**, *52*, 4198–4202; i) G. Henrion, T. E. J. Chavas, X. L. Goff, F. Gagosz, *Angew. Chem.* **2013**, *125*, 6397–6402; *Angew. Chem. Int. Ed.* **2013**, *52*, 6277–6282; j) S. K. Pawar, C.-D. Wang, S. Bhunia, A. M. Jadhav, R.-S. Liu, *Angew. Chem.* **2013**, *125*, 7707–7711; *Angew. Chem. Int. Ed.* **2013**, *52*, 7559–7563; k) K. Graf, C. L. Rühl, M. Rudolph, F. Rominger, A. S. K. Hashmi, *Angew. Chem.* **2013**, *125*, 12960–12964; *Angew. Chem. Int. Ed.* **2013**, *52*, 12727–12731; l) P. Nösel, L. N. dos Santos Comprido, T. Lauterbach, M. Rudolph, F. Rominger, A. S. K. Hashmi, *J. Am. Chem. Soc.* **2013**, *135*, 15662–15666.
- [4] a) N. D. Shapiro, F. D. Toste, *J. Am. Chem. Soc.* **2007**, *129*, 4160–4161; b) A. B. Cuenca, S. Montserrat, K. M. Hossain, G. Mancha, A. Lledós, M. Medio-Simón, G. Ujaque, G. Asensio, *Org. Lett.* **2009**, *11*, 4906–4909; c) C.-W. Li, K. Pati, G.-Y. Lin, S. M. Abu Sohel, H.-H. Hung, R.-S. Liu, *Angew. Chem.* **2010**, *122*, 10087–10090; *Angew. Chem. Int. Ed.* **2010**, *49*, 9891–9894; d) B. Lu, Y. Li, Y. Wang, D. H. Aue, Y. Luo, L. M. Zhang, *J. Am. Chem. Soc.* **2013**, *135*, 8512–8524.
- [5] a) A. Mukherjee, R. B. Dateer, R. Chaudhuri, S. Bhunia, S. N. Karad, R.-S. Liu, *J. Am. Chem. Soc.* **2011**, *133*, 15372–15375; b) S. Bhunia, C.-J. Chang, R.-S. Liu, *Org. Lett.* **2012**, *14*, 5522–5525.
- [6] a) R. S. Varma, K. P. Naicker, *Org. Lett.* **1999**, *1*, 189–191; b) L. McMaster, F. B. Langreck, *J. Am. Chem. Soc.* **1917**, *39*, 103–109.
- [7] a) H. G. Aurich, *Tetrahedron Lett.* **1964**, *5*, 657–658; b) M. S. Kharasch, G. Sosnovsky, *Tetrahedron* **1958**, *3*, 97–104; c) S. S. Kulp, M. J. McGee, *J. Org. Chem.* **1983**, *48*, 4097–4098; d) D. S. Watt, *J. Org. Chem.* **1974**, *39*, 2799–2800; e) S. J. Selikson, D. S. Watt, *J. Org. Chem.* **1975**, *40*, 267–268.
- [8] See selected examples: a) K.-C. Liu, B. R. Shelton, R. K. Howe, *J. Org. Chem.* **1980**, *45*, 3916–3918; b) H. Gnichtel, L. Autenrieth, P. Luger, K. Vangehr, *Liebigs Ann. Chem.* **1982**, 1091–1095; c) J. H. Boyer, T. Manimaran, V. T. Ramakrishnan, *J. Chem. Soc. Perkin Trans. 1* **1987**, 2163–2169; d) T. V. Hansen, P. Wu, V. V. Fokin, *J. Org. Chem.* **2005**, *70*, 7761–7764; e) S. Kanemasa, M. Nishiuchi, *Tetrahedron Lett.* **1993**, *34*, 4011–4014.
- [9] Although some 1,2-oxoarylations of nitriles were named in the literature, such reactions actually proceeded by initial hydration of the nitrile. See: S.-K. Xiang, D.-X. Zhang, H. Hu, J.-L. Shi, L.-G. Liao, C. Feng, B.-Q. Wang, K.-Q. Zhao, P. Hu, H. Yang, W.-H. Yu, *Adv. Synth. Catal.* **2013**, *355*, 1495–1499.
- [10] For a 1,5-shift occurring with quinoline *N*-oxide and imidoyl chlorides or nitriliums, see: R. A. Abramovitch, R. B. Rogers, G. M. Singer, *J. Org. Chem.* **1975**, *40*, 41–42.
- [11] a) R. A. Abramovitch, G. M. Singer, *J. Am. Chem. Soc.* **1969**, *91*, 5672–5673; b) R. A. Abramovitch, G. M. Singer, *J. Org. Chem.* **1974**, *39*, 1795–1802; c) R. A. Abramovitch, P. Tomasik, *J. Heterocycl. Chem.* **1975**, *12*, 501–503.
- [12] a) J. G. Lombardino, *J. Med. Chem.* **1981**, *24*, 39–42; b) J. G. Lombardino, E. H. Wiseman, J. Chiani, *J. Med. Chem.* **1973**, *16*, 493–496; c) G. Caron, G. Ermondi, *J. Med. Chem.* **2005**, *48*, 3269–3279.
- [13] a) S. R. Walker, E. J. Carter, B. C. Huff, J. C. Morris, *Chem. Rev.* **2009**, *109*, 3080–3098, and references therein; b) T. C. Leboho, S. F. Van Vuuren, J. P. Michael, C. B. de Koning, *Org. Biomol. Chem.* **2014**, *12*, 307–315.
- [14] a) W. D. Bechtel, J. Mierau, H. Palzer, *Arzneim.-Forsch.* **1986**, *36*, 793–796; b) W. G. Eberlein, W. W. Engel, G. Trummlitz, G. Schmidt, R. Hammer, *J. Med. Chem.* **1988**, *31*, 1169–1174; c) R. Hammer, A. Giachetti, *Trends Pharmacol. Sci.* **1984**, *5*, 18–20; d) B. H. Jaub, *Scand. J. Gastroenterol. Suppl.* **1981**, *6*, 68.
- [15] a) J. C. Miralles, J. M. Negro, F. Sánchez-Gascón, M. G. García, *Alergol. Immunol. Clin.* **2001**, *16*, 105–108; b) J. Rodríguez-Lozano, J. J. Goday Buján, J. Del Pozo, E. Fonseca, *Contact Dermatit.* **2005**, *52*, 110–111.
- [16] CCDC 983208 (**14**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [17] For catalytic [3+2]-cycloadditions between nitriles and α -oxo or α -imino carbene species, see Ref. [3b] and selected examples: a) Y. Xiao, L. Zhang, *Org. Lett.* **2012**, *14*, 4662–4665; b) K. J. Doyle, C. J. Moody, *Tetrahedron* **1994**, *50*, 3761–3772; c) T. Ibata, K. Fukushima, *Chem. Lett.* **1992**, 2197–2200.
- [18] For synthesis of pyrrole compounds from the [3+2] cycloadditions of nitriles and alkenylmetal carbenes, see: a) R. J. Billèdeau, K. R. Klein, D. Kaplan, Y. Lou, *Org. Lett.* **2013**, *15*, 1421–1423; b) G. Lonzi, L. A. López, *Adv. Synth. Catal.* **2013**, *355*, 1948–1954.
- [19] For a nucleophilic addition at C3 of alkenylgold carbenes, see: Ref. [16b] and a) V. V. Pagar, A. M. Jadhav, R.-S. Liu, *J. Am. Chem. Soc.* **2011**, *133*, 20728–20731; b) J. Barluenga, G. Lonzi, M. Tomás, L. A. López, *Chem. Eur. J.* **2013**, *19*, 1573–1576.
- [20] For the same regioselectivity of alkenylrhodium carbenes, see selected examples: a) X. Xu, P. Y. Zavalij, M. P. Doyle, *Angew. Chem.* **2013**, *125*, 12896–12900; *Angew. Chem. Int. Ed.* **2013**, *52*, 12664–12668; b) X. Wang, X. Xu, P. Y. Zavalij, M. P. Doyle, *J. Am. Chem. Soc.* **2011**, *133*, 16402–16405; c) X. Xu, P. Y. Zavalij, M. P. Doyle, *Angew. Chem.* **2012**, *124*, 9967–9971; *Angew. Chem. Int. Ed.* **2012**, *51*, 9829–9833; d) D. Valette, Y. Lian, J. P. Haydek, K. I. Hardcastle, H. M. L. Davies, *Angew. Chem.* **2012**, *124*, 8764–8767; *Angew. Chem. Int. Ed.* **2012**, *51*, 8636–8639; e) Y. Lian, H. M. L. Davies, *Org. Lett.* **2012**, *14*, 1934–1937.

[21] We attempted to realize an oxoarylation reaction with the nitrilium intermediates **F** generated from diphenylmethanol and acetonitrile (10 equiv) according to Hashmi's report^[22] [Eq. (3)]. We obtained the amide **10** and ether **11** in 53 and 36% yield, respectively, without any oxidant. In the presence of 8-methylquinoline *N*-oxide (1.2 equiv), we obtained no oxoarylation product in tractable amount [Eq. (4)] whereas species **10** and **11** were still formed. Accordingly, not every nitrilium species can be used in the oxoarylation reaction.



[22] N. Ibrahim, A. S. K. Hashmi, F. Rominger, *Adv. Synth. Catal.* **2011**, 353, 461–468.

[23] Unsubstituted pyridine and quinoline *N*-oxides were inapplicable substrates for the reaction in Tables 2–3 because the released pyridine and quinoline were expected to poison gold catalyst.