

# NUTRITION IN THE PERIOPERATIVE PATIENT

Lyn Howard and Christopher Ashley

*Division of Gastroenterology and Clinical Nutrition, Department of Medicine, Albany Medical College, 47 New Scotland Avenue, Albany, New York 12208;  
email: harrinc@mail.amc.edu*

**Key Words** parenteral, enteral, preoperative, postoperative, surgery

■ **Abstract** The association of malnutrition with surgical morbidity and mortality is well recognized. The question of whether this relationship is causal or simply an association in sick patients has been hotly debated. The field of nutrition support has grown out of the belief that correcting malnutrition will modify associated risks for poor outcome. It has been easier to substantiate this belief in some clinical situations than in others. The evidence for nutrition support during the perioperative period is reviewed and recommendations are made about where nutrition support is most useful and where it may be counterproductive. Some of the important unanswered questions about perioperative nutrition support are raised.

## CONTENTS

Introduction .....	263
Preoperative Nutrition Support .....	264
Postoperative Nutrition Support .....	268
Cost of Perioperative Nutrition .....	271
Unresolved Issues in Perioperative Nutrition .....	273

## Introduction

Why is nutrition in the surgical patient such an important issue? The reason is simple: All surgeons know that a nutritionally depleted patient implies a much higher likelihood of postoperative morbidity and mortality. This association was first pointed out by Studley in 1936 (141). Chronic peptic ulcer patients undergoing gastrectomy had a 3.5% mortality rate if they had less than 20% preoperative weight loss, but a 33% mortality rate if they had lost more than 20%. This relationship between preoperative malnutrition and poor outcome has subsequently been confirmed in many other studies (26, 43, 39, 40) and received special emphasis by the Health and Public Policy Committee of the American College of Physicians (64). While the adequately nourished patient usually tolerates major surgery well with rapid recovery of gastrointestinal function, the severely malnourished patient may develop anastomotic breakdown, infection, and may even deteriorate into multiorgan failure.

The impact of malnutrition on wound healing depends in part on the type of healing required. A wound that heals by primary intention, such as a bowel anastomosis, will often heal even in an emaciated patient so long as the healing process is not complicated by infection. Such wounds appear to have metabolic priority for one to two weeks post surgery. In contrast, a wound that heals by secondary intention, such as a decubitus ulcer or grafted burn, strongly reflects the patient's nutritional status. Even from the outset the malnourished patient's wound exhibits reduced or absent granulation.

If weight loss and hypoalbuminemia increase surgical morbidity and mortality, it would seem logical that providing perioperative specialized nutrition support (SNS) would reduce these complications. However, SNS carries its own risk of complications, for example sepsis if the SNS is parenteral and aspiration pneumonia if the SNS is tube enteral. These risks and the cost of SNS make it imperative that risk-benefit and cost-benefit is determined for parenteral and enteral SNS in different perioperative settings.

Parenteral SNS has been more extensively studied than tube enteral SNS. The reasons for this may be the early surgical enthusiasm for parenteral therapy, the fact that delivering enteral SNS can be more challenging, and the recognition that septic complications and the higher cost of parenteral SNS make randomized outcome studies more urgent. It should also be noted that the majority of surgical patients included in SNS trials had some form of cancer.

In theory the goals of perioperative SNS are first to restore the nutritional status of a depleted patient preoperatively, so operative risks are reduced, then to support the depleted patient through the catabolic phase induced by surgery, and finally to speed the healing process and the return of gastrointestinal function so normal oral food intake can be restored as soon as possible. What is the evidence that SNS does this and who are the patients that benefit from SNS intervention?

## Preoperative Nutrition Support

Clearly preoperative SNS can only be considered in patients who do not require urgent surgical intervention.

**PREOPERATIVE PARENTERAL NUTRITION** This intervention is usually continued postoperatively, and in many studies, the control group also receives postoperative parenteral nutrition.

In the past four years there have been four meta-analyses examining prospective randomized controlled trials (PRCTs) of parenteral nutrition in perioperative patients (90, 146, 68, 93). As expected, there was considerable overlap in the trials evaluated in these meta-analyses, but because inclusion criteria and publication dates differed, there was some variation in the pooled data analyzed, as shown in Table 1. Two of the analyses treated pre- and postoperative studies separately (90, 146), while the other two treated them together and also included postoperative studies using hypocaloric parenteral nutrition, so-called protein sparing therapy (68, 93).

**TABLE 1** Meta-analyses of perioperative parenteral nutrition, number of constituent studies (#), and pooled sample size (n)

STUDY	Data analysis			
	Preoperative		Postoperative	
	#	(n)	#	(n)
Klein et al. 1997 (90)	13 <sup>a</sup>	(1358)	9 <sup>b</sup>	(754)
Torosian 1999 (146)	13 <sup>c</sup>	(1245)	8 <sup>d</sup>	(710)
Heyland et al. 2001b (68)	11 <sup>e</sup>	(1165)	16 <sup>f</sup>	(1742) [6 <sup>g</sup> (899) ProtSpar]
Koretz et al. 2001 (93)	25 <sup>h</sup>	(2164)	18 <sup>i</sup>	(482) [13 (1033) ProtSpar]

The PRCTs included in these meta-analyses were:

<sup>a</sup>11, 12, 49, 50, 65, 106, 108, 112, 113, 138, 145, 149, 150

<sup>b</sup>2, 23, 33, 73, 86, 121, 123, 131, 155

<sup>c</sup>11, 12, 49, 50, 65, 106, 108, 112, 113, 138, 145, 149, 150

<sup>d</sup>23, 33, 73, 86, 121, 123, 131, 155

<sup>e</sup>11, 49, 50, 73, 106, 112, 113, 138, 145, 149, 150

<sup>f</sup>2, 6, 23, 46, 51, 54, 57, 60, 72, 78, 79, 87, 123, 131, 155, 157

<sup>g</sup>46, 51, 54, 57, 79, 87

<sup>h</sup>11, 12, 13, 25, 27, 35, 47, 49, 50, 71, 73, 84, 85, 86, 88, 106, 112, 113, 114, 116, 138, 145, 149, 150, 152

<sup>i</sup>2, 14, 23, 28, 29, 60, 62, 78, 79, 98, 100, 104, 119, 123, 125, 137, 156, 157

\*Stated pooled sample size based only on study samples which were primarily verified. True pooled sample sizes are at least this large.

Most of the patients studied had moderate (10 to 20% weight loss, serum albumin <3.2 to >2.5 g/dl) or severe (>20% weight loss, serum albumin <2.5 g/dl) malnutrition. The duration of the preoperative parenteral nutrition varied from 5 to 14 days; for the majority the intervention lasted at least one week. The end points used were postoperative mortality and complications such as sepsis, mechanical catheter problems, anastomotic leaks, pneumonia, and other major surgical problems. Some studies also measured length of hospital stay.

The pooled results from all four meta-analyses showed no overall effect of preoperative parenteral nutrition on reducing surgical mortality.

Three of the four meta-analyses found a reduction in postoperative complications (90, 146, 68). Serious complications were reduced by 10%, from 40% in the control population to 30% in the preoperative parenteral nutrition patients. The fourth more global meta-analysis, which included several studies not published in English, found only a trend toward reduced postoperative complications (93).

The largest single trial of preoperative parenteral nutrition was the cooperative Veterans Administration (VA) study of 395 patients (149). The patients all had some evidence of malnutrition and underwent major abdominal surgery or noncardio thoracic surgical procedures. In the group as a whole preoperative

parenteral nutrition did not reduce postoperative complications (25.5% parenteral nutrition group, 24.6% controls). There were more infectious complications in the parenteral nutrition group (14.1% parenteral nutrition group, 6.4% controls). This increased rate of infection occurred chiefly in patients with only mild malnutrition. In the severely malnourished parenteral nutrition patients, there were fewer noninfectious complications (5% parenteral nutrition group, 43% controls). The noninfectious complications were anastomotic leaks, broncho-pulmonary fistulas, and organ failure. These results led the authors to recommend preoperative parenteral nutrition only in severely malnourished patients, or about 5% of elective surgical patients. Parenteral nutrition clearly created more risk than benefit in mildly or moderately malnourished patients.

It should be noted that two preoperative parenteral nutrition studies, which provided parenteral nutrition therapy for only three days (73) and five days (145) before surgery, found no reduction in postoperative complications, even though the patients studied were severely malnourished. This suggests preoperative parenteral nutrition requires a longer period than three to five days to induce a clinical benefit. All of the studies in severely malnourished patients that did show benefit provided preoperative parenteral nutrition for seven days or more.

**PREOPERATIVE ENTERAL NUTRITION** Although there have been fewer preoperative enteral nutrition studies, the results to date have been fairly encouraging. In the earliest report, from India (136), 110 malnourished patients awaiting surgery were randomized to 10 days of nasogastric feeding or the routine hospital diet. The patients who received the tube feedings showed significant improvement in body weight and serum protein levels compared to controls. Postoperative mortality was reduced 50% in the treated group (6.0% enteral nutrition group, 11.7% controls) and wound infections were reduced 75% (10.4% enteral nutrition group, 37.2% controls). Foschi et al. (53) randomized preoperative patients with obstructive jaundice undergoing transhepatic biliary drainage to 20 days of SNS (86% by the enteral route) or the routine hospital diet. Postoperative mortality was reduced 75% in the treated group (3.5% enteral nutrition group, 12.5% controls). Flynn & Leighty (52) randomized outpatients with squamous cell carcinoma of the head and neck to routine diet with a nocturnal supplement for 10 to 20 days prior to surgery or the routine diet alone. Postoperative complications were reduced 50% in the supplemented group (32% enteral nutrition group, 59% controls). Von Meyenfildt et al. (150) randomized 151 patients with gastric or colorectal cancer to 10 days of either preoperative parenteral nutrition ( $n = 51$ ) or preoperative enteral nutrition ( $n = 50$ ). Fifty controls went to surgery without delay. Stratification for weight loss allowed for subset analysis. In the severely depleted patients, there was a significant decrease of intra-abdominal abscesses in both intervention groups compared to the depleted controls.

Le Cornu et al. (95) randomized 82 malnourished patients awaiting liver transplant to a treatment group who received a high-calorie oral supplement in addition to their usual diet and a control group who just had their usual diet. In this study

supplementation did not affect operative outcome. However, these investigators had carefully recorded dietary intake, and they showed a higher caloric intake from the usual diet in the controls. As a result, despite the 750 kcal, 20g protein supplement taken by the treatment group, their daily intake was only  $2419 \pm 157$  kcal/d and  $79.8 \pm 5.99$  g protein/d compared to the control group of  $2234 \pm 194$  kcal/day and  $86.5 \pm 6.22$  g protein/d. This very modest difference in calories may well explain the lack of effect of the supplement on surgical outcome. Indeed, this study underlines two important issues in regard to specialized nutrition interventions. First, the parenteral or enteral supplement may have an effect on regular dietary intake, and this must be assessed, as it was in the Le Cornu study. Second, it demonstrates the difficulty of providing artificial enteral nutrition. These hepatic failure patients were at high risk for esophageal/gastric varices and ascites, and there would be strong clinical reluctance to insert a nasogastric or percutaneous feeding tube. Unfortunately, an oral supplement, especially in the outpatient setting, is subject to many compliance issues, such as poor appetite, taste barriers, or cost factors if there is no insurance coverage.

The phenomenon of bacterial translocation from the bowel lumen to regional lymph nodes and ultimately the bloodstream has been firmly established in experimental animal models such as the laboratory rat. The translocation was thought to occur because of an increase in mucosal permeability of the poorly nourished or unused bowel. It was postulated that translocated bacteria or their toxins induced a systemic cytokine response that, if severe, could initiate the multiorgan failure syndrome (31, 101). This mechanism has not been entirely confirmed in humans. First, intraoperative studies in patients with inflamed bowel have not consistently cultured luminal bacteria in regional lymph nodes or the bloodstream (89, 144). Second, intestinal permeability, measured by the lactulose/mannitol test, is not altered by the route of feeding, parenteral or enteral (124). Despite this, the concept that dysfunction of the gastrointestinal barrier allows toxins to enter the body and induce a cytokine response and multiple organ failure remains active (99). This has led to a number of trials using enteral immune-enhancing diets.

Gianotti et al. studied the effect of an enteral immune-enhancing diet providing arginine, omega 3 fatty acid, and nucleotides on the nutritional status and inflammatory response of patients with gastric and colorectal cancer (58). Fifty-eight patients were randomized to receiving 1 L/d of the test diet or 1 L/d of a standard isocaloric, isonitrogenous formula. The diet was given orally for seven days preoperatively and continued eight days postoperatively, via a nasogastric tube. The test diet was more effective than the control diet at sustaining normal nutritional parameters through surgery and damping the inflammatory response of interleukin-6 (IL-6) and IL-1R11 to surgery. In a later phase, this same group showed the immune-enhancing diet translated into better clinical outcome with fewer operative complications and a shorter length of hospital stay (18). These benefits were most pronounced when the enriched formula was given both pre- and postoperatively. A similar clinical benefit was described by Senkal

et al. in 154 patients with upper gastrointestinal malignancies using the same immune-enhancing enteral diet (133).

One problem with this immune-enhancing diet is that several immunonutrients are present in the formula, and hence it is not clear if a particular nutrient confers the biological benefit. A recent study from Amsterdam (148) used a standard pre- and postoperative diet with and without added arginine. The arginine did not significantly improve nutritional or clinical outcome, which suggests that arginine may not be the critical factor.

Overall it is reasonable to conclude from these studies that preoperative enteral nutrition is of benefit in malnourished patients and the benefit may be amplified by the use of immune-enhancing diets. Once again the stumbling block is how to effectively deliver the enteral support preoperatively. In many clinical situations there is reluctance to subject the patient to a nasoenteric or percutaneous tube in advance of surgery. Furthermore, the most cost-saving approach would be to initiate preoperative enteral support at home, prior to surgical admission. But insurance carriers may not be willing to cover such therapy at home. For example, this is not an allowable cost under Medicare's home parenteral and enteral nutrition prosthetic device benefit (74). As a consequence, prescribing an expensive immune-enhancing formula for preoperative home buildup is not yet a feasible approach. It will take more studies to prove this is truly cost effective before funding will be provided.

## Postoperative Nutrition Support

Postoperative nutrition support, by convention, refers to nutrition support initiated after surgery rather than preoperative nutrition, which is continued postoperatively. Postoperative support is potentially a larger issue than preoperative support since many operations cannot be delayed pending nutritional intervention, and preoperative support, at least administered parenterally, appears indicated only in the 5% of elective surgical patients who are severely malnourished.

**POSTOPERATIVE PARENTERAL NUTRITION** Nine studies involving more than 700 patients have compared parenteral nutrition to simple intravenous fluids. In many more studies parenteral nutrition was compared to early enteral nutrition, or the parenteral nutrition provided was hypocaloric protein-sparing therapy. The meta-analyses of these nine studies showed parenteral nutrition had no effect on mortality, and unfortunately increased morbidity by 10%, mostly through septic complications.

In Sandstrom's study of 300 patients, 60% of the postoperative patients resumed oral nutrition within eight to nine days regardless of whether they received parenteral nutrition or glucose only (131). The remaining 40% had delayed return of gastrointestinal function; those receiving parenteral nutrition were at increased risk for sepsis, and those receiving hypocaloric glucose were at increased risk for wound breakdown and mortality. These results strongly suggest routine

postoperative parenteral nutrition is inappropriate. However, some patients did appear to need nutrition support. Unfortunately, patients with delayed return of gastrointestinal function could not be identified by preoperative criteria. The Sandstrom study was important also because it showed a high risk of mortality and morbidity if postoperative patients receive only hypocaloric feeding for more than 14 days. The safe interval for withholding nutrition support is not precisely known, but withholding for two weeks is evidently too long.

In the two meta-analyses where hypocaloric protein-sparing therapy was evaluated, there was no benefit seen in terms of operative mortality or complication rates (68, 93).

**POSTOPERATIVE ENTERAL NUTRITION** Two studies in elderly women with hip fractures (8, 41) have shown enteral supplementation shortens the time until weight bearing, speeds return of independent mobility, and reduces length of hospital stay. In the most underweight women, there was a trend—but not a statistical difference—toward lower mortality in the supplemented group, 8% versus 22% in the controls. A similar but more recent study (10) randomized 101 malnourished postoperative patients to a treatment group that drank 400 mg of a formula providing 1 kcal/ml and 0.06 g protein/ml. The patients were encouraged to sip this supplement between meals. They were compared to a control group receiving standard care. In the control group, weight declined for two months after discharge. In the treated group only half the amount of weight was lost; this reached a nadir at four weeks, and then the treatment group began regaining. The treated group had fewer infections and required fewer antibiotics. Quality of life was assessed by the short form-36 questionnaire, and the treatment group showed significantly better physical and mental health compared with the control group. This study suggests nutritional status and quality of life are linked, and postoperative impairment can be substantially redressed by a simple, fairly inexpensive oral supplement.

Barium studies (77, 115) have shown that small bowel motility continues in the postoperative period, even though gastroparesis may be present for 24 to 48 hours and colonic paresis for 3 to 5 days. This finding has encouraged surgeons to try early postoperative jejunal feeding of an elemental diet as an alternative to intravenous isotonic glucose or parenteral nutrition. The jejunum is accessed either by a nasojejunal tube or percutaneous needle catheter jejunostomy placed intraoperatively. There are four PRCTs involving 142 patients, most of whom had gastrointestinal or head and neck cancer (80, 128, 130, 139). Although one study of 14 patients had fewer complications in those treated with the jejunal elemental diet compared to those receiving intravenous isotonic glucose (128), pooled data did not confirm a significant clinical or nutritional advantage. Thus routine early jejunal feeding with standard formulas is feasible but not clearly beneficial.

Given the preserved perioperative motility of the small bowel, Moncure et al. have questioned if jejunostomy feeds really need to be stopped eight hours prior to elective surgery, as is the standard of care for all forms of enteral intake (109). They randomized 82 patients undergoing nonabdominal operations to either

**TABLE 2** Meta-analyses of peri-operative enteral nutrition, number of constituent studies (#), and pooled sample size (n)

STUDY	Data analysis			
	Preoperative		Postoperative	
	#	(n)	#	(n)
Klein et al. 1997 (90)	2 <sup>a</sup>	(211)	4 <sup>b</sup>	(142)
Torosian 1999 (146)	2 <sup>c</sup>	(211)	4 <sup>d</sup>	(142)
Heys et al. 1999 (70)			11 <sup>e</sup>	(1009)*
Heyland et al. 2001c (69)	22 <sup>f</sup>	(2419)*	9 <sup>g</sup>	(1130)**
Beale et al. 1999 (9)	12 <sup>h</sup>	(1482)*		

\*Included both surgical and critically ill patients.

\*\*Included only surgical patients.

The PRCTs included in these meta-analyses were:

<sup>a</sup>136, 150

<sup>b</sup>80, 128, 130, 139

<sup>c</sup>136, 150

<sup>d</sup>80, 128, 130, 139

<sup>e</sup>15, 19, 20, 37, 38, 94, 107, 111, 129, 132, 133

<sup>f</sup>7, 15, 18, 21, 24, 32, 37, 38, 48, 56, 58, 59, 94, 107, 111, 127, 132, 133, 134, 140, 151, Ross Prod. Div. Abbott Labs 1996 (unpublished)

<sup>g</sup>18, 21, 37, 38, 58, 132, 133, 134, 140

<sup>h</sup>7, 15, 19, 32, 37, 38, 55, 94, 111, 132, 133, 151

discontinuing their jejunostomy feedings when transported to the operating room or eight hours before surgery. The clinical outcome was the same in both groups of patients; there was no evidence of aspiration. The group fed until the last moment, as expected, received more calories and protein. While this study showed that continuing jejunal feeds until just before surgery is safe, any clinical benefit from the greater nutrient intake was not demonstrated.

A growing number of early postoperative tube enteral PRCTs have led to meta-analyses as summarized in Table 2.

Since the 1980s a number of PRCTs have compared early enteral nutrition to early parenteral nutrition in postoperative and other critically ill patients (trauma, burns, head injury, acute pancreatitis). These studies found enteral nutrition was superior to parenteral nutrition with fewer postoperative infections and shorter hospital stays (110, 111, 3, 94, 16). The better outcome with enteral nutrition was associated with a lower stress response and less endotoxin translocation (143, 142) and better peripheral protein synthesis (63). The initial interpretation of these studies was that enteral feeding was superior because it more effectively preserved the immunological barrier function of the intestinal mucosa. However, this interpretation has recently come under reappraisal (82, 83). It has been pointed out that most



of these comparative trials provided more calories to the parenteral patients, for while the trials were intended to be isocaloric and isonitrogenous, it was difficult to get the enteral patients up to the planned nutrient amounts. As a consequence the parenteral patients had higher glucose and insulin levels, and this metabolic perturbation may explain their greater stress response and higher infection rate. Recent obesity studies have shown that excessive caloric intake is associated with increased expression of tumor necrosis factor receptors and higher IL-6, leading to greater immunosuppression (34). Thus the enteral nutrition/parenteral nutrition comparisons may have to be repeated with greater attention to isocaloric-isonitrogenous feeding. In one recent study of nonsurgical patients, enteral nutrition was found to be more hazardous than parenteral nutrition. In 2001, Woodcock et al. (154) published what they described as a prospective "pragmatic" comparison of enteral and parenteral nutrition in 562 patients. This was not a strictly randomized study in that the patients' attending physician assessed gastrointestinal function and determined the route of nutrition support in 75% of the patients. The remaining 25% were randomized. The study showed enteral nutrition met with more technical difficulties, making it harder to achieve an adequate nutritional intake. More enteral nutrition patients died. There was no difference between the parenteral and enteral groups in terms of septic morbidity.

One more chapter in the story of postoperative enteral nutrition support has been the use of immune-enhancing diets compared to standard enteral formulas. Two meta-analyses have evaluated this issue in postoperative and in critically ill patients. In 11 PRCTs evaluated by Heys, 6 addressed postoperative patients, and in the 22 PRCTs evaluated by Heyland, 9 addressed postoperative patients. The pooled data found immuno-nutrition did not influence postoperative mortality, but it did reduce infectious complications shortening the length of the hospital stay (70, 69). In a recent subgroup analysis of the critically ill patients with ongoing infection and sepsis, Heyland has shown an increased mortality in those receiving immune-enhancing diets, especially diets high in arginine. Heyland has argued that while the immune-stimulating effect of these diets is likely to benefit the immune-suppressed postoperative patient; the critically ill patients with immune overresponsiveness may respond by fatal aggravation of their inflammatory process (69). Other forms of immune therapy are being tried in critically ill patients, notably antioxidants (134), probiotics (81), and thymopentine (17). All these interventions are aimed at damping, not augmenting, the patient's immune response.

## Cost of Perioperative Nutrition

The cost of perioperative nutrition has chiefly been examined in patients receiving parenteral nutrition. Ten reports were published between 1982 and 1990 (5, 45, 44, 61, 120, 43, 103, 122, 126, 147), and they found parenteral nutrition cost an additional \$75 to \$530 per day. Some of these estimates may have included the fixed costs of hospitalization (47). In a detailed cost analysis of the Veterans Administration Cooperative Study of perioperative parenteral nutrition, Eisenberg et al. showed that incremental costs were highest (\$3921) for patients who were

least likely to benefit from the therapy because they were only mildly or moderately malnourished. Such patients suffered increased infectious complications from parenteral nutrition and were housed in the expensive intensive care unit because of their parenteral therapy. The incremental costs were lowest (\$3071) for the high-risk severely malnourished patients. In this cohort infectious complications were not increased and noninfectious complications were decreased. The cost savings from avoiding different noninfectious complications were not analyzed because of the small number of patients with any particular complication, but it seemed likely that a large fraction of these costs were already included by the altered length of stay. This study could not show a cost saving from parenteral nutrition for any subgroup of patients. The incremental cost of perioperative parenteral nutrition in the VA study was \$196 per day in 1992. This in-depth analysis has potential for helping the VA system and other hospitals spend perioperative nutrition dollars more rationally.

Enteral nutrition has always been thought of as less expensive than parenteral nutrition, both in hospital and at home. Indeed, there is a tenfold difference in the Medicare reimbursement for these therapies in the home setting (76) (Table 3). In recent years, especially in perioperative patients, there has been a swing toward more post pyloric feeding. A PRCT using radioisotope-labeled enteral feeds has shown that feeding beyond the pylorus significantly reduces gastroesophageal regurgitation with a trend toward less microaspiration (67). Placement of post pyloric tubes can be achieved radiologically, endoscopically, or surgically (153). However, all these approaches require special expertise and usually special fluoroscopy units, which translate into significant cost. In addition, there is a tendency for jejunostomy tubes to flip back into the stomach or block because of their small caliber or simultaneous use for feeding and medications. This frequently requires

**TABLE 3** Medicare allowable charges for HPEN therapy in 1992

	Parenteral (per day)	Tube enteral (per day)
NUTRIENT SOLUTION		
Glucose	\$158–298	\$10–35
Amino acids		
Lipids		\$30–40
Additives	\$7	
Dressing kit	\$7	\$0.5–2
Administration set	\$22	\$11
Pump loan (15mo only)	\$12	\$3.60
Mean* (range)	\$280 (\$238–390)	\$33 (\$25–50)

\*The mean daily Medicare allowable charge was calculated from actual Medicare area reimbursement of 1000 days for Medicare patients receiving HPN and 1000 days for Medicare patients receiving HEN to five large nutrition support programs. This mean daily reimbursement was increased from 80% to 100% to derive the mean daily Medicare allowable charge (75).

jejunostomy tube replacement, which is a further expense. Thus, with modern enteral feeding obtaining and sustaining enteric access is a great deal more expensive than with the simple nasogastric tube (96, 22).

## Unresolved Issues in Perioperative Nutrition

A number of important perioperative nutrition issues await clarification.

More studies are needed to test the feasibility of preoperative enteral nutrition, especially in the outpatient setting. This is a two-part issue. First, is tube access necessary to insure an increased intake? Second, how significant is the benefit in terms of reducing expensive postoperative complications? Since the few PRCTs available so far indicate a reduction in postoperative complications of 50 to 75%, more widespread use of preoperative enteral nutrition in the future may be highly appropriate.

Because postoperative enteral nutrition, especially with immune-enhancing formulas, appears to reduce postoperative complications and length of hospital stay, it is desirable that future studies include a control group receiving routine hospital care (isotonic glucose with gradual resumption of oral diet). This control arm has seldom been included, yet it would help to define when more complex enteral nutrition is justified.

Recent animal studies have demonstrated impaired tolerance of enteral nutrition in partial bowel ischemia (92). Currently there are no good clinical methods for assessing the degree of bowel ischemia in patients with poor splanchnic perfusion. This is clearly another important issue in perioperative patients who have blood pressure instability.

As presented in this overview, preoperative parenteral nutrition reduces non-infectious complications in severely malnourished patients by 10%, but postoperative parenteral nutrition increases complications by 10%. Moreover, 60% of postoperative patients resume oral nutrition in eight to nine days, and in this cohort parenteral nutrition has no beneficial effect (131). It seems likely postoperative parenteral nutrition increases infectious rates, perhaps by inducing hyperglycemia. Under these circumstances, perhaps waiting to start or resume parenteral nutrition for a week could be an important step that deserves testing. Such an intervention might avoid the period of acute surgical stress and the risk of hyperglycemia and septic complications. It might also be possible to avoid postoperative parenteral nutrition altogether in patients who appear close to resuming oral nutrition. If postoperative patients who go for two weeks with only hypocaloric glucose have rising morbidity and mortality, the issue becomes a question of when it is not too early and not too late to intervene with parenteral nutrition. This is clearly an important and unresolved series of questions.

The final issue relates to the fact that the majority of perioperative patients studied had some form of malignancy. Cancer, like AIDS, is a wasting disorder that can lead to marked cachexia. Sometimes the weight loss reflects true starvation and is due to an inability to swallow or absorb nutrients. More often, however, the weight loss reflects altered metabolism due to abnormal hormonal

and cytokine factors. In this form of cachexia there is increased loss of lean body mass, and repletion with nutritional intervention is impaired. Shike et al. showed that parenteral nutrition in patients with small cell lung cancer led to weight gain, but this was due to increased body fat and total body water, not to increased lean body mass (135). In nonsurgical cancer patients, such as those undergoing radiation or chemotherapy, parenteral nutrition has been shown in over 40 PRCTs to provide no clinical benefit. Indeed in two meta-analyses (91, 105) the pooled data revealed evidence of net harm, since the parenteral nutrition-treated group showed a higher incidence of infection. In 1989 this finding led to a cautionary statement by the American College of Physicians advising against routine use of parenteral nutrition in cachectic cancer patients undergoing radiation or chemotherapy (4). Enteral studies in cancer patients undergoing radiation or chemotherapy also demonstrated no therapeutic benefit. Keeping these issues in mind the questions that arise are first, how many of the cancer patients included in the perioperative trials were suffering the metabolic syndrome of cancer cachexia and were therefore unlikely to improve with SNS? Second, might the response to perioperative nutrition be more positive if only noncancer patients were studied? These questions need to be answered so these expensive but potentially valuable SNS therapies can be used as safely and cost effectively as possible.

**The Annual Review of Nutrition is online at <http://nutr.annualreviews.org>**

## LITERATURE CITED

1. Abbott WC, Bistrian BR, Blackburn GL. 1986. The effect of dextrose and amino acids on respiratory function and energy expenditure in morbidly obese patients following gastric bypass surgery. *J. Surg. Res.* 41:225–35
2. Abel RM, Fisher JE, Buckley MJ, Barnett GO, Austen WG. 1976. Malnutrition in cardiac surgical patients: results of a prospective randomized evaluation of early postoperative parenteral nutrition. *Arch. Surg.* 111:45–50
3. Adams S, Dellinger EP, Wortz MJ, Oreskovich MR, Simonovitz D, et al. 1986. Enteral versus parenteral nutritional support following laparotomy for trauma: a randomized prospective trial. *J. Trauma* 26(10):882–91
4. American College of Physicians. 1989. Position paper: parenteral nutrition in patients receiving cancer chemotherapy: a meta-analysis. *Ann. Int. Med.* 110(9):734–36
5. Anderson GF, Steinberg EP. 1986. DRGs and specialized nutrition, prospective payment and nutritional support: the need for reform. *J. Parent. Enteral Nutr.* 10:3–8
6. Askanazi J, Hensle TW, Starker PM, Lockhart SH, LaSala PA, et al. 1986. Effect of immediate postoperative nutritional support on length of hospitalization. *Ann. Surg.* 203:236–39
7. Atkinson S, Sieffert E, Bihari D. 1998. A prospective randomized double-blind clinical trial of enteral immunonutrition in the critically ill. *Crit. Care Med.* 26:1164–72
8. Bastow MD, Rawlings J, Allison SP. 1983. Benefits of supplementary tube feeding after fractured neck of femur: a randomized controlled trial. *Br. Med. J.* 287(6405):1589–92

9. Beale RJ, Bryg DJ, Bihari DJ. 1999. Immunonutrition in the critically ill: a systematic review of clinical outcome. *Crit. Care Med.* 27(12):2799–805
10. Beattie AH, Prach AT, Baxter JP, Pennington CR. 2000. A randomized controlled trial evaluating the use of enteral nutritional supplements postoperatively in malnourished surgical patients. *Gut* 46(6):813–18
11. Bellantone R, Doglietto GB, Bossola M, Pacelli F, Negro F, et al. 1988. Preoperative parenteral nutrition in the high-risk surgical patient. *J. Parent. Enteral Nutr.* 12:195–97
12. Bellantone R, Doglietto GB, Bossola M, Pacelli F, Negro F, et al. 1988. Preoperative parenteral nutrition of malnourished patients. *Acta Chir. Scand.* 22:249–51
13. Bellantone R, Doglietto GB, Bossola M, Pacelli F, Negro F, et al. 1990. Preoperative parenteral nutrition in malnourished high-risk surgical patients. *Nutrition* 6:168–70
14. Bonau RA, Jeevanandam M, Daly JM. 1984. High branched-chain amino acid solutions: relationship of composition to efficacy. *J. Parent. Enteral Nutr.* 8:622–27
15. Bower RH, Cerra FB, Bershadsky B, Licari JJ, Hoyt DB, et al. 1995. Early enteral administration of a formula (Impact) supplemented with arginine, nucleotides, and fish oil in intensive care unit patients: results of a multicenter, prospective, randomized, clinical trial. *Crit. Care Med.* 23(3):436–49
16. Bozzetti F, Braga M, Gianotti C, Mariani L. 2001. Postoperative enteral versus parenteral nutrition in malnourished patients with gastrointestinal cancer: a randomized multicentre trial. *Lancet* 358:1487–92
17. Braga M, Constantini E, DiFrancesco A, Gianotti L, Baccari P, et al. 1994. Impact of thymopentin on the incidence and severity of postoperative infection: a randomized controlled trial. *Brit. J. Surg.* 81:205–8
18. Braga M, Gianotti L, Radaelli G, Vignali A, Mari G, et al. 1999. Perioperative immunonutrition in patients undergoing cancer surgery: results of a randomized double-blind phase 3 trial. *Arch. Surg.* 134(4):428–33
19. Braga M, Gianotti L, Vignali A, Cestari A, Bisagni P, et al. 1998. Artificial nutrition after major abdominal surgery: impact of route of administration and composition of the diet. *Crit. Care Med.* 26:24–30
20. Braga M, Vignali A, Gianotti L, Cestari A, Profili M, et al. 1995. Benefits of early postoperative enteral feeding in cancer patients. *Infusionsher. Transfusionsmed.* 22:280–84
21. Braga M, Vignali A, Gianotti L, Cestari A, Profili M, et al. 1996. Immune and nutritional effects of early enteral nutrition after major abdominal operations. *Eur. J. Surg.* 162:105–12
22. Braunschweig CL, Levy P, Sheean PM, Wang X. 2001. Enteral compared with parenteral nutrition: a meta-analysis. *Am. J. Clin. Nutr.* 74(4):534–42
23. Brennan MF, Pisters PW, Posner M, Quesada O, Shike M. 1994. A prospective randomized trial of total parenteral nutrition after major pancreatic resection for malignancy. *Ann. Surg.* 220:436–41
24. Brown RO, Hunt H, Mowatt-Larssen CA, Wojtysiak SL, Henningfield MF, et al. 1994. Comparison of specialized and standard enteral formulas in trauma patients. *Pharmacotherapy* 14:314–20
25. Buzby GP, Knox LS, Crosby LO, Eisenberg JM, Haakenson CM, et al. 1988. Study protocol: a randomized clinical trial of total parenteral nutrition in malnourished surgical patients. *Am. J. Clin. Nutr.* 47:366–81
26. Buzby GP, Mullen JL, Matthews DC, Hobbs CL, Rosato EF. 1980. Prognostic nutritional index in gastrointestinal surgery. *Am. J. Surg.* 139:160–67
27. Buzby GP, Williford WO, Peterson OL, Crosby LO, Page CP, et al. 1988. A randomized clinical trial of total parenteral

- nutrition in malnourished surgical patients: the rationale and impact of previous clinical trials and pilot study on protocol design. *Am. J. Clin. Nutr.* 47:357–65
28. Capella G, Ruiz JM, Hidalgo L, Alonso M, Salvador R, et al. 1990. Valor de la nutrición parenteral en la gastrectomía total por cáncer. Estudio prospectivo. *Nutr. Hosp.* 5:317–21
  29. Cardona D, Del Moral V, Salvador R, Andreu MA, Trias R, et al. 1986. Early postoperative total parenteral nutrition in gastric cancer: a cost-effectiveness study. *J. Clin. Nutr. Gastroenterol.* 1:267–70
  30. Deleted in proof
  31. Carrico CJ. 1993. The elusive pathophysiology of multiple organ failure syndrome. *Ann. Surg.* 218(2):309–10
  32. Cerra FB, Lehman S, Konstantinides N, Konstantinides F, Shronts EP, et al. 1990. Effect of enteral nutrient on in vitro tests of immune function in ICU patients: a preliminary report. *Nutrition* 6(1):84–87
  33. Collins J, Oxby C, Hill G. 1978. Intravenous amino acids and intravenous hyperalimentation as protein-sparing therapy after major surgery: a controlled clinical trial. *Lancet* 1:778–91
  34. Coppack SW. 2001. Pro-inflammatory cytokines and adipose tissue. *Proc. Nutr. Soc.* 60(3):349–56
  35. Cromack D, Moley J, Pass H, Sindelar W, Maher M, et al. 1988. Prospective randomized trial of parenteral nutrition (TPN) compared to ad lib oral nutrition (ON) in patients with upper GI cancer and weight loss undergoing surgical treatment. Presented at Ann. Meet. Assoc. Acad. Surgery (86), 22nd
  36. Culebras-Fernandez JM, de la Hoz Riesco M, Villares García C, Hurtado Fernandez-Llamazares G, Alonos Villalba A. 1987. Improvement of the nutritional condition with hypocaloric peripheral parenteral nutrition (HPPN) in the immediate postoperative period of elective abdominal surgery. *Infusionstherapie* 14:202–8
  37. Daly JM, Lieberman MD, Goldfine J, Shou J, Weintraub FN, et al. 1992. Enteral nutrition with supplemental arginine, RNA, and omega-3 fatty acids in patients after operation: immunologic, metabolic, and clinical outcome. *Surgery* 112:56–67
  38. Daly JM, Weintraub FN, Shou J, Rosato EF, Lucia M. 1995. Enteral nutrition during multimodality therapy in upper gastrointestinal cancer patients. *Ann. Surg.* 221:327–38
  39. Daley J, Khuri SF, Henderson W, Hur K, Gibbs JO, et al. 1997. Risk adjustment of the postoperative morbidity rate for the comparative assessment of the quality of surgical care: results of the National Veterans Affairs Surgical Risk Study. *J. Am. Coll. Surg.* 185(4):328–40
  40. Dempsey DT, Mullen JL, Buzby GP. 1988. The link between nutritional status and clinical outcome: Can nutritional intervention modify it? *Am. J. Clin. Nutr.* 47(2 suppl.):352–56
  41. Delmi M, Rapin CH, Bengoa JM, Delmas PD, Vasey H, et al. 1990. Dietary supplementation in elderly patients with fractured neck of femur. *Lancet* 335(8696):1013–16
  42. Deleted in proof
  43. Detsky AS, Baker JP, O'Rourke K, Johnston N, Whitwell J, et al. 1987. Perioperative nutrition-associated complications for patients undergoing gastrointestinal surgery. *J. Parent. Enteral Nutr.* 11(5):440–46
  44. Detsky AS, Jeejeebhoy KN. 1984. Cost-effectiveness of preoperative parenteral nutrition in patients undergoing major gastro-intestinal surgery. *J. Parent. Enteral Nutr.* 8:632–37
  45. Detsky AS, Mendelson RA, Baker JP, Jeejeebhoy KN. 1984. The choice to treat all, some. Or no patients undergoing gastrointestinal surgery with nutritional support: a decision analysis approach. *J. Parent. Enteral Nutr.* 8:245–53
  46. Doglietto GB, Gallitelli L, Pacelli F, Bellantone R, Malerba M, et al. 1996. Protein-sparing therapy after major abdominal

- surgery: lack of clinical effects. *Ann. Surg.* 223:357–62
47. Eisenberg JM, Glick HA, Buzby GP, Kinoshian B, Williford WO. 1993. Does perioperative total parenteral nutrition reduce medical care costs? *J. Parent. Enteral Nutr.* 17:201–9
48. Engel JM, Menges T, Neuhauser C, Schaefer B, Hempelmann G. 1997. Effects of various feeding regimens in multiple trauma patients on septic complications and immune parameters. *Anesthesiol. Intensivmed. Notfallmed. Schmerzther.* 32: 234–39
49. Fan ST, Lau WY, Wong KK, Chan YPM. 1989. Pre-operative parenteral nutrition in patients with oesophageal cancer: a prospective, randomized clinical trial. *Clin. Nutr.* 8:23–27
50. Fan ST, Lo CM, Lai EC, Chu KM, Liu CL, et al. 1994. Perioperative nutritional support in patients undergoing hepatectomy for hepatocellular carcinoma. *N. Engl. J. Med.* 331:1547–52
51. Figueras J, Puig P, Rafecas A, Bianchi A, Hernandez F, et al. 1988. Postoperative hypocaloric parenteral nutrition. A study in patients without neoplasm. *Acta Chir. Scand.* 154:435–38
52. Flynn MB, Leighty FF. 1987. Preoperative outpatient nutritional support of patients with squamous cancer of the upper aerodigestive tract. *Am. J. Surg.* 154(4):359–62
53. Foschi D, Cavagna G, Callioni F, Morandi E, Rovati V. 1986. Hyperalimentation of jaundiced patients on percutaneous transhepatic biliary drainage. *Br. J. Surg.* 73(9):716–19
54. Freund H, Hoover HC, Atamian S, Fischer JE. 1979. Infusion of the branched-chain amino acids in postoperative patients. *Ann. Surg.* 190:18–23
55. Galban C, Celaya S, Marco P. 1998. An immune-enhancing enteral diet reduces mortality and episodes of bacteremia in septic ICU patients. *J. Parent. Enteral Nutr.* 22:S13 (Abstr.)
56. Galban C, Mortejo JC, Mesejo A, Marco P, Celaya S, et al. 2000. An immune-enhancing enteral diet reduces mortality rate and episodes of bacteremia in septic intensive care unit patients. *Crit. Care Med.* 28(3):643–48
57. Garden OJ, Smith A, Harris NW, Shenkin A, Sim AJ, et al. 1983. The effect of iso-tonic amino acid infusions on serum proteins and muscle breakdown following surgery. *Br. J. Surg.* 70:79–82
58. Gianotti L, Braga M, Vignali A, Balzano G, Zerbi A, et al. 1997. Effects of route of delivery and formulation of postoperative nutritional support in patients undergoing major operations for malignant neoplasms. *Arch. Surg.* 132(11):1222–29
59. Gottschlich MM, Jenkins M, Warden GD, Baumer T, Havens P, et al. 1990. Differential effects of three enteral dietary regimens on selected outcome variables in burn patients. *J. Parent. Enteral Nutr.* 14:225–36
60. Gys T, Peeters R, Hubens A. 1990. The value of short-term peripheral parenteral nutrition after colorectal surgery: a comparative study with conventional postoperative intravenous fluid. *Acta Chir. Belg.* 90:234–39
61. Hamaoui E, Lefkowitz R, Olender L, Krasnopolsky-Levine E, Favale M, et al. 1990. Enteral nutrition in the early postoperative period: a new semi-elemental formula versus total parenteral nutrition. *J. Parent. Enteral Nutr.* 14(5):501–7
62. Hansell DA, Davies JW, Shenkin A, Garden OJ, Burns HJG, et al. 1989. The effects of an anabolic steroid and peripherally administered intravenous nutrition in the early postoperative period. *J. Parent. Enteral Nutr.* 13:349–58
63. Harrison LE, Hochwald SN, Heslin MJ, Berman R, Burt M, et al. 1997. Early postoperative enteral nutrition improves peripheral protein kinetics in upper gastrointestinal cancer patients undergoing complete resection: a randomized trial. *J. Parent. Enteral Nutr.* 21(4):202–7

64. Health and Public Policy Committee, American College of Physicians. 1987. Perioperative parenteral nutrition. *Ann. Int. Med.* 107:252–53
65. Heatley RV, Williams RHP, Lewis MH. 1979. Pre-operative intravenous feeding: a controlled trial. *Postgrad. Med. J.* 55: 541–45
66. Hensle TW. 1978. Protein-sparing therapy in cystectomy patients. *J. Urol.* 119:355–58
67. Heyland DK, Drover JW, MacDonald S, Novak F, Lam M. 2001. Effect of postpyloric feeding on gastroesophageal regurgitation and pulmonary microaspiration: results of a randomized controlled trial. *Crit. Care Med.* 29(8):1495–501
68. Heyland DK, Montalvo M, MacDonald S, Keefe L, Su XY, et al. 2001. Total parenteral nutrition in the surgical patient: a meta-analysis. *J. Can. Chir.* 44(2):102–11
69. Heyland DK, Novak F, Drover JW, Jain M, Su X, et al. 2001. Should immunonutrition become routine in critically ill patients? A systematic review of the evidence. *JAMA* 286(8):944–53
70. Heys SD, Walker LG, Smith I, Eremin O. 1999. Enteral nutritional supplementation with key nutrients in patients with critical illness and cancer. A meta-analysis of randomized controlled clinical trials. *Ann. Surg.* 229(4):467–77
71. Hoffman E, Hansen BB, Owen-Falkenberg T, Djorup J. 1988. Total parenteral emearing ved colonkirurgi. *Ugeskr. Laeger* 150:1277–79
72. Hogbin BM, Smith AM, Craven AH. 1984. An evaluation of peripheral essential amino acid infusion following major surgery. *J. Parent. Enteral Nutr.* 8:511–14
73. Holter AR, Fischer JE. 1977. The effects of perioperative hyperalimentation on complications in patients with carcinoma and weight loss. *J. Surg. Res.* 23:31–34
74. Howard L. 2000. A global perspective of home parenteral and enteral nutrition. *Nutrition* 16(7-8):625–8
75. Howard L, Ament M, Fleming CR, Shike M, Steiger E. 1995. Current use and clinical outcome of home parenteral and enteral nutrition therapies in the United States. *Gastroenterology* 109:355–65
76. Howard L, Hassan N. 1998. Home parenteral nutrition. *Gastroenterol. Clin. North Am.* 27(2):481–512
77. Hoyer A. 1950. Abdominal distention and intestinal activity following laparotomy. *Acta Radiol.* 83(Suppl):1
78. Hu SS, Fontaine F, Kelly B, Bradford DS. 1998. Nutritional depletion in staged spinal reconstructive surgery. The effect of total parenteral nutrition. *Spine* 23:1401–5
79. Hwang TL, Mou SC, Chen MF. 1993. The importance of a source of sufficient protein in postoperative hypocaloric partial parenteral nutritional support. *J. Parent. Enteral Nutr.* 17:254–56
80. Iovinelli G, Marsili I, Varrassi G. 1993. Nutrition support after total laryngectomy. *J. Parent. Enteral Nutr.* 17:445–48
81. Isolauri E, Sutas Y, Kankaanpää P, Arvilommi H, Salminen S. 2001. Probiotics: effects on immunity. *Am. J. Clin. Nutr.* 73(S):444–50S
82. Jeejeebhoy KN. 2001. Enteral and parenteral nutrition: evidence based approach. *Proc. Nutr. Soc.* 60(3):399–402
83. Jeejeebhoy KN. 2001. TPN: potion or poison? *Am. J. Clin. Nutr.* 74(2):160–63
84. Jensen S. 1982. Parenteral nutrition and cancer surgery. *J. Parent. Enteral Nutr.* 6:335
85. Jensen S, Ginnerup P. 1982. Total parenteral emearing perioperativt til patienter med adenocarcinoma recti. *Ugeskr. Laeger* 144:460–63
86. Jensen S. 1985. Clinical effects of enteral and parenteral nutrition preceding cancer surgery. *Med. Oncol. Tumor Pharmacother.* 2:225–29
87. Jimenez FJ, Leyba CO, Jimenez LM, Valdecasas MS, Montero JG. 1995. Study of hypocaloric peripheral parenteral



- nutrition in postoperative patients. *Clin. Nutr.* 14:88–96
88. Jin D, Phillips M, Byles JE. 1999. Effects of parenteral nutrition support and chemotherapy on the phasic composition of tumor cells in gastrointestinal cancer. *J. Parent. Enteral Nutr.* 23:237–41
89. Kanwar S, Windsor AC, Welsh F, Barclay GR, Guillou PJ, et al. 2000. Lack of correlation between failure of gut barrier function and septic complications after major upper gastrointestinal surgery. *Ann. Surg.* 231(1):88–95
90. Klein S, Kinney J, Jeejeebhoy K, Alpers D, Hellerstein M, et al. 1997. Nutrition support in clinical practice: review of published data and recommendations for future research directions. National Institutes of Health, American Society for Parenteral and Enteral Nutrition and American Society for Clinical Nutrition. *J. Parent. Enteral Nutr.* 21:133–56
91. Klein S, Simes J, Blackburn GL. 1986. Total parenteral nutrition and cancer clinical trials. *Cancer* 58(6):1378–86
92. Kles KA, Wallig MA, Tappenden KA. 2001. Luminal nutrients exacerbate intestinal hypoxia in the hypoperfused jejunum. *J. Parent. Enteral Nutr.* 25(5):246–53
93. Koretz RL, Lipman TO, Klein S. 2001. AGA technical review on parenteral nutrition. *Gastroenterology* 121:970–1001
94. Kudsk DA, Minard G, Croce MA, Brown RO, Lowrey TS, et al. 1996. A randomized trial of isonitrogenous enteral diets after severe trauma. An immune-enhancing diet reduces septic complications. *Ann. Surg.* 224(4):531–40
95. Le Cornu KA, McKiernan FJ, Kapadia SA, Neuberger JM. 2000. A prospective randomized study of preoperative nutritional supplementation in patients awaiting elective orthotopic liver transplantation. *Transplantation* 69(7):1364–69
96. Lipman TO. 1998. Grains or veins: Is enteral nutrition really better than parenteral nutrition? A look at the evidence. *J. Parent. Enteral Nutr.* 22:167–82
97. Lopez-Hellin J, Lopez-Lara M, Mercader S, Gemar E, Garcia-Arumi C, et al. 1997. Early curbing of protein hypercatabolism in postoperative patients by nutritional support with glucose plus amino acids, but not with glucose alone. *Clin. Nutr.* 16:67–73
98. Lorenzoni K. 1981. Postoperative infusiontherapie in der abdominal-chirurgie. *Wien. Med. Wochenschr.* 131:485–91
99. MacFie J, O'Boyle C, Mitchell CJ, Buckley PM, Johnstone D, et al. 1999. Gut origin of sepsis: a prospective study investigating associations between bacterial translocation, gastric microflora, and septic morbidity. *Gut* 45(2):223–28
100. Marin VB, Rebollo MGJ, Castillo-Duran CD, Lopez MTS, Sanabria MS, et al. 1999. Controlled study of early postoperative parenteral nutrition in children. *J. Ped. Surg.* 34:1330–35
101. Marshall JC, Christou NV, Meakins JL. 1993. The gastrointestinal tract. The “undrained abscess” of multiple organ failure. *Ann. Surg.* 218(2):111–19
102. Massar EL, Daly JM, Copeland EM, Johnson DE, Von Eshenbach AC, et al. 1983. Peripheral vein complications in patients receiving amino acid/dextrose solutions. *J. Parent. Enteral Nutr.* 7:159–62
103. Maswosse JJ, Newcomer DR, Quandt CM. 1987. Achieving parenteral nutrition cost savings through prescribing guidelines and formulary restrictions. *Am. J. Hosp. Pharm.* 44:1376–81
104. Mattioli S, Lassari A, Lerro F, DiSimone MP, Pilotti V, et al. 1993. Nutrizione parenterale totale dopo interventi di chirurgia digestiva maggiore. *Chirurgia* 6:129–33
105. McGeer AJ, Detsky AS, O'Rourke K. 1990. Parenteral nutrition in cancer patients undergoing chemotherapy: a meta-analysis. *Nutrition* 6(3):233–40
106. Meguid MM, Curtas MS, Meguid V, Campos AC. 1988. Effects of pre-operative TPN on surgical risk—preliminary status

- report. *Br. J. Clin. Pract.* 42(S63):53–58
107. Mendez C, Jurkovich GJ, Garcia I, Davis D, Parker A, et al. 1997. Effects of an immune-enhancing diet in critically injured patients. *J. Trauma* 42:933–42
  108. Moghissi K, Hornshaw J, Teasdale PR, Dawes EA. 1977. Parenteral nutrition in carcinoma of the oesophagus treated by surgery: nitrogen balance and clinical studies. *Br. J. Surg.* 64:125–28
  109. Moncure M, Samaha E, Moncure K, Mitchell J, Rehm C, et al. 1999. Jejunostomy tube feedings should not be stopped in the perioperative patient. *J. Parent. Enteral Nutr.* 23(6):356–59
  110. Moore FA, Moore EE, Jones TN, McCroskey BL, Peterson VM. 1989. TEN versus TPN following major abdominal trauma-reduced septic morbidity. *J. Trauma* 29(7):916–22
  111. Moore FA, Moore EE, Kudsk KA, Brown RO, Bower RH, et al. 1994. Clinical benefits of an immune-enhancing diet for early postinjury enteral feeding. *J. Trauma* 37(4):607–15
  112. Muller J, Brenner U, Dienst C, Pichlmaier H. 1982. Preoperative parenteral feeding in patients with gastrointestinal carcinoma. *Lancet* 1:68–71
  113. Muller J, Keller H, Brenner U, Walter M, Holtzmuller W. 1986. Indications and effects of preoperative parenteral nutrition. *World J. Surg.* 10:53–63
  114. Muller J, Thul P, Halber M, Huber P. 1988. Parenteral nutrition in tumor patients. *Recent Results Cancer Res.* 108:185–93
  115. Nachlas MM, Younis MT, Roda CP, Wityk JJ. 1972. Gastrointestinal motility studies as a guide to postoperative management. *Ann. Surg.* 175(4):510–22
  116. Neuvonen P, Salo M. 1984. Effects of preoperative parenteral nutrition on cell-mediated immunity in malnourished patients. *Clin. Nutr.* 3:197–201
  117. Neuvonen P, Salo M, Perttila J, Havia T. 1986. Lack of modulation of postoperative immunosuppression by isotonic amino acid infusion. *J. Parent. Enteral Nutr.* 10:160–65
  118. Nissila M, Salo M, Granberg C, Perttila J, Neuvonen P. 1988. Activity of natural killer cells after postoperative amino acid infusion. *J. Parent. Enteral Nutr.* 12:346–50
  119. O'Mahoney JB, McIrvine AJ, Palder SB, See-Young L, Saporoschetz IB, et al. 1984. The effect of short-term postoperative intravenous feeding upon cell-mediated immunity and serum suppressive activity in well-nourished patients. *Surg. Gynecol. Obstet.* 159:27–32
  120. Perry S, Pillar B, Radany MH. 1990. The appropriate use of high-cost, high-risk technologies: the case of total parenteral nutrition. *Qual. Rev. Bull.* 16:214–17
  121. Preshaw R, Attisha R, Hollingworth W. 1979. Randomized sequential trial of parenteral nutrition in healing of colonic anastomoses in man. *Can. J. Surg.* 22:437–39
  122. Reilly JJ, Hull SF, Albert N, Waller A, Bringardener S. 1988. Economic impact of malnutrition: a model for hospitalized patients. *J. Parent. Enteral Nutr.* 12(4):371–76
  123. Reilly J, Mehta R, Teperman L, Cemaj S, Tzakis A, et al. 1990. Nutritional support after liver transplantation: a randomized prospective study. *J. Parent. Enteral Nutr.* 14:386–91
  124. Reynolds JV, Kanwar S, Welsh FK, Windsor AC, Murchan P, et al. 1997. Does the route of feeding modify gut barrier function and clinical outcome in patients after major upper gastrointestinal surgery? *J. Parent. Enteral Nutr.* 21(4):196–201
  125. Rimbaud V, Cardona D, Escudero JR, Artega R, Mestress JM, et al. 1989. Efecto de la nutricion parenteral periferica en el postoperatorio inmediato de la cirugia del sector aortico. *Angiologia* 41:87–91
  126. Roberts D, Thelen D, Weinstein S. 1982. Parenteral and enteral nutrition: a cost-benefit audit. *Minn. Med.* 65:707–10
  127. Rodrigo CM, Garcia PJ. 1997. The effect

- of the composition of the enteral nutrition on infection in the critical patient. *Nutr. Hosp.* 12:80–84
128. Ryan JA, Page CP, Babcock L. 1981. Early postoperative jejunal feeding of elemental diet in gastrointestinal surgery. *Am. Surg.* 47:393–403
129. Saffle JR, Wiebke G, Jennings K, Morris SE, Barton RG. 1997. Randomized trial of immune-enhancing enteral nutrition in burn patients. *J. Trauma* 42:793–802
130. Sagar S, Harland P, Shields R. 1979. Early postoperative feeding with elemental diet. *Br. Med. J.* 1:293–95
131. Sandstrom R, Drott C, Hyltander A, Arfvidsson B, Schersten T, et al. 1993. The effect of postoperative intravenous feeding (TPN) on outcome following major surgery evaluated in a randomized study. *Ann. Surg.* 217:185–95
132. Schilling J, Vranjes N, Fierz W, Joller H, Gyurech D, et al. 1996. Clinical outcome and immunology of postoperative arginine, w-3 fatty acids, and nucleotide-enriched enteral feeding: a randomized prospective comparison with standard enteral and low calorie/low fat IV solutions. *Nutrition* 12:423–29
133. Senkal M, Mumme A, Eickhoff U, Geier B, Spath G, et al. 1997. Early postoperative enteral immunonutrition: clinical outcome and cost-comparison analysis in surgical patients. *Crit. Care Med.* 25: 1489–96
134. Senkal M, Zumtobel V, Bauer KH, Marpe B, Wolfram G, et al. 1999. Outcome and cost-effectiveness of perioperative enteral immunonutrition in patients undergoing elective upper gastrointestinal tract surgery: a prospective randomized study. *Arch. Surg.* 134(12):1309–16
135. Shike M, Russel DM, Detsky AS, Harrison JE, McNeil KG, et al. 1984. Changes in body composition in patients with small-cell lung cancer. The effect of total parenteral nutrition as an adjunct to chemotherapy. *Ann. Intern. Med.* 101(3):303–9
136. Shukla HS, Rao RR, Banu N, Gupta RM, Yadav RC. 1984. Enteral hyperalimentation in malnourished surgical patients. *Indian J. Med. Res.* 80:339–46
137. Simms JM, Oliver E, Smith JAR. 1980. A study of total parenteral nutrition (TPN) in major gastric and esophageal resection for neoplasia. *J. Parent. Enteral Nutr.* 4:422
138. Smith R, Hartemink R. 1988. Improvement of nutritional measures during preoperative parenteral nutrition in patients selected by the prognostic nutritional index: a randomized controlled trial. *J. Parent. Enteral Nutr.* 12:587–91
139. Smith RC, Hartemink RJ, Hollinshead JW, Gillett DJ. 1985. Fine bore jejunostomy feeding following major abdominal surgery: a controlled randomized clinical trial. *Br. J. Surg.* 72:458–61
140. Snyderman CH, Kachman K, Molsead L, Wagner R, D'Amico F, et al. 1999. Reduced postoperative infections with an immune-enhancing nutritional supplement. *Laryngoscope* 109(6):915–21
141. Studley HO. 1936. Percentage of weight loss: a basic indicator of surgical risk in patients with chronic peptic ulcer. *JAMA* 106:458–60
142. Takagi K, Yamamori H, Tashiro T. 2001. TPN induces bacterial translocation. *Nippon Rinsho* 59(55):191–95
143. Takagi K, Yamamori H, Yasuyoshi T, Nakejima N, Tashiro T. 2000. Modulating effects of the feeding route on stress response and endotoxin translocation in severely stressed patients receiving thoracic esophagectomy. *Nutrition* 16:355–60
144. Takesue Y, Ohge H, Uemura K, Imamura Y, Murakami Y, et al. 2002. Bacterial translocation in patient with Crohn's disease undergoing surgery. *Dis. Colon Rectum* 45(12):1665–71
145. Thompson B, Julian T, Stremple J. 1981. Perioperative total parenteral nutrition in patients with gastrointestinal cancer. *J. Surg. Res.* 30:497–500
146. Torosian MH. 1999. Perioperative

- nutrition support for patients undergoing gastrointestinal surgery: critical analysis and recommendations. *World J. Surg.* 23:565–69
147. Twomey PL, Patching SC. 1985. Cost-effectiveness of nutritional support. *J. Parent. Enteral Nutr.* 9:3–10
  148. van Bokhorst-de van der Schueren M, Quak JJ, von Blomberg-van der Flier BM, Kuik DJ, Langendoen SI, et al. 2001. Effect of perioperative nutrition with and without arginine supplementation, on nutritional status, immune function, postoperative morbidity, and survival in severely malnourished head and neck cancer patients. *Am. J. Clin. Nutr.* 73:323–32
  149. Veterans Affairs Total Parenteral Nutrition Cooperative Study. 1991. Perioperative total parenteral nutrition in surgical patients. *N. Engl. J. Med.* 325:525–32
  150. von Meyenfeldt MF, Meijerink WJHJ, Rouflart MMJ, Builmaassen MTHJ, Soeters PB. 1992. Perioperative nutritional support: a randomized clinical trial. *Clin. Nutr.* 11:180–86
  151. Weimann A, Bastian L, Bischoff WE, Grotz M, Hansel M, et al. 1998. The influence of arginine, omega-3 fatty acids and nucleotide-supplemented enteral support on systemic inflammatory response syndrome and multiple organ failure in patients after severe trauma. *Nutrition* 14(2):165–72
  152. Williford WO, Krol WF, Buzby GP. 1993. Comparison of eligible randomized patients with two groups of ineligible patients: Can the results of the VA total parenteral nutrition clinical trial be generalized? *J. Clin. Epidemiol.* 46:1025–34
  153. Wollman B, D'Agostino HB, Walas-Wigle JR, Easter DW, Beale A. 1995. Radiologic, endoscopic, and surgical gastrostomy: an institutional evaluation and meta-analysis of the literature. *Radiology* 197:699–704
  154. Woodcock NP, Ziegler D, Palmer MD, Buckley P, Mitchell CJ, et al. 2001. Enteral versus parenteral nutrition: a pragmatic study. *Nutrition* 17(1):1–12
  155. Woolfson A, Smith J. 1989. Elective nutritional support after major surgery: a prospective randomized trial. *Clin. Nutr.* 8:15
  156. Wu C-W, Meng HC, Mok K-T, Kung S-P, Lin S-H, et al. 1995. Effect of total parenteral nutrition on the postoperative outcome in aged patients with gastric cancer. *Dig. Surg.* 12:164–70
  157. Yamada N, Koyama J, Hioki K, Yamada T, Yamamoto M. 1983. Effect of postoperative total parenteral nutrition (TPN) as an adjunct to gastrectomy for advanced gastric carcinoma. *Br. J. Surg.* 70:267–74

## CONTENTS

---

FRONTISPIECE— <i>Frank Chytil</i>	xiv
ROUGH AND ROCKY ROAD TO THE RETINOID REVOLUTION, <i>Frank Chytil</i>	1
MECHANISM AND REGULATION OF SELENOPROTEIN SYNTHESIS, <i>Donna M. Driscoll and Paul R. Copeland</i>	17
IRON STATUS AND NEURAL FUNCTIONING, <i>John L. Beard and James R. Connor</i>	41
INSIGHTS INTO THE PATHOGENESIS OF GALACTOSEMIA, <i>Nancy D. Leslie</i>	59
DIET AND NUTRITION IN POOR AND MINORITY COMMUNITIES IN THE UNITED STATES 100 YEARS AGO, <i>Robert Dirks</i>	81
DIFFERENT APPROACHES TO DEFINE INDIVIDUAL AMINO ACID REQUIREMENTS, <i>Paul B. Pencharz and Ronald O. Ball</i>	101
VITAMIN D AND ITS ANALOGS AS REGULATORS OF IMMUNE ACTIVATION AND ANTIGEN PRESENTATION, <i>Matthew D. Griffin, Nianzeng Xing, and Rajiv Kumar</i>	117
NUTRITION AND PREVENTION OF TYPE 2 DIABETES, <i>T. Costacou and E.J. Mayer-Davis</i>	147
BIOLOGIC MECHANISMS OF THE PROTECTIVE ROLE OF LUTEIN AND ZEAXANTHIN IN THE EYE, <i>Norman I. Krinsky, John T. Landrum, and Richard A. Bone</i>	171
NUTRITIONAL REGULATION OF MILK FAT SYNTHESIS, <i>Dale E. Bauman and J. Mikko Griinari</i>	203
TROPHIC AND CYTOPROTECTIVE NUTRITION FOR INTESTINAL ADAPTATION, MUCOSAL REPAIR, AND BARRIER FUNCTION, <i>Thomas R. Ziegler, Mary E. Evans, Concepción Fernández-Estívariz, and Dean P. Jones</i>	229
NUTRITION IN THE PERIOPERATIVE PATIENT, <i>Lyn Howard and Christopher Ashley</i>	263
PHYSIOLOGY AND MOLECULAR BIOLOGY OF DIETARY IRON ABSORPTION, <i>Silvia Miret, Robert J. Simpson, and Andrew T. McKie</i>	283

GUGULIPID: A NATURAL CHOLESTEROL-LOWERING AGENT, <i>Nancy L. Urizar and David D. Moore</i>	303
CHALLENGES AND APPROACHES TO REDUCING FOODBORNE ILLNESS, <i>Catherine E. Woteki and Brian D. Kineman</i>	315
DIETARY, EVOLUTIONARY, AND MODERNIZING INFLUENCES ON THE PREVALENCE OF TYPE 2 DIABETES, <i>Leslie Sue Lieberman</i>	345
IN VIVO MEASUREMENT OF FLUXES THROUGH METABOLIC PATHWAYS: THE MISSING LINK IN FUNCTIONAL GENOMICS AND PHARMACEUTICAL RESEARCH, <i>Marc K. Hellerstein</i>	379
COMMON ENDOCRINE CONTROL OF BODY WEIGHT, REPRODUCTION, AND BONE MASS, <i>Shu Takeda, Florent Elefteriou, and Gerard Karsenty</i>	403
INDEXES	
Subject Index	413
Cumulative Index of Contributing Authors, Volumes 19–23	433
Cumulative Index of Chapter Titles, Volumes 19–23	436
ERRATA	
An online log of corrections to <i>Annual Review of Nutrition</i> chapters (if any, 1997 to the present) may be found at <a href="http://nutr.annualreviews.org/">http://nutr.annualreviews.org/</a>	