

Forum Review

Mitochondrial Uncoupling Proteins in the Central Nervous System

JEONG SOOK KIM-HAN¹ and LAURA L. DUGAN^{1,2,3}

ABSTRACT

Mitochondrial uncoupling proteins (UCPs), a subfamily of the mitochondrial transporter family, are related by sequence homology to UCP1. This protein, which is located in the inner mitochondrial membrane, dissipates the proton gradient between the intermembrane space and the mitochondrial matrix to uncouple electron transport from ATP synthesis. UCP1 (thermogenin) was first discovered in brown adipose tissue and is responsible for non-shivering thermogenesis. Expression of mRNA for three other UCP isoforms, UCP2, UCP4, and BMCP1/UCP5, has been found at high levels in brain. However, the physiological function(s) of UCPs in the brain have not been determined, although it has recently been postulated that UCPs regulate free radical flux from mitochondria by physiologically modulating mitochondrial membrane potential. In the CNS, this hypothesis has been studied primarily for UCP2. UCP2 message has been shown to be up-regulated in the CNS by stress signals such as kainate administration or ischemia, and overexpression of UCP2 has been reported to be neuroprotective against oxidative stress *in vivo* and *in vitro*, although the exact mechanism has not been fully established. In this review, studies on UCPs in the nervous system will be reviewed, and the potential roles of these intriguing proteins in acute and chronic diseases of the nervous system will be discussed.

Antioxid. Redox Signal. 7, 1173–1181.

GENERAL INTRODUCTION TO UNCOUPLING PROTEINS

THE ELECTROCHEMICAL PROTON GRADIENT (the proton motive force, $\Delta p = \Delta\psi + \Delta pH$) between the intermembrane space and the matrix of mitochondria is produced by the electron transport chain residing in the mitochondrial inner membrane (Fig. 1). As first proposed by Mitchell (61), this gradient is now known to be the driving force for ATP synthesis, called “coupling.” Many proteins, including members of the mitochondrial carrier family, harness Δp for their function, some using its electrical component (ADP/ATP carrier and glutamate/aspartate carrier) and others using the ΔpH gradient (phosphate carrier and other carriers using substrate- H^+ symport). Any process using Δp *without* ATP synthesis has been termed “uncoupling.” Uncoupling protein 1 (UCP1; thermogenin) was first recognized as the protein-mediated uncoupling activity in brown adipose tissues (BAT), in which

it acts as the non-shivering heat generator to maintain the body temperature in newborns and hibernating mammals (9, 47, 64).

A number of other protein members of the SLC25 subfamily of the mitochondrial transporter superfamily (for review, see 67) have been designated as “uncoupling proteins” (UCPs) based initially on their sequence homology with UCP1 and evidence that they had many of the same functional motifs. Of those studied to date, all appear to be located in the inner mitochondrial membrane and to uncouple the proton gradient. More than 45 genes encoding UCP isoforms have been described in single-cell organisms, plants, and animals, including humans (54). In addition to UCP1, many of these additional UCP isoforms have been cloned from plants and mammals, and their physiological roles have begun to be investigated (8). UCPs are expressed in a tissue-specific manner, suggesting distinct roles and regulation of their function in different locations. UCP2 has broad tissue

¹Department of Neurology and ²Department of Medicine, Washington University School of Medicine, St. Louis, MO.

³Department of Medicine, University of California San Diego, San Diego, CA.




FIG. 1. Effect of UCP on the production of ROS. (A) Under normal conditions, the proton gradient established by the electron transport chain (ETC) is coupled to ATP synthesis through F_1F_0 -ATPase. When the membrane is highly polarized, electron transfer from the ETC to oxygen is enhanced, increasing levels of ROS (superoxide, H_2O_2). In the presence of purine nucleotides (GDP), UCP activity is inhibited and elevated membrane potential and ROS production are maintained. (B) When overexpression of UCP or activation of the protein occurs, *e.g.*, by binding of free fatty acids (FFA) or coenzyme Q10, the proton gradient is brought below the critical point for ROS production, and mitochondrial ROS formation decreases.

expression, whereas UCP1 is almost exclusively expressed in BAT, and UCP3 is found mainly in BAT and skeletal muscle. UCP2 and UCP3 have the highest degree of homology to UCP1. Although they have significantly lower degrees of homology to UCP1, UCP4 and BMCP1/UCP5 have been added as putative members of the UCP family because they show most of the functional motifs of UCP1. Another isoform, SLC25A30, was recently reported by the RIKEN cDNA library to have a high degree of sequence homology to BMCP1/UCP5, but has not yet been studied. In BAT, UCP1 is highly expressed as one of the major mitochondrial proteins and, when activated, results in sufficient heat generation to maintain body temperature in newborns. Evidence suggests that UCP2 is *not* important for thermogenesis, however (2, 87). All UCPs are expressed at significantly lower levels in their respective tissues compared with UCP1 in BAT, suggesting that the function of these UCP isoforms may not be thermogenesis. The observation that the newer UCPs are expressed at such low levels prompted introduction of the concept of “mild uncoupling” (76)—in which functional regulation of mitochondrial reactive oxygen species (ROS) production occurs through modest decreases in Ψ_m —for the non-UCP1 family members (Fig. 1). Mitochondria are considered to be one of the main sources for intracellular ROS, and the mitochondrial membrane potential is the driving force for ROS generation as well as ATP production. A small decrease in Ψ_m can dramatically reduce production of ROS, specifically hydrogen peroxide (H_2O_2) (49). In addition, this mild uncoupling activity has been proposed to be one of the key modulators of cellular metabolism and signal transduction through regulation of mitochondrial ROS production (11, 32), because many critical signaling pathways have redox-sensitive components (30, 31). As the involvement of mitochondrial ROS dysregulation in neurodegenerative diseases has been a focus of research for many decades, and altered production or decomposition of ROS has been observed in many experimental models of CNS injury and neurodegenerative diseases, the possibility that UCPs, by regulating mitochondrial ROS levels, could be important modifiers of CNS damage has recently emerged. In this review, the involvement of UCPs in regulation of ROS will be

described, and the physiological and pathological roles of UCPs in the nervous system will be discussed.

UCP EXPRESSION: TRANSCRIPTIONAL AND POSTTRANSCRIPTIONAL REGULATION

Although regulation of UCP gene expression at the mRNA level has been demonstrated in response to a diverse range of stimuli, including exposure to cold, high fat and ketogenic diets, starvation, leptin, insulin, inflammatory cytokines, and endotoxin (for review, see 14), regulation of protein is rarely reported. In part, this reflects problems with currently available UCP antibodies, including both custom-prepared and commercially available antibodies. Most are not exclusively specific to each isoform (69), and we have found that several antibodies detect 32–36-kDa bands in various knockout mice (Dugan and Kim-Han, unpublished observations), suggesting that specific, well characterized antibodies would be extremely useful reagents for future studies on the UCPs.

Regulation of UCP mRNA has not, in many cases, been followed by changes in UCP protein expression (43, 75), suggesting posttranscriptional regulation of UCP expression (23, 69). Recently, message for UCP2 has been found without concomitant protein expression (84). This disconnect between mRNA and protein expression results from an upstream AUG stop codon and alternative open reading frames (ORFs) in the 5′-untranslated region as a key determinant of posttranscriptional regulation (69). One or three upstream ORFs have been identified in UCP2/3 and BMCP1/UCP5, respectively, and these result in dramatically reduced efficiency of translation (Fig. 2). In addition, more than six mRNA splicing variants have been detected for BMCP1/UCP5 mRNA (Dugan and Kim-Han, unpublished observations). We have also experienced inhibition of UCP expression after *in vitro* transfection in mammalian cells, although mRNA was clearly present (Dugan and Kim-Han, unpublished observations). Other posttranscriptional mechanisms, including changes in mRNA stability, need to be studied for these pro-




FIG. 2. The putative ORFs at the upstream of the starting codon in mouse UCP mRNAs. Among UCPS, UCP1 does not have any putative ORF. UCP2 and UCP5/BMCP1 have three ATG codons at the upstream of the starting codon with one and three ORFs, respectively. Rat UCP4 mRNA has an ORF found in the 5'-untranslated region. Slc25a30 is highly homologous to UCP5/BMCP1. (M, ATG codon. The size of mRNA is shown in the box at the right side.)

teins. In summary, it is important to bear in mind that the regulation of UCP mRNA expression may not correspond to that of protein expression, and that mRNA data as predictive of UCP protein or activity should be interpreted with some degree of caution.

REGULATION OF UCP ACTIVITY

Well known regulators of UCP activity are free fatty acids and purine nucleotides. Polyunsaturated fatty acids activate UCP, whereas bovine serum albumin, a fatty acid binding protein, inhibits UCP activity (25, 47). The purine nucleotides, ADP and GDP, also inhibit the proton conductance induced by various stimuli *in vitro* by binding to a specific allosteric site. The specificity and preference of a given UCP isoform for specific free fatty acid species has not been fully defined, but would be of interest because different fatty acid pools are released by varying stimuli, including ischemia and traumatic injury (1, 4, 36, 52, 53, 70, 79). In addition, there remains a controversy about whether free fatty acids are not only "activators" of UCP activity through a specific, high-affinity binding site, but "transported substrates" of UCPS, being cotransported with protons under selected circumstances (34). This alternative fatty acid transport activity has been most convincingly shown to date for UCP1 and UCP3. Regardless of their mechanisms of action, the requirement for nanomolar to micromolar concentrations of free fatty acids for full UCP activation has been well established (25, 35).

It was recently demonstrated that specific free radicals and oxidized macromolecules can modify the activity of UCPS. Echter *et al.* reported that superoxide increases mitochondrial proton conductance through UCPS (26). This superoxide-mediated increase in conductance was sensitive to GDP and

required fatty acids. Superoxide-stimulated proton conductance through UCPS was reversible, pH-dependent, and sensitive to superoxide dismutase (SOD), but not to catalase (26). Breakdown products of lipid peroxidation, such as 4-hydroxynonenal and its related compounds, specifically induce proton conductance of mitochondria through the uncoupling proteins (UCP1, UCP2, and UCP3) and the adenine nucleotide translocase in animals (27) and through StUCP (UCP homologue in *Solanum tuberosum*) in plants (77). Moreover, products of lipid peroxidation, such as cleaved hydroperoxy-fatty acids and hydroxy-fatty acids, have been reported to activate UCP2 and promote feedback down-regulation of mitochondrial ROS production (10).

Endogenously produced superoxide also increases proton conductance through activation of UCP3 in skeletal muscle, subsequently limiting mitochondrial production of superoxide (82). Recently, Krauss and colleagues reported UCP2 activation by superoxide introduced exogenously using mitochondria from kidney and spleen (50). Exogenously generated superoxide also activated UCP2 in thymocytes and B cells in mice, and activation was inhibited by MnTBAP, a SOD mimetic, or manganese superoxide dismutase (MnSOD), but not by glutathione peroxidase (GPX1). Although UCP activation by superoxide was absent in UCP2 knockout (KO) mice, implying that UCP2 may be activated by superoxide, in another study (18), UCP2 was clearly not activated by superoxide. As other UCP isoforms, specifically UCP1 (26) and UCP3 (26, 82), can be activated by superoxide, the expression of other UCPS in UCP2 KO mice should be determined to verify that other UCPS were not induced as has been shown previously (19). Specific activation of UCPS by superoxide also needs to be further confirmed by comparison with other inner mitochondrial membrane proteins. In addition, Li and colleagues pointed out that the superoxide-generating system, xanthine oxidase metabolism of xanthine, used for some of

these studies, consumes oxygen to produce superoxide, thus reducing oxygen content in the reaction mixture and modifying mitochondrial respiratory studies (56). Experiments using an *in vitro* reconstituted system might give more direct answers to this controversial question.

Ubiquinone has also been reported to be an activator of UCP1 (24). Short-term oral administration of ubiquinone induced nigral mitochondrial uncoupling and prevented dopamine cell loss after 1-methyl-4-phenyl-1,2,5,6-tetrahydropyridine administration in monkeys (41), although whether the effects of ubiquinone were due to activation of a UCP isoform, or to the multiple other actions (*e.g.* antioxidant) of this electron transport chain component, are not yet established. The requirement of ubiquinone for UCP activation has not been clearly demonstrated in other model systems, such as yeast and reconstituted liposomes (29, 42), so this remains an important area of UCP research.

Information regarding the regulation of UCP activity is likely to be more relevant than that of expression of UCP mRNA or protein *per se*. Why? With the exception of UCP1, expression of other UCP isoforms appears to be quite low, at least two orders of magnitude lower than that of UCP1, measured by [³H]GDP binding (86). Furthermore, even in BAT, with its high content of UCP1, it is only after activation of UCP1 by fatty acids that uncoupling and thermogenesis can be detected, indicating that the presence of protein will not necessarily predict uncoupling activity. Experiments performed to determine UCP activity have generally been performed with isolated mitochondria from tissues that express mixed pools of UCPs, as well as other mitochondrial carrier proteins which may contribute to a small extent to "uncoupling" activity. Many studies using tissues and transfected cells have assigned uncoupling activity to a specific UCP without knowing the isoforms present in their systems. The contribution of one individual UCP in such preparations and the degree to which mitochondria "compensate" for the absence or overexpression of a single UCP remain an open question. Finally, the UCP expression profile in a given experimental model may differ from that reported in the literature. It is clear that there are strain differences in UCP expression even at baseline, and there are differing responses reported for the same treatment in different strains (37, 81).

MODULATION OF MITOCHONDRIAL ROS PRODUCTION BY UCPs

One of the main sites for intracellular ROS formation is the mitochondrial electron transport chain, in which superoxide is generated as a by-product of electron transport. In isolated mitochondria, inhibitors of complex I and III significantly increase the production of ROS (10, 13, 15). Production is closely dependent on mitochondrial membrane potential, *i.e.*, a small increase in Ψ_m can cause a robust production of superoxide and H₂O₂. A 10% decrease in Ψ_m resulted in a 55% decrease in H₂O₂ production (49). A 10-mV decrease in Ψ_m reduced superoxide production up to 70% by complex I following reverse electron flow from glycerol 3-phosphate (62). Physiological activation of UCPs has therefore been

proposed to decrease ROS production. Although UCPs act as uncouplers, because of their low level of expression and the requirement for small-molecule activators for UCPs to be fully activated, the contribution of UCPs to regulation of Ψ_m is likely to be quite small, leading to the previously-mentioned concept of "mild uncoupling" (62, 76). This concept explains why there are no significant changes in basal levels of proton conductance in either UCP-overexpressing or KO animals until nonphysiological conditions are present. The role of UCPs in regulating ROS was first reported in 1997. In this study, GDP, an inhibitor of UCPs, increased both Ψ_m and production of H₂O₂ in mitochondria from BAT and liver (63). Macrophages from UCP2 KO mice produce more ROS, increasing their resistance to toxoplasmosis infection (2). Antisense treatment to decrease UCP2 expression increased ROS production and lipid peroxidation in murine endothelial cells (22). Mitochondrial H₂O₂ production is lowered by calorie restriction, and this appears to involve up-regulation of UCP3 in rat muscle mitochondria (7). Glucose-induced ROS production through early mitochondrial hyperpolarization was inhibited by UCP3 and UCP1 overexpression in dorsal root ganglia (DRG) neurons (83). In brain mitochondria, UCP2 overexpression also reduced mitochondrial ROS content, determined by flow cytometry using dichlorofluorescein fluorescence (59, 63, 66). Taken together, UCPs modulate ROS production possibly through decreasing mitochondrial membrane potential.

EXPRESSION AND FUNCTION OF UCPs IN BRAIN

Message for three UCP isoforms, UCP2, UCP4, and BMCP1/UCP5, has been found in brain. UCP2 mRNA is ubiquitously expressed in various tissues, including brain in rodents (38, 72), and is mainly neuronal, although there is some debate about whether UCP2 protein is expressed predominantly in macrophages under basal conditions (73). UCP4 message is exclusively in brain (58), whereas BMCP1/UCP5 is expressed in brain and several other tissues, including heart and kidney (45, 48, 74). Although no expression of UCP1 or UCP3 mRNA has been reported for brain, recently, UCP3 protein was found in rat DRG neurons immediately after dissection, but not in cultured DRG neurons (83). The authors suggested that UCP3 was protective against glucose-induced neuronal death in DRG neurons. Carefully performed quantitative RT-PCR analysis to determine the expression of UCPs in brain cortex indicated the following order: UCP5 > UCP4 > UCP2 >>> UCP3, UCP1 (55). Recently, expression of an isoform of BMCP1/UCP5 in fruit flies was also found predominantly in the head of adult flies (33), supporting an important role for this putative UCP in the CNS. The amount of overall UCP protein, shown by [³H]GTP binding in brain, has been reported to be substantially lower than in BAT or spleen, but higher than in kidney, skeletal muscle, or liver (44). Despite the presence of at least three isoforms of UCP in brain, UCP2 has been the focus of the majority of studies on brain injury and neurodegenerative diseases. In normal brain, UCP2 mRNA expression is localized to specific brain regions, whereas in pathological conditions, UCP2 mRNA is

up-regulated in the regions affected by the injury. UCP2 mRNA is induced by transient ischemia in lesioned entorhinal cortex (5, 57) and in the periinfarct area in the neocortex and caudate putamen after 3–4 days (19). Both sublethal ischemia (59) and chemically induced seizures (21) increase mRNA expression in mouse hippocampus. Kainic acid (KA), given by intraperitoneal injection to mice, also induced UCP2 mRNA in the CA1 subfield of the hippocampus, the dorsal endopiriform nucleus, and the piriform cortex, areas sensitive to KA excitotoxicity (6, 51). Induction was primarily neuronal and reached a maximum at 24 h, returning to basal levels within 72 h post injection. This neuronal induction of UCP2 mRNA by KA was more extensive in 129T2SvEmsJ, a KA-sensitive mouse strain, than in C57BL/6J, a KA-insensitive one, suggesting a specific link between UCP2 in KA excitotoxicity (17). Lipopolysaccharide, a commonly-used activator of oxidative stress, induced UCP2 mRNA, but not BMCP1/UCP5 mRNA in mice (12). Administration of a ketogenic diet increased mitochondrial respiration by fatty acids, possibly through induction of one or more uncoupling proteins (80). It has been postulated that ROS generation is one of the main factors for both neurodegeneration and aging, so UCP-dependent reduction of ROS in the nervous system has the potential to be neuroprotective in age-related neurodegenerative diseases, as well (28, 46).

Although UCP4 and BMCP1/UCP5 have sequence homology to UCP1, and UCP2–3 (29–34% and 34–39%, respectively) and show uncoupling activity in *in vitro* systems (58, 74), their function in brain has yet to be established. BMCP1/UCP5 mRNA was down-regulated by hypoxia and up-regulated by oxidative stressors, such as hyperoxia, 4-hydroxynonenal, or *tert*-butyl hydroperoxide in SH-SY5Y neuroblastoma cells (71). BMCP1/UCP5 message was decreased by chronic hypoxia and increased by transient global cerebral ischemia in rats (71), but was unaltered by cerebral ischemia in adult (19) and neonatal (Kim-Han, Reichert, and Dugan, unpublished observations) mice.

Among the three main isoforms of UCP expressed in brain, UCP2 is the most well characterized. UCP2 appears to reduce ROS production in brain under pathological, but not physiological, conditions. Focal cerebral ischemia produced by middle cerebral artery occlusion (MCAO) was used to test the role of UCP2 in CNS injury using both UCP2 transgenic and UCP2 KO mice. Both overexpressing and KO mice were resistant to ischemic injury (19, 59). Overexpression of UCP2/3 was neuroprotective against MCAO *in vivo* and oxygen-glucose deprivation *in vitro*. Levels of ROS in the mitochondrial matrix were lower in mitochondria from UCP2/3 transgenic mice than from nontransgenic controls, although there was no difference in total production of ROS (59, 68). In response to MCAO, UCP2 KO mice also showed a smaller infarct volume, less TUNEL-positive cells, no cytochrome *c* translocation, and less lipid peroxidation products compared with the control mice. The paradox that both overexpression and targeted deletion of UCP2 were neuroprotective has been explained by the observation that UCP2 KO mice show higher expression of MnSOD and had higher mitochondrial [GSH] than wild-type mice. Like the dicarboxylate and 2-oxoglutarate carriers (16), UCP2 may regulate mitochondrial GSH uptake. As UCP2 KO mice are presumed to produce more mitochondrial ROS, up-

regulation of MnSOD and lowered consumption of GSH after ischemia have been proposed as compensatory mechanisms (19). These reports suggested that UCP2 can modify mitochondrial antioxidant defense systems. Overexpression of UCP2 also resulted in neuroprotection and lower caspase-3 immunoreactivity after entorhinal cortex lesioning (5) and was protective against oxidative stress-induced cell death in PC12 cells (21). Although UCP2 overexpression does not prevent seizures after scopolamine/pilocarpine treatment, it reduced cell death after the seizures (21). On the other hand, overexpression of UCP2 may cause cells to have difficulty providing enough ATP production for cellular metabolism, leading to mitochondrial biogenesis to support ATP production (40). In fact, UCP2 overexpression increased mitochondrial number, presumably to compensate for decreased ATP production by uncoupled mitochondria (21). In addition, UCP2 negatively regulates secretion of insulin (87) and dopamine (85) through an ATP-dependent process. How these effects factor into neuroprotection or neurodegeneration has not yet been determined, and may be complex.

Ethanol sensitivity was decreased in UCP2-overexpressing mice and increased in UCP2 KO mice. UCP2 also modified the impairment of pain and temperature sensation induced by ethanol, which is consistent with localization of UCP2 in primary sensory afferents of the spinal cord (38). Overexpression of UCP1 or UCP3 is also neuroprotective in glucose-induced neuronal cell death through glucose-induced superoxide production and glucose-induced mitochondrial hyperpolarization (83). Recently, the “uncoupling to survive” hypothesis was suggested by the positive correlation between metabolic rate and life span across many species (78). ROS production induced by oligomycin was significantly ($p < 0.05$) reduced in ketogenic diet-fed mice compared to *ad libitum* controls, suggesting that a ketogenic diet may have neuroprotective effects by diminishing ROS production through activation of mitochondrial UCPs (80), although the effects of ketogenesis on metabolism are quite complex, and include modulation of insulin signaling.

CONCLUSION AND PERSPECTIVES

UCPs are normally expressed in brain and regulated by CNS injury in affected brain regions, where they are postulated to cause mild uncoupling, reducing electron leakage from the electron transport chain, and thereby decrease superoxide generation. Mitochondrial ROS generation in pathological conditions is dependent on mitochondrial membrane potential, and despite a number of questions that remain to be answered, evidence suggests that overexpression of UCPs can be neuroprotective. However, in some situations, UCP activation may result in a balance between beneficial and detrimental effects. UCP expression can be proapoptotic, as well as antiapoptotic, depending on transcriptional and biochemical regulation (20, 60).

Mitochondrial membrane potential is the main regulator of mitochondria Ca^{2+} uptake, and many Ca^{2+} -dependent pathways are involved in neuronal cell death. Understanding the role of mild uncoupling in Ca^{2+} homeostasis and mitochondrial free radical generation in excitotoxicity may help our

understanding of a key mechanism of neurotoxicity (64). Finally, overexpression of UCP results in growth inhibition in yeast (3, 74), an effect that may be related not only to changes in energy metabolism such as reduced ATP production, but may also reflect dysregulation of redox-sensitive pathways, including activation of mitogen-activating protein kinase signaling.

The idea that this is due to modification of mitochondrial ROS production places these proteins in the intriguing position of integrating both metabolic function and oxidative stress. Although it is suggested that modification of UCPs can lead to many other cellular responses, including changes in antioxidant defense systems and mitochondrial biogenesis, careful investigation will be needed to identify the detailed mechanisms behind their neuroprotective effects. Elucidating the mechanism(s) of UCP action in the regulation of energy metabolism and ROS generation would advance our understanding of how cells regulate their "power plants" to respond to normal physiology and stress conditions.

ACKNOWLEDGMENTS

This work was supported by NS 41796 (L.L.D.). The authors would like to thank Scott Gianino for excellent technical assistance with this work.

ABBREVIATIONS

BAT, brown adipose tissues; DRG, dorsal root ganglia; GSH, reduced glutathione; H_2O_2 , hydrogen peroxide; KA, kainic acid; KO, knockout; MCAO, middle cerebral artery occlusion; MnSOD, manganese superoxide dismutase; ORF, open reading frame; ROS, reactive oxygen species; SOD, superoxide dismutase; UCP, mitochondrial uncoupling protein; UCP1, thermogenin, uncoupling protein 1.

REFERENCES

- Agardh CD, Chapman AG, Pelligrino D, and Siesjo BK. Influence of severe hypoglycemia on mitochondrial and plasma membrane function in rat brain. *J Neurochem* 38: 662–668, 1982.
- Arsenijevic D, Onuma H, Pecqueur C, Raimbault S, Manning BS, Miroux B, Couplan E, Alves-Guerra MC, Goubern M, Surwit R, Bouillaud F, Richard D, Collins S, and Ricquier D. Disruption of the uncoupling protein-2 gene in mice reveals a role in immunity and reactive oxygen species production. *Nat Genet* 26: 435–439, 2000.
- Bathgate B, Freebairn EM, Greenland AJ, and Reid GA. Functional expression of the rat brown adipose tissue uncoupling protein in *Saccharomyces cerevisiae*. *Mol Microbiol* 6: 363–370, 1992.
- Bazan NG Jr. Effects of ischemia and electroconvulsive shock on free fatty acid pool in the brain. *Biochim Biophys Acta* 218: 1–10, 1970.
- Bechmann I, Diano S, Warden CH, Bartfai T, Nitsch R, and Horvath TL. Brain mitochondrial uncoupling protein 2 (UCP2): a protective stress signal in neuronal injury. *Biochem Pharmacol* 64: 363–367, 2002.
- Ben-Ari Y and Cossart R. Kainate, a double agent that generates seizures: two decades of progress. *Trends Neurosci* 23: 580–587, 2000.
- Bevilacqua L, Ramsey JJ, Hagopian K, Weindrich R, and Harper ME. Effects of short- and medium-term calorie restriction on muscle mitochondrial proton leak and reactive oxygen species production. *Am J Physiol Endocrinol Metab* 286: E852–E861, 2004.
- Bouillaud F, Couplan E, Pecqueur C, and Ricquier D. Homologues of the uncoupling protein from brown adipose tissue (UCP1): UCP2, UCP3, BMCP1 and UCP4. *Biochim Biophys Acta* 1504: 107–119, 2001.
- Boyer BB, Barnes BM, Lowell BB, and Grujic D. Differential regulation of uncoupling protein gene homologues in multiple tissues of hibernating ground squirrels. *Am J Physiol* 275: R1232–R1238, 1998.
- Brand MD, Affourtit C, Esteves TC, Green K, Lambert AJ, Miwa S, Pakay JL, and Parker N. Mitochondrial superoxide: production, biological effects, and activation of uncoupling proteins. *Free Radic Biol Med* 37: 755–767, 2004.
- Brookes PS, Levonen AL, Shiva S, Sarti P, and Darley-Usmar VM. Mitochondria: regulators of signal transduction by reactive oxygen and nitrogen species. *Free Radic Biol Med* 33: 755–764, 2002.
- Busquets S, Alvarez B, Van Royen M, Figueras MT, Lopez-Soriano FJ, and Argiles JM. Increased uncoupling protein-2 gene expression in brain of lipopolysaccharide-injected mice: role of tumour necrosis factor- α ? *Biochim Biophys Acta* 1499: 249–256, 2001.
- Cadenas E, Boveris A, Ragan CI, and Stoppani AO. Production of superoxide radicals and hydrogen peroxide by NADH-ubiquinone reductase and ubiquinol-cytochrome c reductase from beef-heart mitochondria. *Arch Biochem Biophys* 180: 248–257, 1977.
- Cannon B and Nedergaard J. Brown adipose tissue: function and physiological significance. *Physiol Rev* 84: 277–359, 2004.
- Casteilla L, Rigoulet M, and Penicaud L. Mitochondrial ROS metabolism: modulation by uncoupling proteins. *IUBMB Life* 52: 181–188, 2001.
- Chen Z and Lash LH. Evidence for mitochondrial uptake of glutathione by dicarboxylate and 2-oxoglutarate carriers. *J Pharmacol Exp Ther* 285: 608–618, 1998.
- Clavel S, Paradis E, Ricquier D, and Richard D. Kainic acid upregulates uncoupling protein-2 mRNA expression in the mouse brain. *Neuroreport* 14: 2015–2017, 2003.
- Couplan E, del Mar Gonzalez-Barroso M, Alves-Guerra MC, Ricquier D, Goubern M, and Bouillaud F. No evidence for a basal, retinoic, or superoxide-induced uncoupling activity of the uncoupling protein 2 present in spleen or lung mitochondria. *J Biol Chem* 277: 26268–26275, 2002.
- de Bilbao F, Arsenijevic D, Vallet P, Hjelle OP, Ottersen OP, Bouras C, Raffin Y, Abou K, Langhans W, Collins S, Plamondon J, Alves-Guerra MC, Haguenaer A, Garcia I, Richard D, Ricquier D, and Giannakopoulos P. Resistance to cerebral ischemic injury in UCP2 knockout mice: evidence for a role of UCP2 as a regulator of mitochondrial glutathione levels. *J Neurochem* 89: 1283–1292, 2004.

20. Dejean L, Camara Y, Sibille B, Solanes G, and Villarroya F. Uncoupling protein-3 sensitizes cells to mitochondrial-dependent stimulus of apoptosis. *J Cell Physiol* 201: 294–304, 2004.
21. Diano S, Matthews RT, Patrylo P, Yang L, Beal MF, Barnstable CJ, and Horvath TL. Uncoupling protein 2 prevents neuronal death including that occurring during seizures: a mechanism for preconditioning. *Endocrinology* 144: 5014–5021, 2003.
22. Duval C, Negre-Salvayre A, Dogilo A, Salvayre R, Penicaud L, and Casteilla L. Increased reactive oxygen species production with antisense oligonucleotides directed against uncoupling protein 2 in murine endothelial cells. *Biochem Cell Biol* 80: 757–764, 2002.
23. Ealey KN, El-Sohehy A, and Archer MC. Effects of dietary conjugated linoleic acid on the expression of uncoupling proteins in mice and rats. *Lipids* 37: 853–861, 2002.
24. Echtay KS, Winkler E, and Klingenberg M. Coenzyme Q is an obligatory cofactor for uncoupling protein function. *Nature* 408: 609–613, 2000.
25. Echtay KS, Winkler E, Frischmuth K, and Klingenberg M. Uncoupling proteins 2 and 3 are highly active H⁺ transporters and highly nucleotide sensitive when activated by coenzyme Q (ubiquinone). *Proc Natl Acad Sci U S A* 98: 1416–1421, 2001.
26. Echtay KS, Roussel D, St-Pierre J, Jekabsons MB, Cadenas S, Stuart JA, Harper JA, Roebuck SJ, Morrison A, Pickering S, Clapham JC, and Brand MD. Superoxide activates mitochondrial uncoupling proteins. *Nature* 415: 96–99, 2002.
27. Echtay KS, Esteves TC, Pakay JL, Jekabsons MB, Lambert AJ, Portero-Otin M, Pamplona R, Vidal-Puig AJ, Wang S, Roebuck SJ, and Brand MD. A signalling role for 4-hydroxy-2-nonenal in regulation of mitochondrial uncoupling. *EMBO J* 22: 4103–4110, 2003.
28. Emerit I, Edeas M, and Bricaire F. Neurodegenerative diseases and oxidative stress. *Biomed Pharmacother* 58: 39–46, 2004.
29. Esteves TC, Echtay KS, Jonassen T, Clarke CF, and Brand MD. Ubiquinone is not required for proton conductance by uncoupling protein 1 in yeast mitochondria. *Biochem J* 379: 309–315, 2004.
30. Finkel T. Reactive oxygen species and signal transduction. *IUBMB Life* 52: 3–6, 2001.
31. Finkel T. Oxidant signals and oxidative stress. *Curr Opin Cell Biol* 15: 247–254, 2003.
32. Forman HJ, Fukuto JM, and Torres M. Redox signaling: thiol chemistry defines which reactive oxygen and nitrogen species can act as second messengers. *Am J Physiol Cell Physiol* 287: C246–C256, 2004.
33. Fridell YW, Sanchez-Blanco A, Silvia BA, and Helfand SL. Functional characterization of a *Drosophila* mitochondrial uncoupling protein. *J Bioenerg Biomembr* 36: 219–228, 2004.
34. Garlid KD, Jaburek M, Jezek P, and Varecha M. How do uncoupling proteins uncouple? *Biochim Biophys Acta* 1459: 383–389, 2000.
35. Gonzalez-Barroso MM, Fleury C, Arechaga I, Zaragoza P, Levi-Meyreuis C, Raimbault S, Ricquier D, Bouillaud F, and Rial E. Activation of the uncoupling protein by fatty acids is modulated by mutations in the C-terminal region of the protein. *Eur J Biochem* 239: 445–450, 1996.
36. Hillered L and Chan PH. Role of arachidonic acid and other free fatty acids in mitochondrial dysfunction in brain ischemia. *J Neurosci Res* 20: 451–456, 1988.
37. Hioki C, Yoshida T, Kogure A, Takakura Y, Umekawa T, Yoshioka K, Shimatsu A, and Yoshikawa T. Effects of growth hormone (GH) on mRNA levels of uncoupling proteins 1, 2, and 3 in brown and white adipose tissues and skeletal muscle in obese mice. *Horm Metab Res* 36: 607–613, 2004.
38. Horvath B, Spies C, Horvath G, Kox WJ, Miyamoto S, Barry S, Warden CH, Bechmann I, Diano S, Heemskerk J, and Horvath TL. Uncoupling protein 2 (UCP2) lowers alcohol sensitivity and pain threshold. *Biochem Pharmacol* 64: 369–374, 2002.
39. Horvath TL, Warden CH, Hajos M, Lombardi A, Goglia F, and Diano S. Brain uncoupling protein 2: uncoupled neuronal mitochondria predict thermal synapses in homeostatic centers. *J Neurosci* 19: 10417–10427, 1999.
40. Horvath TL, Diano S, and Barnstable C. Mitochondrial uncoupling protein 2 in the central nervous system: neuromodulator and neuroprotector. *Biochem Pharmacol* 65: 1917–1921, 2003.
41. Horvath TL, Diano S, Leranthe C, Garcia-Segura LM, Cowley MA, Shanabrough M, Elsworth JD, Sotonyi P, Roth RH, Dietrich EH, Matthews RT, Barnstable CJ, and Redmond DE Jr. Coenzyme Q induces nigral mitochondrial uncoupling and prevents dopamine cell loss in a primate model of Parkinson's disease. *Endocrinology* 144: 2757–2760, 2003.
42. Jaburek M and Garlid KD. Reconstitution of recombinant uncoupling proteins: UCP1, -2, and -3 have similar affinities for ATP and are unaffected by coenzyme Q10. *J Biol Chem* 278: 25825–25831, 2003.
43. Jakus PB, Sipos K, Kispal G, and Sandor A. Opposite regulation of uncoupling protein 1 and uncoupling protein 3 in vivo in brown adipose tissue of cold-exposed rats. *FEBS Lett* 519: 210–214, 2002.
44. Jezek P, Zackova M, Ruzicka M, Skobisova E, and Jaburek M. Mitochondrial uncoupling proteins—facts and fantasies. *Physiol Res* 53 Suppl 1: S199–S211, 2004.
45. Kim-Han JS, Reichert SA, Quick KL, and Dugan LL. BMCP1: a mitochondrial uncoupling protein in neurons which regulates mitochondrial function and oxidant production. *J Neurochem* 79: 658–668, 2001.
46. Klein NA, Houmard BS, Hansen KR, Woodruff TK, Sluss PM, Bremner WJ, and Soules MR. Age-related analysis of inhibin A, inhibin B, and activin a relative to the intercycle monotropic follicle-stimulating hormone rise in normal ovulatory women. *J Clin Endocrinol Metab* 89: 2977–2981, 2004.
47. Klingenberg M. Uncoupling protein—a useful energy dissipator. *J Bioenerg Biomembr* 31: 419–430, 1999.
48. Kondou S, Hidaka S, Yoshimatsu H, Tsuruta Y, Itateyama E, and Sakata T. Molecular cloning of rat brain mitochondrial carrier protein-1 cDNA and its up-regulation during postnatal development. *Biochim Biophys Acta* 1457: 182–189, 2000.

49. Korshunov SS, Skulachev VP, and Starkov AA. High protonic potential actuates a mechanism of production of reactive oxygen species in mitochondria. *FEBS Lett* 416: 15–18, 1997.
50. Krauss S, Zhang CY, Scorrano L, Dalgaard LT, St-Pierre J, Grey ST, and Lowell BB. Superoxide-mediated activation of uncoupling protein 2 causes pancreatic beta cell dysfunction. *J Clin Invest* 112: 1831–1842, 2003.
51. Kunz WS, Goussakov IV, Beck H, and Elger CE. Altered mitochondrial oxidative phosphorylation in hippocampal slices of kainate-treated rats. *Brain Res* 826: 236–242, 1999.
52. Kuwashima J, Nakamura K, Fujitani B, Kadokawa T, Yoshida K, and Shimizu M. Relationship between cerebral energy failure and free fatty acid accumulation following prolonged brain ischemia. *Jpn J Pharmacol* 28: 277–287, 1978.
53. Lazarewicz JW, Strosznajder J, and Gromek A. Effects of ischemia and exogenous fatty acids on the energy metabolism in brain mitochondria. *Bull Acad Pol Sci Biol* 20: 599–606, 1972.
54. Ledesma A, de Lacoba MG, and Rial E. The mitochondrial uncoupling proteins. *Genome Biol* 3: 3015, 2002.
55. Lengacher S, Magistretti PJ, and Pellerin L. Quantitative rt-PCR analysis of uncoupling protein isoforms in mouse brain cortex: methodological optimization and comparison of expression with brown adipose tissue and skeletal muscle. *J Cereb Blood Flow Metab* 24: 780–788, 2004.
56. Li G, Klein RL, Matheny M, King MA, Meyer EM, and Scarpace PJ. Induction of uncoupling protein 1 by central interleukin-6 gene delivery is dependent on sympathetic innervation of brown adipose tissue and underlies one mechanism of body weight reduction in rats. *Neuroscience* 115: 879–889, 2002.
57. MacManus JP, Graber T, Luebbert C, Preston E, Rasquinha I, Smith B, and Webster J. Translation-state analysis of gene expression in mouse brain after focal ischemia. *J Cereb Blood Flow Metab* 24: 657–667, 2004.
58. Mao W, Yu XX, Zhong A, Li W, Brush J, Sherwood SW, Adams SH, and Pan G. UCP4, a novel brain-specific mitochondrial protein that reduces membrane potential in mammalian cells. *FEBS Lett* 443: 326–330, 1999.
59. Mattiasson G, Shamloo M, Gido G, Mathi K, Tomasevic G, Yi S, Warden CH, Castilho RF, Melcher T, Gonzalez-Zulueta M, Nikolich K, and Wieloch T. Uncoupling protein-2 prevents neuronal death and diminishes brain dysfunction after stroke and brain trauma. *Nat Med* 9: 1062–1068, 2003.
60. Mills EM, Xu D, Fergusson MM, Combs CA, Xu Y, and Finkel T. Regulation of cellular oncogenesis by uncoupling protein 2. *J Biol Chem* 277: 27385–27392, 2002.
61. Mitchell P. Coupling of phosphorylation to electron and hydrogen transfer by a chemi-osmotic type of mechanism. *Naturwissenschaften* 191: 144–148, 1961.
62. Miwa S and Brand MD. Mitochondrial matrix reactive oxygen species production is very sensitive to mild uncoupling. *Biochem Soc Trans* 31: 1300–1301, 2003.
63. Negre-Salvayre A, Hirtz C, Carrera G, Cazenave R, Trolly M, Salvayre R, Penicaud L, and Casteilla L. A role for uncoupling protein-2 as a regulator of mitochondrial hydrogen peroxide generation. *FASEB J* 11: 809–815, 1997.
64. Nicholls DG, Snelling R, and Rial E. Proton and calcium circuits across the mitochondrial inner membrane. *Biochem Soc Trans* 12: 388–390, 1984.
65. Nicholls DG, Vesce S, Kirk L, and Chalmers S. Interactions between mitochondrial bioenergetics and cytoplasmic calcium in cultured cerebellar granule cells. *Cell Calcium* 34: 407–424, 2003.
66. Nishikawa T, Edelstein D, Du XL, Yamagishi S, Matsumura T, Kaneda Y, Yorek MA, Beebe D, Oates PJ, Hammes HP, Giardino I, and Brownlee M. Normalizing mitochondrial superoxide production blocks three pathways of hyperglycaemic damage. *Nature* 404: 787–790, 2000.
67. Palmieri F. The mitochondrial transporter family (SLC25): physiological and pathological implications. *Pflugers Arch* 447: 689–709, 2004.
68. Paradis E, Clavel S, Bouillaud F, Ricquier D, and Richard D. Uncoupling protein 2: a novel player in neuroprotection. *Trends Mol Med* 9: 522–525, 2003.
69. Pecqueur C, Alves-Guerra MC, Gelly C, Levi-Meyrueis C, Couplan E, Collins S, Ricquier D, Bouillaud F, and Miroux B. Uncoupling protein 2, in vivo distribution, induction upon oxidative stress, and evidence for translational regulation. *J Biol Chem* 276: 8705–8712, 2001.
70. Phillis JW and O'Regan MH. A potentially critical role of phospholipases in central nervous system ischemic, traumatic, and neurodegenerative disorders. *Brain Res Brain Res Rev* 44: 13–47, 2004.
71. Pichiule P, Chavez JC, and LaManna JC. Oxygen and oxidative stress modulate the expression of uncoupling protein-5 in vitro and in vivo. *Adv Exp Med Biol* 540: 103–107, 2003.
72. Richard D, Rivest R, Huang Q, Bouillaud F, Sanchis D, Champigny O, and Ricquier D. Distribution of the uncoupling protein 2 mRNA in the mouse brain. *J Comp Neurol* 397: 549–560, 1998.
73. Richard D, Clavel S, Huang Q, Sanchis D, and Ricquier D. Uncoupling protein 2 in the brain: distribution and function. *Biochem Soc Trans* 29: 812–817, 2001.
74. Sanchis D, Fleury C, Chomiki N, Goubert M, Huang Q, Neverova M, Gregoire F, Easlick J, Raimbault S, Levi-Meyrueis C, Miroux B, Collins S, Seldin M, Richard D, Warden C, Bouillaud F, and Ricquier D. BMCP1, a novel mitochondrial carrier with high expression in the central nervous system of humans and rodents, and respiration uncoupling activity in recombinant yeast. *J Biol Chem* 273: 34611–34615, 1998.
75. Sivitz WI, Fink BD, and Donohoue PA. Fasting and leptin modulate adipose and muscle uncoupling protein: divergent effects between messenger ribonucleic acid and protein expression. *Endocrinology* 140: 1511–1519, 1999.
76. Skulachev VP. Uncoupling: new approaches to an old problem of bioenergetics. *Biochim Biophys Acta* 1363: 100–124, 1998.
77. Smith AM, Ratcliffe RG, and Sweetlove LJ. Activation and function of mitochondrial uncoupling protein in plants. *J Biol Chem* 279: 51944–51952, 2004.
78. Speakman JR, Talbot DA, Selman C, Snart S, McLaren JS, Redman P, Krol E, Jackson DM, Johnson MS, and Brand MD. Uncoupled and surviving: individual mice with high

- metabolism have greater mitochondrial uncoupling and live longer. *Aging Cell* 3: 87–95, 2004.
79. Stephenson D, Rash K, Smalstig B, Roberts E, Johnstone E, Sharp J, Panetta J, Little S, Kramer R, and Clemens J. Cytosolic phospholipase A2 is induced in reactive glia following different forms of neurodegeneration. *Glia* 27: 110–128, 1999.
 80. Sullivan PG, Rippy NA, Dorenbos K, Concepcion RC, Agarwal AK, and Rho JM. The ketogenic diet increases mitochondrial uncoupling protein levels and activity. *Ann Neurol* 55: 576–580, 2004.
 81. Takahashi Y, Kushiro M, Shinohara K, and Ide T. Dietary conjugated linoleic acid reduces body fat mass and affects gene expression of proteins regulating energy metabolism in mice. *Comp Biochem Physiol B Biochem Mol Biol* 133: 395–404, 2002.
 82. Talbot DA, Lambert AJ, and Brand MD. Production of endogenous matrix superoxide from mitochondrial complex I leads to activation of uncoupling protein 3. *FEBS Lett* 556: 111–115, 2004.
 83. Vincent AM, Olzmann JA, Brownlee M, Sivitz WI, and Russell JW. Uncoupling proteins prevent glucose-induced neuronal oxidative stress and programmed cell death. *Diabetes* 53: 726–734, 2004.
 84. Watanabe A and Hirai A. Two uncoupling protein genes of rice (*Oryza sativa* L.): molecular study reveals the defects in the pre-mRNA processing for the heat-generating proteins of the subtropical cereal. *Planta* 215: 90–100, 2002.
 85. Yamada S, Isojima Y, Yamatodani A, and Nagai K. Uncoupling protein 2 influences dopamine secretion in PC12h cells. *J Neurochem* 87: 461–469, 2003.
 86. Zackova M, Skobisova E, Urbankova E, and Jezek P. Activating omega-6 polyunsaturated fatty acids and inhibitory purine nucleotides are high affinity ligands for novel mitochondrial uncoupling proteins UCP2 and UCP3. *J Biol Chem* 278: 20761–20769, 2003.
 87. Zhang CY, Baffy G, Perret P, Krauss S, Peroni O, Grujic D, Hagen T, Vidal-Puig AJ, Boss O, Kim YB, Zheng XX, Wheeler MB, Shulman GI, Chan CB, and Lowell BB. Uncoupling protein-2 negatively regulates insulin secretion and is a major link between obesity, beta cell dysfunction, and type 2 diabetes. *Cell* 105: 745–755, 2001.

Address reprint requests to:

Laura L. Dugan, M.D.

Department of Medicine

Division of General Medicine/Geriatrics

University of California San Diego

San Diego, CA 92093

E-mail: ladugan@UCSD.edu

Received for publication April 12, 2005; accepted April 13, 2005.

This article has been cited by:

1. T. D. Papkovskaia, K.-Y. Chau, F. Inesta-Vaquera, D. B. Papkovsky, D. G. Healy, K. Nishio, J. Staddon, M. R. Duchon, J. Hardy, A. H. V. Schapira, J. M. Cooper. 2012. G2019S leucine-rich repeat kinase 2 causes uncoupling protein-mediated mitochondrial depolarization. *Human Molecular Genetics* **21**:19, 4201-4213. [[CrossRef](#)]
2. Luping Huang, Tatiana Belousova, Minyi Chen, Gabriel DiMattia, Dajun Liu, David Sheikh-Hamad. 2012. Overexpression of stanniocalcin-1 inhibits reactive oxygen species and renal ischemia/reperfusion injury in mice. *Kidney International* **82**:8, 867-877. [[CrossRef](#)]
3. David B. Ramsden, Philip W.-L. Ho, Jessica W.-M. Ho, Hui-Fang Liu, Danny H.-F. So, Ho-Man Tse, Koon-Ho Chan, Shu-Leong Ho. 2012. Human neuronal uncoupling proteins 4 and 5 (UCP4 and UCP5): structural properties, regulation, and physiological role in protection against oxidative stress and mitochondrial dysfunction. *Brain and Behavior* **2**:4, 468-478. [[CrossRef](#)]
4. Rommy von Bernhardt, Jaime Eugén. 2012. Alzheimer's Disease: Redox Dysregulation As a Common Denominator for Diverse Pathogenic Mechanisms. *Antioxidants & Redox Signaling* **16**:9, 974-1031. [[Abstract](#)] [[Full Text HTML](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]
5. Zhe Ying, Alejandro Covalin, Jack Judy, Fernando Gomez-Pinilla. 2012. Hypothalamic stimulation enhances hippocampal BDNF plasticity in proportion to metabolic rate. *Brain Stimulation*. [[CrossRef](#)]
6. Yao-Chung Chuang, Tsu-Kung Lin, Hsuan-Ying Huang, Wen-Neng Chang, Chia-Wei Liou, Shang-Der Chen, Alice YW Chang, Samuel HH Chan. 2012. Peroxisome proliferator-activated receptors gamma/mitochondrial uncoupling protein 2 signaling protects against seizure-induced neuronal cell death in the hippocampus following experimental status epilepticus. *Journal of Neuroinflammation* **9**:1, 184. [[CrossRef](#)]
7. Zhihua Sun, Jing Zhang, Yingmin Chen, Yunting Zhang, Xuejun Zhang, Hong Guo, Chunshui Yu. 2012. Differential Temporal Evolution Patterns in Brain Temperature in Different Ischemic Tissues in a Monkey Model of Middle Cerebral Artery Occlusion. *Journal of Biomedicine and Biotechnology* **2012**, 1-8. [[CrossRef](#)]
8. Malgorzata Slocinska, Nina Antos-Krzeminska, Grzegorz Rosinski, Wieslawa Jarmuszkiewicz. 2011. Identification and characterization of uncoupling protein 4 in fat body and muscle mitochondria from the cockroach *Gromphadorhina cocquerelliana*. *Journal of Bioenergetics and Biomembranes* **43**:6, 717-727. [[CrossRef](#)]
9. Rocío Guevara, Magdalena Gianotti, Jordi Oliver, Pilar Roca. 2011. Age and sex-related changes in rat brain mitochondrial oxidative status. *Experimental Gerontology*. [[CrossRef](#)]
10. M.-T. Besson, P. Dupont, Y.-W. C. Fridell, J.-C. Lievens. 2010. Increased energy metabolism rescues glia-induced pathology in a *Drosophila* model of Huntington's disease. *Human Molecular Genetics* **19**:17, 3372-3382. [[CrossRef](#)]
11. Sic L. Chan, Zelan Wei, Srinivasulu Chigurupati, Weihong Tu. 2010. Compromised respiratory adaptation and thermoregulation in aging and age-related diseases. *Ageing Research Reviews* **9**:1, 20-40. [[CrossRef](#)]
12. Zhaofei Wu, Jie Zhang, Baolu Zhao. 2009. Superoxide Anion Regulates the Mitochondrial Free Ca²⁺ Through Uncoupling Proteins. *Antioxidants & Redox Signaling* **11**:8, 1805-1818. [[Abstract](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)] [[Supplemental material](#)]
13. S. Sharma, Y. Zhuang, Z. Ying, A. Wu, F. Gomez-Pinilla. 2009. Dietary curcumin supplementation counteracts reduction in levels of molecules involved in energy homeostasis after brain trauma. *Neuroscience* **161**:4, 1037-1044. [[CrossRef](#)]
14. Andrew Chi-Yuen Chu, Philip Wing-Lok Ho, Ken Hon-Hung Kwok, Jessica Wing-Man Ho, Koon-Ho Chan, Hui-Fang Liu, Michelle Hiu-Wai Kung, David Boyer Ramsden, Shu-Leong Ho. 2009. Mitochondrial UCP4 attenuates MPP⁺ and dopamine-induced oxidative stress, mitochondrial depolarization, and ATP deficiency in neurons and is interlinked with UCP2 expression. *Free Radical Biology and Medicine* **46**:6, 810-820. [[CrossRef](#)]
15. Tetsuro Matsunaga, Ning Gu, Hanae Yamazaki, Mariko Tsuda, Tetsuya Adachi, Koichiro Yasuda, Toshio Moritani, Kinsuke Tsuda, Masahiko Nonaka, Tohru Nishiyama. 2009. Association of UCP2 and UCP3 polymorphisms with heart rate variability in Japanese men. *Journal of Hypertension* **27**:2, 305-313. [[CrossRef](#)]
16. Fernando Gomez-Pinilla, Shoshanna Vaynman, Zhe Ying. 2008. Brain-derived neurotrophic factor functions as a metabotrophin to mediate the effects of exercise on cognition. *European Journal of Neuroscience* **28**:11, 2278-2287. [[CrossRef](#)]
17. Matthew T. Scharf, Nirinjini Naidoo, John E. Zimmerman, Allan I. Pack. 2008. The energy hypothesis of sleep revisited. *Progress in Neurobiology* **86**:3, 264-280. [[CrossRef](#)]

18. Ulrich Dirnagl, Andreas Meisel. 2008. Endogenous neuroprotection: Mitochondria as gateways to cerebral preconditioning?. *Neuropharmacology* **55**:3, 334-344. [[CrossRef](#)]
19. B. Karaszewski, J. M. Wardlaw, I. Marshall, V. Cvorovic, K. Wartolowska, K. Haga, P. A. Armitage, M. E. Bastin, M. S. Dennis. 2008. Early brain temperature elevation and anaerobic metabolism in human acute ischaemic stroke. *Brain* **132**:4, 955-964. [[CrossRef](#)]
20. L. Li, K. Prabhakaran, X. Zhang, L. Zhang, H. Liu, J.L. Borowitz, G.E. Isom. 2008. 1#,25-dihydroxyvitamin D3 attenuates cyanide-induced neurotoxicity by inhibiting uncoupling protein-2 up-regulation. *Journal of Neuroscience Research* **86**:6, 1397-1408. [[CrossRef](#)]
21. Jullie W. Pan, Anne Williamson, Idil Cavus, Hoby P. Hetherington, Hitten Zaveri, Ognen A. C. Petroff, Dennis D. Spencer. 2008. Neurometabolism in human epilepsy. *Epilepsia* **49**:s3, 31-41. [[CrossRef](#)]
22. X. Zhang, L. Li, K. Prabhakaran, L. Zhang, H.B. Leavesley, J.L. Borowitz, G.E. Isom. 2007. Uncoupling protein-2 up-regulation and enhanced cyanide toxicity are mediated by PPAR# activation and oxidative stress. *Toxicology and Applied Pharmacology* **223**:1, 10-19. [[CrossRef](#)]
23. S CHEN, H WU, D YANG, S LEE, F SHAW, T LIN, C LIOU, Y CHUANG. 2006. Effects of rosiglitazone on global ischemia-induced hippocampal injury and expression of mitochondrial uncoupling protein 2. *Biochemical and Biophysical Research Communications* **351**:1, 198-203. [[CrossRef](#)]
24. Tatjana Haitina, Jonas Lindblom, Thomas Renström, Robert Fredriksson. 2006. Fourteen novel human members of mitochondrial solute carrier family 25 (SLC25) widely expressed in the central nervous system. *Genomics* **88**:6, 779-790. [[CrossRef](#)]
25. Philip Wing-Lok Ho, Andrew Chi-Yuen Chu, Ken Hon-Hung Kwok, Michelle Hiu-Wai Kung, David Boyer Ramsden, Shu-Leong Ho. 2006. Knockdown of uncoupling protein-5 in neuronal SH-SY5Y cells: Effects on MPP+-induced mitochondrial membrane depolarization, ATP deficiency, and oxidative cytotoxicity. *Journal of Neuroscience Research* **84**:6, 1358-1366. [[CrossRef](#)]
26. Barbara Cannon, Irina G. Shabalina, Tatiana V. Kramarova, Natasa Petrovic, Jan Nedergaard. 2006. Uncoupling proteins: A role in protection against reactive oxygen species—or not?. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1757**:5-6, 449-458. [[CrossRef](#)]
27. Janet M. Dubinsky . 2005. CNS Mitochondria in Neurodegenerative Disorders. *Antioxidants & Redox Signaling* **7**:9-10, 1089-1091. [[Citation](#)] [[Full Text PDF](#)] [[Full Text PDF with Links](#)]