

The Neuroprotection of Hydrogen Sulfide Against MPTP-Induced Dopaminergic Neuron Degeneration Involves Uncoupling Protein 2 Rather Than ATP-Sensitive Potassium Channels

Ming Lu,¹ Fang-Fang Zhao,¹ Juan-Juan Tang,¹ Cun-Jin Su,¹ Yi Fan,¹ Jian-Hua Ding,¹
Jin-Song Bian,^{1,2} and Gang Hu¹

Abstract

Aims: Hydrogen sulfide (H₂S), a novel gaseous mediator, has been recognized to protect neurons from over-excitation by enhancing the activity of the adenosine triphosphate-sensitive potassium (K-ATP) channel. However, no direct evidence supports that the K-ATP channel contributes to the neuroprotective effect of H₂S in neurodegeneration. Herein, wild-type and Kir6.2 knockout (*Kir6.2*^{-/-}) mice were used to establish the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) mouse model of Parkinson's disease (PD) so as to investigate the involvement of K-ATP channels in the neuroprotection of H₂S. **Results:** Systemic administration of sodium hydrosulfide (NaHS) (an H₂S donor, 5.6 mg/kg/day) for 7 days rescued MPTP-induced loss of dopaminergic (DA) neurons in substantia nigra compacta of both *Kir6.2*^{+/+} and *Kir6.2*^{-/-} mice. Consistently, NaHS (100 μM) protected primary mesencephalic neurons against 1-methyl-4-phenylpyridinium (MPP⁺)-induced cytotoxicity in both genotypes. We further found that deficiency of mitochondrial uncoupling protein 2 (UCP2), which reduces reactive oxygen species (ROS) production and functions as upstream to the K-ATP channel in determining vulnerability of DA neurons, abolished the protective effects of H₂S against either DA neuron degeneration in the PD mouse model or MPP⁺-induced injury in primary mesencephalic neurons. Rationally, UCP2 evokes mild uncoupling, which in turn diminishes ROS accumulation in DA neurons. Furthermore, H₂S exerted neuroprotective effect *via* enhancing UCP2-mediated antioxidation and subsequently suppressing ROS-triggered endoplasmic reticulum stress as well as ultimately inhibiting caspase 12-induced neuronal apoptosis. **Innovation and Conclusion:** H₂S protects DA neurons against degeneration in a UCP2 rather than Kir6.2/K-ATP channel-dependent mechanism, which will give us an insight into the potential of H₂S in terms of opening up new therapeutic avenues for PD. *Antioxid. Redox Signal.* 17, 849–859.

Introduction

PARKINSON'S DISEASE (PD) is characterized by the selective degeneration of dopaminergic (DA) neurons in the substantia nigra (SN) and aggregation of Lewy bodies in neurons. The incidence of the disorder increases with age, and more than 2% of the population aged over 65 are attacked by the disease. Therefore, PD is the second most prevalent neurodegenerative disease with only symptomatic treatment available (18). Clinically, the first-class drug to treat PD is L-DOPA, which solely ameliorates the symptoms but fails to retard DA neuron degeneration. It thus is crucial to develop effective therapeutic drugs and strategies that can reverse or

alleviate the pathological process occurring in the DA neurons in PD patients.

Over the last decade, hydrogen sulfide (H₂S) has been recognized as a novel gaseous mediator alongside nitric oxide and carbon monoxide (1). Endogenous H₂S in mammalian body is produced by cystathionine β-synthase and cystathionine γ-lyase, which are two types of pyridoxal-5'-phosphate-dependent enzymes. Later on, Shibuya *et al.* found that 3-mercaptopyruvate sulfurtransferase, in combination with cysteine aminotransferase, also produces H₂S with cysteine as a substrate in neurons (27). H₂S is present at a relatively high level in brain of human, rat, and bovine (15). Furthermore, multiple lines of evidence suggest that H₂S may

¹Jiangsu Key Laboratory of Neurodegeneration, Department of Pharmacology, Nanjing Medical University, Nanjing, Jiangsu, People's Republic of China.

²Department of Pharmacology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore, Singapore.

Innovation

We here demonstrate that hydrogen sulfide (H_2S) protects dopaminergic (DA) neurons against 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced degeneration in an adenosine triphosphate-sensitive potassium channel-independent, but uncoupling protein 2 (UCP2)-dependent mechanism. Accordingly, we propose that H_2S may enhance UCP2-induced mild uncoupling of mitochondrial oxidative phosphorylation. As such, reactive oxygen species generation is reduced, which in turn ameliorates oxidative products evoked endoplasmic reticulum stress and subsequent neuronal apoptosis. Ultimately, H_2S alleviates the loss of substantia nigra compacta DA neurons and protects against MPTP-induced neurodegeneration. These findings will give us an insight into the potential of H_2S in terms of opening up new therapeutic avenues for Parkinson's disease (PD). Promisingly, H_2S -based neuroprotective therapies against PD may one day be developed.

function as a neuromodulator in the brain. In peripheral systems, H_2S protects heart against ischemic injury, regulates vascular tone, controls the release of insulin, and modulates cell proliferation and apoptosis (2, 9, 29). Although the contribution is minor in comparison to the effect on increasing glutathione levels, H_2S protects neurons from excessive excitation by enhancing the activity of adenosine triphosphate-sensitive potassium (K-ATP) channels (12, 13). However, no direct evidence supports that the K-ATP channel is responsible for the neuroprotective effect of H_2S *in vivo*.

K-ATP channels are unique channel proteins that directly couple the metabolic state of a cell to its electrical activity and distribute widely in brain. They are heterooctamers composed of pore-forming Kir6.x (6.1 or 6.2) subunits and sulfonylurea receptor (SUR1 or SUR2) regulatory subunits (25–26). Kir6.2 and SUR1 or SUR2B are expressed in neurons, whereas Kir6.1 and SUR1 or SUR2 are in astrocyte and microglia (31, 32, 36). H_2S stimulates the K-ATP channel in the neurons, vascular smooth muscle cells, cardiomyocytes, and pancreatic β -cells, exerting its biological function. The direct evidence was from electrophysiological investigations since H_2S increased K-ATP channel currents, which were blocked by antagonists like glibenclamide, in rat aorta and arteria mesenterica. H_2S may bind to the cysteine residues (Cys6 and Cys26) of SUR1 or break the disulfide bonds between them. This will change the steric configuration and therefore open K-ATP channels (8, 19, 30). Nevertheless, the exact association of H_2S and K-ATP channels is still in need of further exploration.

Although we reported that H_2S may be a potential neuroprotectant for PD therapy (6), it remains largely unknown whether this neuroprotection of H_2S is mediated by K-ATP channels. Therefore, we established a PD model by injection of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP), a classic neurotoxin that destroys DA neurons, in both *Kir6.2^{+/+}* and *Kir6.2^{-/-}* mice to investigate the involvement of K-ATP channels in the neuroprotection of H_2S and to gain an insight into the potential strategy for PD therapy. Herein, we found unexpectedly that H_2S protected DA neurons against MPTP-induced degeneration in a K-ATP channel-independent manner. However, mitochondrial mild uncoupling has been shown to predominantly reduce generation of reactive oxy-

gen species (ROS), and decreased ROS will reduce the open probability of K-ATP channels. Therefore, uncoupling protein 2 (UCP2) may function as the upstream of the K-ATP channel in determining vulnerability of DA neurons (3). Our study showed that UCP2 mediated the protective effects of H_2S against either DA neuron degeneration in the PD mouse model or MPP⁺-injured primary mesencephalic neurons. These findings indicate that the UCP2 rather than K-ATP channel is responsible for the neuroprotection of H_2S .

Results

NaHS protects DA neurons against MPTP-induced degeneration in a Kir6.2/K-ATP channel-independent manner

To demonstrate whether K-ATP channels contribute to the effect of H_2S on DA neuron degeneration, we established an MPTP-induced subacute model of PD in wild type (WT) and *Kir6.2^{-/-}* mice. As shown in Figure 1a, administration of MPTP (20 mg/kg, s.c.) for 5 days resulted in the dramatic degeneration of tyrosine-hydroxylase-positive (TH^+) neurons in substantia nigra compacta (SNc) and ventral tegmental area (VTA). There was no significant difference in the number of TH^+ neurons between WT and *Kir6.2^{-/-}* mice in response to MPTP treatment. Treatment with sodium hydrosulfide (NaHS) at 5.6 mg/kg for 7 days remarkably rescued the loss of TH^+ neurons induced by MPTP in both genotypic mice. Stereological statistics revealed that NaHS almost fully recovered the MPTP-reduced TH^+ cell numbers in SNc and VTA of both *Kir6.2^{+/+}* and *Kir6.2^{-/-}* mice to the basal level (Fig. 1b, c). These results indicate that H_2S may exert a beneficial effect on DA neuron degeneration in a Kir6.2/K-ATP channel-independent manner.

Depletion of nigrostriatal dopamine is an important pathological character for PD. To examine the effect of H_2S on DA neurotransmitters in the MPTP-induced PD mouse model, we measured the levels of dopamine and its metabolic products by high-performance liquid chromatography coupled with electrochemical detection (HPLC-ECD). We found that MPTP treatment resulted in a dramatic reduction of dopamine (Fig. 1d), dihydroxyphenylacetic acid (DOPAC; Fig. 1e), and homovanillic acid (HVA; Fig. 1f) in striatum of both genotypes. However, NaHS (5.6 mg/kg), which alone had no significant effect on DA or its metabolic products, failed to rescue the decreased levels of neurotransmitters caused by MPTP. These data imply that H_2S may ameliorate the structural degeneration of DA neuron before the rescue of dopamine metabolism and release.

NaHS protects primary mesencephalic neurons against MPP⁺-induced cytotoxicity in a Kir6.2/K-ATP channel-independent manner

To investigate the potential mechanisms for the protective effect of H_2S on PD, we isolated and cultured the primary mesencephalic neurons from WT and *Kir6.2^{-/-}* mice. Immunocytochemistry showed that the number and morphology of TH^+ neurons were equivalent under basal condition in both genotypes (Fig. 2a). Incubation of MPP⁺ (10 μM) for 48 h visibly decreased the mean length of neuritis and the number of TH^+ neurons (Fig. 2a, b) and increased lactate dehydrogenase (LDH) release (Fig. 2c). Pretreatment with NaHS (100 μM ,




FIG. 1. NaHS protected DA neurons against MPTP-induced degeneration in midbrain of wild type and *Kir6.2*^{-/-} mice. Immunohistochemical staining of THir neurons in SNc of mice with $\times 40$ magnifications (**a**). Stereological counts of THir cells in mouse SNpc (**b**) and VTA (**c**). Data are presented as the mean \pm S.E.M., $n = 5$ for each group. NaHS had no significant effect on MPTP-reduced levels of DA (**d**), DOPAC (**e**), and HVA (**f**) in the striatum of both genotypic mice. Data expressed as means \pm S.E.M. from eight mice per group. * $p < 0.05$, ** $p < 0.01$, and *** $p < 0.001$ versus saline group in corresponding genotype; # $p < 0.05$ and ## $p < 0.01$ versus corresponding MPTP-treated groups. MPTP, 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; DA, dopaminergic; TH, tyrosine-hydroxylase; TH⁺, tyrosine-hydroxylase positive; THir, tyrosine-hydroxylase immunoreactive; VTA, ventral tegmental area; DOPAC, dihydroxyphenylacetic acid; HVA, homovanillic acid; NaHS, sodium hydrosulfide; SNc, substantia nigra compacta; SNpc, substantia nigra pars compacta; S.E.M., standard error of mean. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).

15 min) significantly attenuated MPP⁺-reduced number of TH⁺ neurons and suppressed the MPP⁺-elevated LDH level to the same extent in both genotypes (Fig. 2a–c). These results further reveal that the protective effect of H₂S on DA neuron degeneration *in vitro* is also independent of the Kir6.2/K-ATP channel.

NaHS inhibits endoplasmic reticulum stress-triggered apoptosis in primary mesencephalic neurons isolated from WT and *Kir6.2*^{-/-} mice

Glucose-regulated protein (GRP) 78, which induces expression in endoplasmic reticulum (ER) lumen and also acts as an apoptotic regulator during endoplasmic reticulum stress (ERS), has been thought to be a biomarker of ERS. As shown in Figure 2d, preincubation with NaHS (100 μ M) for 15 min significantly suppressed the upregulation of GRP78 induced by MPP⁺ in both genotypes (Fig. 2e). Furthermore, the inductions of caspase 12 and p65 represent the activation level of ERS-triggered death pathway and inflammation, respectively. NaHS suppressed MPP⁺-upregulated expression of

caspase 12 (Fig. 2f) and the translocation of p65 into nucleus (Fig. 2g) to a similar extent in both genotypic mesencephalic neurons. These data suggest that H₂S may protect DA neurons against degeneration *via* inhibiting ERS-triggered apoptosis and inflammation.

NaHS attenuates MPTP-triggered glial cell activation and proliferation in the presence or absence of the Kir6.2/K-ATP channel

Glial proliferation and overactivation occur during the process of PD. Accompanied by the sustained secretion of inflammatory mediators, the persistent activation of microglia is regarded to be involved in the aggravation of DA neuron degeneration in PD. We therefore examined glial activation in the SNc of mice. As shown in Figure 3, the basal numbers of Glial fibrillary acidic protein (GFAP⁺, a marker for astrocytes) and Mac-1⁺ (also called CD11b/C18, a marker for microglia activation) cells were identical in SNc of WT and *Kir6.2*^{-/-} mice. Administration of MPTP significantly increased the numbers of GFAP⁺ (426% and 587%, WT




FIG. 2. NaHS protected against MPP⁺-induced cytotoxicity in primary cultured mesencephalic neurons from wild-type and *Kir6.2*^{-/-} mice. (a) Typical photograph of immunocytochemical staining showing TH⁺ neurons at ×100 magnifications. (b, c) Effects of NaHS on the number of TH⁺ neurons (b) and LDH release (c) in MPP⁺-treated DA neurons. (d) Typical western blotting revealed that NaHS alleviated ERS and apoptosis in primary mesencephalic neurons isolated from wild-type and *Kir6.2*^{-/-} mice. Statistical analysis of GRP78 (e), caspase 12 (f), and p65/H₃ (g) protein levels in neurons. Data are presented as the mean ± S.E.M. of four individual experiments. ***p* < 0.01 and **p* < 0.05 versus control group; ##*p* < 0.01 and #*p* < 0.05 versus corresponding MPP⁺-treated groups; \$*p* < 0.05 versus *Kir6.2*^{-/-} group. ERS, endoplasmic reticulum stress; GRP, glucose-regulated protein; LDH, lactate dehydrogenase; MPP⁺, 1-methyl-4-phenylpyridinium. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).

and *Kir6.2*^{-/-}, respectively) and Mac-1⁺ cells (271% and 261%, WT and *Kir6.2*^{-/-}, respectively) in SNc compared with those in saline-treated mice. NaHS treatment attenuated the proliferation of GFAP⁺ (306% and 378% of vehicle, WT and *Kir6.2*^{-/-}, respectively) (Fig. 3a, b) and Mac-1⁺ cells (167% and 177% of vehicle, WT, and *Kir6.2*^{-/-}, respectively) (Fig. 3c, d). These data suggest that low expression of Kir6.2 in astrocytes may be important to maintain the resting status of the glial cells and Kir6.2 knockout aggravated astrocyte activation. Moreover, inhibition of MPTP-induced glial excessive activation and subsequent neuroinflammation also contributes to the potential protective effect of H₂S on PD.

UCP2 deficiency abolishes the protective effects of NaHS against DA neuron degeneration in vivo and in vitro

UCP2, located in the inner membrane of mitochondria, reduces ROS production in the respiratory chain and functions as the upstream to K-ATP channels in determine the vulnerability of DA neurons (3). To identify the potential target of H₂S, we investigated the effect of H₂S in MPTP-treated WT and *Ucp2*^{-/-} mice. Immunostaining showed that MPTP induced obvious TH⁺ neuron loss in SNc of WT and *Ucp2*^{-/-} mice. Notably, NaHS treatment could reverse SNc TH⁺ neuron damage in WT mice (Fig. 4a, b). However, UCP2

knockout abolished the protective effect of NaHS against MPTP-injured TH⁺ neurons. Similar results were also exhibited in the VTA region (Fig. 4c). *In vitro* study showed that MPP⁺ significantly reduced TH⁺ cell number and neurites length (Fig. 5a–c) as well as elevated LDH release (Fig. 5d) in the neurons isolated from WT and *Ucp2*^{-/-} mice. In line with what we found in the *in vivo* study, NaHS treatment could only markedly attenuate MPP⁺-induced neuronal injuries in WT mice, but not in *Ucp2*^{-/-} mice. These results suggest that UCP2 may be the action target of H₂S and contribute to the neuroprotective effects of H₂S against DA neuron degeneration.

UCP2 knockout revokes the prevention of NaHS on ERS-triggered neuronal apoptosis in the mesencephalon

It has been demonstrated that ERS involves DA neuron degeneration in PD and that expression of C/EBP homologous protein (CHOP), which is an ERS-induced transcription factor, can be stimulated by MPTP in the PD mouse model. In this study, we found that treatment with NaHS for 7 days reversed the upregulated protein expression of GRP78, CHOP, and caspase 12 in the mesencephalon of WT mice treated with MPTP. However, these suppressive effects of NaHS on ERS were abolished by UCP2 knockout (Fig. 6a, b).




FIG. 3. NaHS attenuated MPTP-triggered glial cell activation and proliferation in the SN of wild-type and *Kir6.2*^{-/-} mice. (a) Microphotographs of GFAP-ir cells in the SN with $\times 40$ magnification. (b) Stereological counts of GFAP-ir cells in mouse SN. (c) Microphotographs of Mac1-ir cells in the SN with $\times 200$ magnification. (d) Stereological counts of Mac-1-ir cells in mouse SN. $n = 5$. Data are presented as the mean \pm S.E.M., ** $p < 0.01$ and * $p < 0.05$ versus saline group; # $p < 0.05$ versus corresponding MPTP-treated groups; \$ $p < 0.05$ versus corresponding MPTP-treated *Kir6.2*^{-/-} groups. SN, substantia nigra; GFAP, Glial fibrillary acidic protein. Arrows indicate Mac-1⁺ cells. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).

These results suggest that H₂S may suppress ERS-triggered apoptosis owing to the existence of UCP2.

UCP2 deletion abolishes the inhibitory effect of NaHS on ROS accumulation in TH⁺ neurons in SN

Given that intracellular ROS accumulation evokes ERS, we continued to examine ROS production in TH⁺ neurons in the presence or absence of UCP2 to explore the potential mechanism whereby UCP2 deficiency abolished the effect of H₂S on DA neuron degeneration. It was found that MPTP triggered ROS accumulation not only in TH⁺ neurons but also in astrocytes surrounding TH⁺ neurons in our previous study (data not shown). As shown in Figure 6c, ROS levels were much higher in SNc TH⁺ neurons of *Ucp2*^{-/-} mice treated with and without MPTP compared with those in WT mice, respectively. Moreover, H₂S treatment significantly scavenged the accumulation of ROS in WT TH⁺ neurons, but failed to diminish ROS level in TH⁺ neurons of *Ucp2*^{-/-} mice. These findings indicate that existence of UCP2 suppresses intracellular ROS production and mediates the antioxidative role of H₂S, whereas UCP2 deletion increases mitochondrial ROS production and further abolishes the protective effects of H₂S.

NaHS fails to alleviate MPTP-triggered glial cell activation and proliferation in the absence of UCP2

As shown in Figure 7, there was no significant difference of the morphology and basal amounts of GFAP⁺ as well as Mac-1⁺ cells in SNc between WT and *Ucp2*^{-/-} mice. Administration of MPTP resulted in a significant activation and proliferation of GFAP⁺ and Mac-1⁺ cells in SNc compared with those in saline-treated mice. NaHS treatment attenuated the MPTP-enhanced proliferation of GFAP⁺ (Fig. 7a, b) and Mac-1⁺ cells (Fig. 7c, d) in WT mice. However, NaHS failed to alleviate MPTP-induced glial cell proliferation in the SNc of *Ucp2*^{-/-} mice. These data suggest that UCP2-mediated antioxidation involves the inhibitory effect of NaHS on glial cell activation and the subsequent neuroinflammation.

Discussion

We demonstrated in the present study that the Kir6.2/K-ATP channel, which is abundantly expressed in SN DA neurons, was not responsible for the neuroprotective effects of H₂S in the MPTP mouse model of PD. Deficiency of UCP2, which functions as the upstream of K-ATP channels (3),




FIG. 4. UCP2 deficiency abolished the protective effect of NaHS on MPTP-injured TH⁺ neurons in SNc and VTA. (a) Microphotographs of THir neurons in SNpc of mice with $\times 40$ magnification. Stereological counts of THir cells in SNpc (b) and VTA (c). Data are presented as the mean \pm S.E.M., $n=5$ for each group. * $p<0.05$ and ** $p<0.01$ versus saline group in corresponding genotype; # $p<0.05$ versus corresponding MPTP-treated *Ucp2*^{+/+} groups; \$ $p<0.05$ versus corresponding *Ucp2*^{+/+} groups. UCP2, uncoupling protein 2. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).

abolished the protective effect of H₂S on DA neuronal degeneration. Our findings indicate that H₂S may target UCP2 to attenuate ROS-triggered ERS and subsequent neuronal apoptosis, suggesting the potential of H₂S in prospective therapy for PD.

Among numerous ion channels, the K-ATP channel is a unique one to integrate energy cues with membrane excitability-dependent processes (17, 21). It is inhibited by ATP at the physiologic concentration range, but stimulated when ATP falls during metabolic depletion (28). K-ATP channels contributed to multiple pathological and physiological functions of H₂S (30). However, they were not found to mediate the antioxidant effects of H₂S in the central neurons system (33). The present study was, therefore, designed to investigate the contribution of K-ATP channels in the therapeutic effects of H₂S in the MPTP-induced subacute model of PD with *Kir6.2*^{-/-} mice. Although Birgit Liss *et al.* reported that *Kir6.2* knockout recovered DA neuron loss in SNc in the chronic PD model (16), we found that deficiency of *Kir6.2* had no effect on DA neuron degeneration in both SNc and VTA in the subacute MPTP-induced PD model. Notably, our results showed that H₂S reversed the loss of DA neurons induced by MPTP in both WT and *Kir6.2*^{-/-} mice. *In vitro* study further revealed that H₂S protected mesencephalic TH⁺ neurons against MPP⁺-induced cytotoxicity in both genotypes. These findings indicate that the beneficial effect of H₂S on PD model mice is independent of the *Kir6.2*/K-ATP channel. In agreement of previous findings (16), we found that dopamine production and survival of DA neurons seemed out of step. H₂S failed to rescue MPTP-induced depletion of dopamine and its metabolic products in

striatum of both genotypic mice at current experiment condition, suggesting that although NaHS rescued DA neuron loss in SNc, the TH in these survived neurons may not able to produce DA synchronously. This is consistent with the previous finding that structural recovery of DA neuron degeneration is before the amelioration of neuronal function *in vivo* (37).

So far, the exact mechanisms for DA neuron degeneration in the progress of PD have not been fully understood. It involves multiple pathological impairments such as mitochondrial oxidative stress, inflammation, dysfunction of ubiquitin-protease system, and deposition of abnormal protein. Among these, ER is the major signal-transducing organelle that continuously responds to environmental cues (22, 24) and provides a unique oxidizing compartment for the folding of membrane and secretory proteins (11). Recent studies reveal that ER stress also occurs in the neurodegeneration of PD and MPTP evokes the accumulation of unfolding or misfolding protein and up-regulate expression of GRP78, which in turn initiates the unfolding protein response (4, 23, 38). Recently, Krishnan *et al.* reported that suppression of endogenous H₂S production decreased the phosphorylation of protein kinase-like endoplasmic reticulum kinase, thus reducing its activation in response to ER stress (14). In the present study, we found consistently that H₂S attenuated MPTP-induced ER stress and the subsequent apoptosis. Moreover, absence of the *Kir6.2*/K-ATP channel could not impair the suppressive effect of H₂S on ER stress in MPP⁺-injured DA neurons. These data suggest that the K-ATP channel is not responsible for the neuroprotective effect of H₂S in the MPTP-induced PD model.




FIG. 5. UCP2 knockout abolished the protective effect of NaHS on MPP⁺-induced neurotoxicity in primary cultured mesencephalic neurons. THir neurons were stained by immunocytochemistry with $\times 200$ magnifications (a). Effects of NaHS on MPP⁺-induced neurotoxicity in DA neurons as assessed by the number of TH⁺ neurons (b), mean neurites length per cell (c), and LDH release (d). Four independent experiments were performed in duplicate. Data were presented as the mean \pm S.E.M. *** $p < 0.001$ and ** $p < 0.01$ versus control group; ## $p < 0.01$ and # $p < 0.05$ versus MPTP-treated *Ucp2*^{+/+} group; \$ $p < 0.05$ *Ucp2*^{+/+} group versus *Ucp2*^{-/-} group. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).

UCP2, localized in the inner mitochondrial membrane, uncouples respiration from ATP synthesis by providing an alternative route for the protons to enter into the mitochondrial matrix and also functions as a transporter of fatty acids into the mitochondrial matrix (20). The primary physiological function of mild uncoupling is to suppress oxygen radical production accompanied by restricting mitochondrial permeability transition pore opening and Ca²⁺ accumulation (20). It also plays an important role in regulating the energy balance in many tissues such as pancreas, liver, and brain (10). It has been reported that UCP2 modulated intracellular energy metabolism and thus was involved in aging, diabetes, and neurodegeneration (5, 34). It has been proposed that UCP2 may be upstream to the K-ATP channel in determining vulnerability of DA neurons (3). We therefore identified whether UCP2 was a potential target for the action of H₂S by establishing an MPTP PD model in *Ucp2*^{-/-} mice. Our findings showed that UCP2 knockout could abolish the protective effects of H₂S against DA neuron damage in both *in vivo* and *in vitro* studies, indicating that UCP2 may be responsible for the neuroprotective action of H₂S.

MPTP is converted to MPP⁺ by monoamine oxidase-B expressed in glial cells and then transported into DA neurons by dopamine transporter. MPP⁺ selectively inhibits mitochondrial complex I, triggering ROS production and ERS. As a result, upregulated GRP78 and CHOP activate gene transcription, leading to glial cell activation-induced neuroinflammation or caspase 12-mediated neuronal apoptosis. H₂S may act on UCP2 and facilitate it to inhibit MPTP-induced ROS accumulation and subsequently triggered ERS, whereby preventing DA neurons from degeneration (Fig. 8). UCP2 deficiency impairs the inhibitory action on mitochondrial ROS production and further leads to the accumulation of higher level oxygen radicals. Consequently, this deficit may largely weaken or abolish the antioxidative effect of H₂S on both neurons and glial cells. These findings suggest that UCP2 may act as a potential target of H₂S on protecting DA neurons *via* antioxidation.

Although our study revealed the neuroprotection of H₂S was independent of Kir6.2/K-ATP channels in the PD mouse model, we does not exclude the possibility that K-ATP channels contribute to the cytoprotective actions of H₂S in other pathophysiological processes. Since UCP2 uncouples oxidative




FIG. 6. UCP2 mediated the suppressive effect of NaHS on ROS-triggered ERS in the midbrain. Western blotting analysis of GRP78, CHOP, and caspase 12 protein levels in the midbrain of mice (**a**, **b**). UCP2 knockout abolished the inhibitory effect of NaHS on MPTP-induced ROS accumulation in SNc TH⁺ neurons (**c**). Photos were taken at $\times 600$ magnification. Green color represents TH neuron and red color represents ROS. Scale bar = 22 μ m. Data are presented as the mean \pm S.E.M., $n = 4$ for each group. * $p < 0.05$ versus saline group; # $p < 0.05$ versus corresponding MPTP-treated groups. ROS, reactive oxygen species; CHOP, C/EBP homologous protein. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).

phosphorylation and reduces the synthesis of ATP (5, 7), it acts as a potential modulator of the K-ATP channel *via* regulating the ratio of ATP and ADP. H₂S may therefore activate UCP2, which in turn decreases ATP level and subsequently opens K-ATP channels. In this case, UCP2 may be upstream molecule of K-ATP channels and is responsible for the neuroprotective effect of H₂S on PD. Therefore, although it is not essential for the neuroprotection of H₂S in the PD model, the K-ATP channel can still be activated by H₂S and mediates multiple functions in other pathophysiological processes.

Materials and Methods

The study protocol was approved by the Institutional Animal Care and Use Committee of Nanjing Medical University.

Animals and reagents

Kir6.2 knockout C57BL/6J mice were donated by Professor Miki (Chiba University, Japan) (25), of which 12–16-weeks-


old, 26–32-g male Kir6.2^{+/+} and Kir6.2^{-/-} mice were used. UCP2 knockout C57BL/6J mice were donated by Professor Chen-Yu Zhang (Nanjing University, P.R. China) with the same age and weight. The expressions of Kir6.2 and UCP2 were detected by RT-PCR (Supplementary Fig. S1; Supplementary Data are available online at www.liebertonline.com/ars).

NaHS and MPTP were purchased from Sigma-Aldrich, while probenecid was from Jinan Times Pharmacology Co. Ltd. MPTP was dissolved in saline and preserved at 0°C, while probenecid was dissolved in dimethyl sulfoxide (DMSO) just 30 min before injection.

Animal model and drug administration

Each type of mice (Kir6.2^{+/+}, Kir6.2^{-/-}, Ucp2^{+/+}, and Ucp2^{-/-}) was divided into four groups: vehicle, MPTP, MPTP+NaHS (5.6 mg kg⁻¹·day⁻¹, intraperitoneal injection [i.p.]), and NaHS alone groups. MPTP (20 mg kg⁻¹, subcutaneous [s.c.]) was injected 1 h before injection of probenecid

FIG. 7. UCP2 deletion abolished the inhibitory effect of NaHS on MPTP-triggered glial cell activation and proliferation. (a) Microphotographs of GFAP-ir cells in the SN with $\times 40$ magnification. (b) Stereological counts of GFAP-ir cells in mouse SN. (c) Microphotographs of Mac1-ir cells in the SN with $\times 200$ magnification. (d) Stereological counts of Mac-1-ir cells in mouse SN. $n=4$. Data are presented as the mean \pm S.E.M. $**p < 0.01$ versus saline group; $^{\#}p < 0.05$ versus corresponding MPTP-treated groups; $^{\$}p < 0.05$ and $^{\$\$}p < 0.01$ versus corresponding *Ucp2*^{-/-} groups. Arrows indicate Mac-1⁺ cells. Scale bar = 100 μ m. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).



(250 mg kg⁻¹, i.p.) for 5 days in MPTP group. Mice in NaHS treatment group received NaHS (5.6 mg kg⁻¹, i.p.) injection for 7 days and MPTP administration 30 min after NaHS injection from day 3 to day 7. In NaHS alone group, NaHS (5.6 mg kg⁻¹, ip) was given for 7 days. The same volume of

saline was injected in the 7 days in the vehicle group. Animals were sacrificed 3.5 days after the last injection of MPTP.

Immunohistochemical studies and quantitative evaluation

Immunostaining method was described in the previous publication (6). Images were observed and photos were taken under a confocal microscope (Axiovert LSM510; Carl Zeiss Co.). The immunostaining signals were quantitatively analyzed using the Optical Fractionator method with Microbrightfield Stereo-Investigator software (Stereo Investigator software; Microbrightfield). The total number of TH-immunoreactive (IR) neurons, GFAP-IR astrocytes, and Mac-1-IR microglia in entire extent of substantia nigra pars compacta were counted from four samples per group.

High-performance liquid chromatography

Striatum tissues were collected for measurement of DA, DOPAC, and HVA with HPLC with ECD. HPLC/ECD consisted of BAS LC-4C, a reversed-phase C18 column (Ultrasphere ODS 4.6 \times 250 mm, 5 μ m), chromatograph interface DA-5, and solvent delivery system. The mobile phase consisted of 0.1 M citrate, 0.075 M Na₂HPO₄, 0.1 M EDTA, 1.0 mM 1-heptanesulfonic acid, and 10% methanol, pH 3.9.

In situ detection of ROS

Dihydroethidium (Molecular Probes) was used to investigate the local *in situ* production of ROS. An intravenous injection of 200 μ l dihydroethidium (DHE; stock solution, 100 mg/ml in DMSO, diluted 1:100 with sterile saline before injection) was administered through the caudal vein. Three hours after DHE injection, mice were overdosed with dimethyl ether and transcardially perfused with 4% paraformaldehyde (0.1% glutaraldehyde and 15% picric acid in phosphate buffer). After postfixing overnight in paraformaldehyde without glutaraldehyde, sections at 30 μ m were cut




FIG. 8. Schematic model for the mechanism of H₂S in protecting against MPP⁺-induced neuronal injury. MPP⁺ is selectively transported into DA neurons by DAT and subsequently inhibits the activity of mitochondrial complex I. Accumulated ROS evokes the upregulation of GRP78 and further triggers ERS. Consequently, p65 nuclear translocation-induced gene transcription or caspase 12-mediated apoptosis occurs in DA neurons. H₂S-stimulated mitochondria uncoupling reduces the ratio of ATP/ADP, which in turn opens K-ATP channels. On the other hand, H₂S enhances UCP2 function and attenuates ROS production, suppressing ERS and apoptosis induced by MPP⁺. DAT, dopamine transporter; K-ATP, adenosine triphosphate-sensitive potassium; H₂S, hydrogen sulfide. (To see this illustration in color the reader is referred to the web version of this article at www.liebertonline.com/ars).

through the SN using a Leica freezing microtome. To observe ROS in SNc DA cells, sections were incubated overnight with TH antibody (1:3000). After washing for three times, sections were then incubated with the fluorescent secondary antibody, donkey anti-mouse IgG 488 Alexafluore (1:200; Molecular Probes), to visualize TH-IR cells.

Identification and quantification of TH⁺ neurons and neuronal processes

Mesencephalic neuron primary culture was performed according to the previous publication (35). After incubation with MPP⁺ (10 μ M) for 48 h with or without pretreatment of NaHS (100 μ M), cells were rinsed carefully with 0.1 M phosphate buffer solution (PBS) (pH 7.2) and fixed with 4% paraformaldehyde, followed by blocking with PBST (0.03% Triton) containing 10% bovine serum albumin. Cells were then incubated at 4°C overnight with the primary antibody (the mouse monoclonal anti-TH at 1:3000; Sigma). After washing for three times, cells were exposed to 1:800 dilutions of HRP-conjugated goat anti-mouse IgG (Chemicon) for 1 h at room temperature. Immunostaining was visualized with 3,3'-diaminobenzidine. The number of THir neurons was counted in 10 randomly selected fields (1.13 mm²/field) at $\times 100$ magnification on a Nikon Optical TE2000-S inverted microscope. The values were normalized to that obtained from control culture. The average number of THir cells in control cultures ranged from 20 to 30 per field, while THir cells made up about 5% of all cells in primary culture. Cell processes were determined at $\times 200$ magnification. Photomicrographs were taken with a Nikon Optical TE2000-S inverted microscope. Each THir cell process was traced from soma to the end of the process, realized by the measurement function of Image Pro Plus 5.1.

Assay of LDH release

After incubation with MPP⁺ for 48 h, LDH release was measured in the culture medium with an LDH diagnostic kit (Jiancheng Bioengineering) in accordance with the manufacturer's instructions. LDH activity was calculated by data measured from absorbance at 490 nm. Data were expressed as percentages of LDH release gained from control treatment.

Western blotting analysis

The method was described in previous publication (6). Different primary antibodies (mouse anti-Actin, CHOP, and H3 at 1:1000 [Cell Signaling]; rabbit anti-GRP78 and Caspase 12 at 1:1000 and rabbit anti-p65 at 1:600 [Cell Signaling]) and the secondary antibody (1:1000) were used in the present study.

Statistical analysis

All data were presented as mean \pm standard error of mean. Statistical significance was assessed with one-way analysis of variance followed by the *post hoc* Student-Newman-Keuls test between control and samples treated with various factors. Differences with *p*-values less than 0.05 were considered statistically significant.

Acknowledgments

We appreciate Prof. Miki (Chiba University, Japan) for donating *Kir6.2*^{-/-} mice and appreciate Prof. Chen-Yu Zhang

(Nanjing University, P.R. China) for donating *Ucp2*^{-/-} mice. The work reported herein was supported by the grants from the National Key Program of Basic Research of China (No. 2011CB504103), the National Natural Science Foundation of China (No. 81030060 and No. 30873055), the National Science & Technology Major Project (No. 2012ZX09304-001), and Jiangsu Province's Key Discipline of Medicine (XK201117).

Author Disclosure Statement

The authors declare no conflicts of interest.

References

1. Abe K and Kimura H. The possible role of hydrogen sulfide as an endogenous neuromodulator. *J Neurosci* 16: 1066–1071, 1996.
2. Bannenberg GL and Vieira HL. Therapeutic applications of the gaseous mediators carbon monoxide and hydrogen sulfide. *Expert Opin Ther Pat* 19: 663–682, 2009.
3. Deutch AY and Winder DG. A channel to neurodegeneration. *Nat Med* 12: 17–18, 2006.
4. Holtz WA and O'Malley KL. Parkinsonian mimetics induce aspects of unfolded protein response in death of dopaminergic neurons. *J Biol Chem* 278: 19367–19377, 2003.
5. Hong Y, Fink BD, Dillon JS, and Sivitz WI. Effects of adenoviral overexpression of uncoupling protein-2 and -3 on mitochondrial respiration in insulinoma cells. *Endocrinology* 142: 249–256, 2001.
6. Hu LF, Lu M, Tiong CX, Dawe GS, Hu G, and Bian JS. Neuroprotective effects of hydrogen sulfide on Parkinson's disease rat models. *Aging Cell* 9: 135–146, 2010.
7. Hurtaud C, Gelly C, Chen Z, and Levi-Meyrueis C, Bouilaud F. Glutamine stimulates translation of uncoupling protein 2mRNA. *Cell Mol Life Sci* 64: 1853–1860, 2007.
8. Jiang B, Tang G, Cao K, Wu L, and Wang R. Molecular mechanism for H(2)S-induced activation of K(ATP) channels. *Antioxid Redox Signal* 12: 1167–1178, 2010.
9. Kapoor A and Thiemermann C. Hydrogen sulfide, neurogenic inflammation, and cardioprotection: a tale of rotten eggs and vanilloid receptors. *Crit Care Med* 38: 728–730, 2010.
10. Kashemsant N and Chan CB. Impact of uncoupling protein-2 overexpression on proinsulin processing. *J Mol Endocrinol* 37: 517–526, 2006.
11. Kaufman RJ. Stress signaling from the lumen of the endoplasmic reticulum: coordination of gene transcriptional and translational controls. *Genes Dev* 13: 1211–1233, 1999.
12. Kimura Y, Dargusch R, Schubert D, and Kimura H. Hydrogen sulfide protects HT22 neuronal cells from oxidative stress. *Antioxid Redox Signal* 8: 661–670, 2006.
13. Kimura Y, Goto Y, and Kimura H. Hydrogen sulfide increases glutathione production and suppresses oxidative stress in mitochondria. *Antioxid Redox Signal* 12: 1–13, 2010.
14. Krishnan N, Fu C, Pappin DJ, and Tonks NK. H2S-Induced sulphydration of the phosphatase PTP1B and its role in the endoplasmic reticulum stress response. *Sci Signal* 4: ra86, 2011.
15. Lee SW, Hu YS, Hu LF, Lu Q, Dawe GS, Moore PK, Wong PT, and Bian JS. Hydrogen sulphide regulates calcium homeostasis in microglial cells. *Glia* 54: 116–124, 2006.
16. Liss B, Haeckel O, Wildmann J, Miki T, Seino S, and Roeper J. K-ATP channels promote the differential degeneration of dopaminergic midbrain neurons. *Nat Neurosci* 8: 1742–1751, 2005.
17. Miki T, Liss B, Minami K, Shiuchi T, Saraya A, Kashima Y, Horiuchi M, Ashcroft F, Minokoshi Y, Roeper J, and Seino S. ATP-sensitive K⁺ channels in the hypothalamus are essential for the maintenance of glucose homeostasis. *Nat Neurosci* 4: 507–512, 2001.

18. Moore DJ, West AB, Dawson VL, and Dawson TM. Molecular pathophysiology of Parkinson's disease. *Annu Rev Neurosci* 28: 57–87, 2005.
19. Mustafa AK, Gadalla MM, Sen N, Kim S, Mu W, Gazi SK, Barrow RK, Yang G, Wang R, and Snyder SH. H₂S signals through protein S-sulfhydration. *Sci Signal* 2: ra72, 2009.
20. Nagy TR, Blaylock ML, and Garvey WT. Role of UCP2 and UCP3 in nutrition and obesity. *Nutrition* 20: 139–144, 2004.
21. Nichols CG. KATP channels as molecular sensors of cellular metabolism. *Nature* 440: 470–476, 2006.
22. Ron D and Walter P. Signal integration in the endoplasmic reticulum unfolded protein response. *Nat Rev Mol Cell Biol* 8: 519–529, 2007.
23. Ryu EJ, Harding HP, Angelastro JM, Vitolo OV, Ron D, and Greene LA. Endoplasmic reticulum stress and the unfolded protein response in cellular models of Parkinson's disease. *J Neurosci* 22: 10690–10698, 2002.
24. Schroder M and Kaufman RJ. The mammalian unfolded protein response. *Annu Rev Biochem* 74: 739–789, 2005.
25. Seino S and Miki T. Gene targeting approach to clarification of ion channel function: studies of Kir6.x null mice. *J Physiol* 554: 295–300, 2004.
26. Seino S and Miki T. Physiological and pathophysiological roles of ATP-sensitive K⁺ channels. *Prog Biophys Mol Biol* 81: 133–176, 2003.
27. Shibuya N, Tanaka M, Yoshida M, Ogasawara Y, Togawa T, Ishii K, and Kimura H. 3-Mercaptopyruvate sulfurtransferase produces hydrogen sulfide and bound sulfane sulfur in the brain. *Antioxid Redox Signal* 11: 703–714, 2009.
28. Sun XL and Hu G. ATP-sensitive potassium channels: a promising target for protecting neurovascular unit function in stroke. *Clin Exp Pharmacol Physiol* 37: 243–252, 2010.
29. Szabo C. Hydrogen sulphide and its therapeutic potential. *Nat Rev Drug Discov* 6: 917–935, 2007.
30. Tang G, Wu L, and Wang R. Interaction of hydrogen sulfide with ion channels. *Clin Exp Pharmacol Physiol* 37: 753–763, 2010.
31. Thomzig A, Pruss H, and Veh RW. The Kir6.1-protein, a pore-forming subunit of ATP-sensitive potassium channels, is prominently expressed by giant cholinergic interneurons in the striatum of the rat brain. *Brain Res* 986: 132–138, 2003.
32. Thomzig A, Wenzel M, Karschin C, Eaton MJ, Skatchkov SN, Karschin A, and Veh RW. Kir6.1 is the principal pore-forming subunit of astrocyte but not neuronal plasma membrane K-ATP channels. *Mol Cell Neurosci* 18: 671–690, 2001.
33. Tyagi N, Moshal KS, Sen U, Vacek TP, Kumar M, Hughes WM, Jr., Kundu S, and Tyagi SC. H₂S protects against methionine-induced oxidative stress in brain endothelial cells. *Antioxid Redox Signal* 11: 25–33, 2009.
34. Vogler S, Goedde R, Mitterski B, Gold R, Kroner A, Koczan D, Zettl UK, Rieckmann P, Epplen JT, and Ibrahim SM. Association of a common polymorphism in the promoter of UCP2 with susceptibility to multiple sclerosis. *J Mol Med (Berl)* 83: 806–811, 2005.
35. Xie J, Duan L, Qian X, Huang X, Ding J, and Hu G. K(ATP) channel openers protect mesencephalic neurons against MPP⁺-induced cytotoxicity via inhibition of ROS production. *J Neurosci Res* 88: 428–437, 2010.
36. Zhou F, Yao HH, Wu JY, Ding JH, Sun T, and Hu G. Opening of microglial K(ATP) channels inhibits rotenone-induced neuroinflammation. *J Cell Mol Med* 12: 1559–1570, 2008.
37. Zhou QY and Palmiter RD. Dopamine-deficient mice are severely hypoactive, adipsic, and aphagic. *Cell* 83: 1197–1209, 1995.
38. Zhu JH, Horbinski C, Guo F, Watkins S, Uchiyama Y, and Chu CT. Regulation of autophagy by extracellular signal-regulated protein kinases during 1-methyl-4-phenylpyridinium-induced cell death. *Am J Pathol* 170: 75–86, 2007.

Address correspondence to:

Dr. Gang Hu

Jiangsu Key Laboratory of Neurodegeneration

Department of Pharmacology

Nanjing Medical University

140 Hanzhong Road

Nanjing

Jiangsu 210029

People's Republic of China

E-mail: ghu@njmu.edu.cn

Dr. Jin-Song Bian

Department of Pharmacology

National University of Singapore

10 Medical Drive

Singapore 117597

Singapore

E-mail: phcbjs@nus.edu.sg

Date of first submission to ARS Central, December 31, 2011; date of final revised submission, February 21, 2012; date of acceptance, February 22, 2012.

Abbreviations Used

CHOP = C/EBP homologous protein

DA = dopaminergic

DHE = dihydroedithium

DMSO = dimethyl sulfoxide

DOPAC = dihydroxyphenylacetic acid

ECD = electrochemical detection

ERS = endoplasmic reticulum stress

GFAP = glial fibrillary acidic protein

GRP = glucose-regulated protein

HPLC = high-performance liquid chromatography

H₂S = hydrogen sulfide

HVA = homovanillic acid

IR = immunoreactive

K-ATP = adenosine triphosphate-sensitive potassium

LDH = lactate dehydrogenase

MPP⁺ = 1-methyl-4-phenylpyridinium

MPTP = 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine

NaHS = sodium hydrosulfide

PBS = phosphate buffer solution

PD = Parkinson's disease

ROS = reactive oxygen species

S.E.M. = standard error of mean

SN = substantia nigra

SNc = substantia nigra compacta

TH = tyrosine-hydroxylase

UCP2 = uncoupling protein 2

VTA = ventral tegmental area

WT = wild type